

Questionnaire nr. 1

1. * CA 125 increase most frequently in
 - a) pancreatic cancer
 - b) lung cancer
 - ☒ c) ovarian cancer
 - d) breast cancer
 - e) colorectal cancer

C
2. * Chemotherapy-induced anemia is
 - ☒ a) normocyte normochrome
 - b) microcyte hypochrome
 - c) macrocyte hyperchrome
 - d) microcyte normochrome
 - e) macrocyte normochrome

A
3. * Which of the following drugs is a topoisomerase inhibitor
 - a) cyclophosphamide
 - b) docetaxel
 - c) doxorubicin
 - d) oxaliplatin
 - ☒ e) irinotecan

E
4. * Which of the following drugs is NOT chemotherapy
 - a) cisplatin
 - b) paclitaxel
 - c) exemestane
 - d) oxaliplatin
 - e) etoposide

C
5. * Anemia occurring during chemotherapy is based on a mechanism
 - a) of iron deficiency
 - b) of B12 deficiency
 - c) of tumor-induced bone marrow inhibition
 - d) of treatment-induced bone marrow inhibition
 - e) autoimmune

D
6. * Which of the following tyrosine kinase inhibitors is used in chronic myeloid leukemia
 - a) sunitinib
 - ☒ b) imatinib
 - c) sorafenib
 - d) erlotinib
 - e) lapatinib

B
7. * What is the most common histology of breast cancer
 - a) invasive lobular carcinoma
 - ☒ b) invasive ductal carcinoma
 - c) Phyllodes tumor
 - d) Paget's disease
 - e) angiosarcoma

B
8. * Which of the following is NOT a risk factor for esophageal cancer
 - a) alcohol
 - b) Plummer-Vinson syndrome
 - c) smoking
 - ☒ d) sun exposure
 - e) Barrett's esophagus

D
9. * Risk factors in pancreatic cancer are the following, except
 - a) smoking
 - b) alcohol
 - c) obesity
 - d) Peutz-Jeghers syndrome
 - ☒ e) personal history of liver cancer

E
10. * Which of the following is NOT a risk factor in lung cancer
 - a) smoking
 - ☒ b) diet
 - c) environmental exposure to radon
 - d) radiation exposure
 - e) occupational exposure to asbestos

B

11. * In which type of testicular cancer may be used radiation therapy?

- a) only in Yolk sac tumors
- b) only in embryonal carcinoma
- ☒ c) only in seminomas
- d) only in teratomas
- e) in all histological types of testicular cancer

C

12. * CA 19-9 increases most frequently in

- a) lung cancer
- b) ovarian cancer
- c) breast cancer
- d) thyroid cancer
- ☒ e) pancreatic cancer

E

13. * Which is the first most common clinical sign of breast cancer

- ☒ a) breast lump
- b) retracted nipple
- c) nipple discharge
- d) carcinomatous mastitis
- e) "orange peel" skin

A

14. * The most common type of liver cancer is

- a) cholangiocarcinoma
- b) clear cell carcinoma
- ☒ c) hepatocellular carcinoma
- d) angiosarcoma
- e) lymphoma

C

15. * Which of the following is the most common histology in renal cancer

- a) transitional cell carcinoma
- b) papillary adenocarcinoma
- ☒ c) clear cell adenocarcinoma
- d) chromofobic adenocarcinoma
- e) embryonal carcinoma

C

16. Which of the following viruses have oncogenic capacity

- ☒ a) Epstein Barr virus
- ☒ b) hepatitis C virus
- ☒ c) HPV strain 16
- d) influenza virus
- ☒ e) hepatitis B virus

ABCE

17. CBC of a cancer patient under chemotherapy treatment often present

- ☒ a) anemia
- b) polycythemia
- ☒ c) thrombocytopenia
- d) neutrophilia
- ☒ e) neutropenia

ACE

18. Which of the following markers are used in colorectal cancer

- a) AFP
- ☒ b) CEA
- c) CA 15-3
- d) CA 125
- ☒ e) CA 19-9

BE A?

19. Ultrasound

- a) has the advantage that it is not invasive
- ☒ b) may be carried out even in pregnancy
- c) is the method of choice in the diagnosis of lung cancer
- d) has a higher resolution than MRI in liver cancer
- ☒ e) can differentiate between a cystic tumor and a solid tumor

ABE

20. Endoscopic investigations have the following advantages

- a) allow biopsy sampling
- b) are not invasive
- c) are easily tolerated by the patient
- ☒ d) visualize the epithelium of hollow organs
- e) never require sedation

AD

21. Core needle biopsy

- ☒ a) is used in breast cancer
- ☒ b) is carried out with special needles
- ☒ c) is used in prostate cancer
- ☒ d) can be carried out under ultrasound guidance
- ☐ e) allow only a limited cytological diagnosis

ABCD

22. The TNM staging

- ☒ a) there are two distinct types of staging - clinical staging and pathological staging
- ☒ b) clinical staging is based on results obtained by microscopic examination of the tumor
- ☒ c) the clinical and pathological staging are always identical
- ☒ d) clinical staging sometimes differ from the pathological staging
- ☒ e) pathological staging is based on imaging investigations

AD

23. Which of the following cancers are more common in men

- ☒ a) lung cancer
- ☒ b) thyroid cancer
- ☒ c) gastric cancer
- ☒ d) esophageal cancer
- ☒ e) bladder cancer

ACDE

24. Which statements are true regarding occupational cancers

- ☒ a) are becoming more common lately
- ☒ b) asbestos can cause cervical carcinoma
- ☒ c) dyes and paints industry workers frequently develop bladder cancer
- ☒ d) exposure to benzene compounds may develop leukemia
- ☒ e) exposure to radiation increases the risk of leukemia

BCDE

25. Brachytherapy is

- ☐ a) an antalgic neurolysis procedure for brachial plexus
- ☒ b) an intra-arterial chemotherapy technique on isolated upper limb
- ☒ c) a procedure in which the tumor gets in contact with a radioactive source
- ☒ d) a radiotherapy technique
- ☐ e) a method of implantation of a chemotherapy infusion device

CD

26. Chemotherapy performed before curative surgery is called

- ☐ a) primary chemotherapy
- ☐ b) adjuvant chemotherapy
- ☒ c) neoadjuvant chemotherapy
- ☐ d) palliative chemotherapy
- ☐ e) accelerated chemotherapy

Ac

27. Which of the following statements are true about gastric cancer

- ☒ a) approximately 95% of gastric cancers are adenocarcinomas
- ☒ b) diffuse adenocarcinoma is located on the distal stomach
- ☒ c) intestinal adenocarcinoma is located on the distal stomach
- ☒ d) diffuse adenocarcinoma has a better prognosis
- ☒ e) intestinal adenocarcinoma has a better prognosis

ACE

28. Which statements are true about colorectal cancer

- ☐ a) is rarely curable, even in early stages
- ☐ b) incidence decreases with age
- ☐ c) physical activity and obesity does not influence the risk of developing colorectal cancer
- ☒ d) smokers have an increased risk
- ☒ e) familial adenomatous polyposis (FAP) is associated with an increased risk

DE

29. Which statements are true about liver cancer

- ☒ a) malignant tumors of the liver can be both primary tumors and secondary tumors
- ☐ b) liver cancer is one of the rarest cancers worldwide
- ☐ c) occurs exclusively in females
- ☒ d) hepatocellular carcinoma is the most common form
- ☒ e) overall 5 year survival rate is close to 50%

AD

30. Symptoms that may be encountered in patients with esophageal cancer are

- ☒ a) regurgitation
- ☒ b) dysphagia
- ☐ c) vomiting
- ☒ d) weight loss
- ☐ e) hematuria

ABD

31. Clinical examination in patients with hepatocellular carcinoma can detect

- ☒ a) hepatomegaly
- ☒ b) splenomegaly
- ☒ c) abdominal collateral circulation
- ☒ d) ascites
- ☐ e) tumor on digital rectal examination

ABCD

32. Which of the following histologies are characteristic for testicular cancer

- ☐ a) "signet ring" carcinoma
- ☒ b) Yolk sac tumor
- ☒ c) choriocarcinoma
- ☒ d) embryonal carcinoma
- ☐ e) transitional cell carcinoma

BCD

33. Renal adenocarcinoma is

- ☐ a) sensitive to chemotherapy
- ☒ b) resistant to chemotherapy
- ☐ c) sensitive to radiation therapy
- ☒ d) resistant to radiation therapy
- ☐ e) sensitive to antiangiogenic therapies

BD

34. Carcinoembryonic antigen is frequently elevated in

- ☒ a) colorectal cancer
- ☒ b) lung cancer
- ☒ c) breast cancer
- ☐ d) head and neck cancer
- ☐ e) bacterial meningitis

ABC

35. The following statements are true about MRI

- ☐ a) is based on the use of X-rays
- ☒ b) is based on the use of a magnetic field
- ☒ c) is contraindicated in the presence of various metal devices or pacemakers
- ☒ d) is more efficient than tomography in the evaluation of brain or pelvic lesions
- ☐ e) the time to perform MRI is very short, several seconds

BCD

36. A number of microscopic criteria suggest malignancy

- ☒ a) variation in cell size and shapes
- ☐ b) low mitotic activity
- ☒ c) variable shaped nuclei
- ☒ d) organized arrangement of cells
- ☒ e) increased mitotic activity

ACE

37. Malignant tumor grading systems

- ☒ a) appreciate the degree of tissue differentiation
- ☐ b) consider the number of mitoses
- ☐ c) determine the size of the primary tumor
- ☒ d) consider nuclear pleomorphism
- ☐ e) consider the existence of distant metastases

ABD

38. Follow-up of cancer patients

- ☒ a) detect early relapse
- ☐ b) is called screening
- ☐ c) is called primary prevention
- ☒ d) is called secondary prevention
- ☒ e) is called tertiary prevention

AE

39. Which of the following statements are true about brachytherapy

- ☒ a) in brachytherapy radiation sources are placed in intimate contact with the tumor tissue
- ☐ b) surface brachytherapy is used in prostate cancer
- ☒ c) intracavity implants are used in cervical cancer
- ☒ d) afterloading techniques protect medical staff
- ☐ e) interstitial brachytherapy is used for skin cancer

ACD

40. Which of the following antiemetic drugs are serotonin antagonists

- ☒ a) ondansetron
- ☐ b) aprepitant
- ☐ c) metoclopramide
- ☐ d) palonosetron
- ☐ e) prochlorperazine

AD

41. Which of the following are risk factors in ...
- a) late-onset menstrual cycle and early menopause
 - ☒ b) oral contraceptives
 - c) obesity
 - ☒ d) alcohol
 - e) coming from a poor country

42. Trastuzumab

- ☒ a) is an anti-HER2 monoclonal antibody
- b) is an anti-CD20 monoclonal antibody
- c) is an anti-VEGF monoclonal antibody
- ☒ d) may be associated with chemotherapy
- ☒ e) is used in certain types of breast cancer

43. Which statements are true about the prognosis of esophageal cancer

- ☒ a) esophageal cancer has a good prognosis, 5-year survival rate is over 40%
- ☒ b) TNM stage is the most important prognostic factor
- ☒ c) Weight loss of more than 10% of body weight is accompanied by a worse prognosis
- d) liver metastases does not influence prognosis
- ☒ e) tumor in the lower and middle third has a worse prognosis

44. Which of the following are paraneoplastic dermatologic syndromes

- ☒ a) acquired ichthyosis
- b) syndrome of inappropriate ADH secretion
- c) Eaton-Lambert syndrome
- ☒ d) acanthosis nigricans
- ☒ e) myasthenia gravis

45. Nausea and vomiting caused by chemotherapy can be

- ☒ a) acute during the first 24 hours of treatment
- ☒ b) delayed, starting 24 hours after treatment
- c) constant intensity throughout the chemotherapy
- ☒ d) anticipatory, before administration
- ☒ e) chemotherapy never induce nausea and vomiting

36. A number of microscopic criteria suggest malignancy ... in cell size and shapes

46. In breast cancer, castration can be done
- ☒ a) surgically
 - ☒ b) by irradiation
 - c) chemically using aromatase inhibitors
 - ☒ d) chemically using GnRH analogues
 - ☒ e) chemically using antiestrogens

47. Risk factors for liver cancer are

- ☒ a) HBV infection
- ☒ b) HCV infection
- ☒ c) cirrhosis
- ☒ d) ingestion of aflatoxin contaminated food
- ☒ e) sun exposure

48. Bleeding in colorectal cancer may be

- ☒ a) one of the first clinical signs of colorectal cancer
- ☒ b) in the form of melena
- ☒ c) in the form of fresh blood in stool
- ☒ d) in the form of occult hemorrhage
- ☒ e) in the form of haematemesis

49. The following statements regarding the treatment of colorectal cancer are true

- ☒ a) chemotherapy is useful only in patients with metastatic disease
- ☒ b) radiotherapy is useful in the treatment of rectal cancer
- ☒ c) combination of chemotherapy and monoclonal antibodies prolongs survival in patients with metastatic colorectal cancer
- ☒ d) commonly used combination chemotherapy is based on 5-FU or capecitabine
- ☒ e) radiotherapy is useful in the treatment of proximal colon cancer

50. Which of these drugs are tyrosine kinase inhibitors for use in renal cancer

- ☒ a) sunitinib
- ☒ b) temsirolimus
- c) bevacizumab
- ☒ d) everolimus
- ☒ e) sorafenib