



UN SINGUR RASPUNS:

Single Answer

1. Which of the following etiologies is the most frequent in kidney cancer

- a) transitional carcinoma
- b) papillary adenocarcinoma
- ☒ c) clear cell adenocarcinoma
- d) cromophobe adenocarcinoma
- e) embrionic carcinoma

2. The most frequent neoplasm of prostate metastasis:

- a) hepatic
- b) cerebral
- ☒ c) bone
- d) lung
- e) skin

3. Which of the following is not a negative prognostic element for a patient with bronchopulmonary adenocarcinoma

- a) Unintentional weight loss of 10 kg in the last 6 months
- ☒ b) EGFR mutation present
- c) Performance index ECOG 3
- d) Contralateral lung nodule
- e) Hepatic metastasis

4. What is the frequency of hormone dependent breast cancers

- a) Less than 5%
- b) 20-30%
- c) 40-50%
- ☒ d) 60-70%
- e) Over 90%

5. Which is the principal risk factor in cervical cancer

- a) Prolonged exposure to estrogen
- b) Prolonged exposure to sun rays
- c) Unprotected sexual contact
- ☒ d) Infection with HP
- e) Diet rich in animal proteins and fats

6. The most frequent type of hepatic cancer is

- ☒ a) Hepatocarcinoma
- b) Cholangiocarcinoma
- c) Hepatoblastoma
- d) Angiosarcoma
- e) Lymphoma

7. Paraneoplastic syndromes that can be found in gastric cancer are

- a) acanthosis nigricans
- b) dermatomyositis
- c) Trousseau Syndrome
- d) Nephrotic Syndrome
- ☒ e) All of the above



8. Which of the following affirmations about biochemical resistance to chemotherapy is not true

- a) The tumor can be incapable of converting the antineoplastic agent to its active form
- b) Pleiotropic resistance is given by a type of membrane pump cellular structure
- c) the cell may increase the level of target enzymes that the cytotoxic agent intends to inactivate
- d) resistance is due to deficit in irrigation of malignant tissue
- ☒ e) the cell may increase the expression of DNA repair enzymes

9. Which is the most frequent clinical sign at debut of breast cancer

- ☒ a) tumoral breast nodule
- b) axillary adenopathy
- c) nipple leakage
- d) carcinomatous mastitis
- e) Peau D'Orange

10. Which of the following medications are antitumoral antibiotics

- a) mustard nitrogen analogues
- b) nitrosureas
- c) purine analogs
- ☒ d) anthracyclines
- e) taxanes

11. Which of the following antiemetic medications is a neurokinin 1 antagonist (NK-1)

- a) Ondansetron
- b) Aprepitant

11. Which of the following antiemetic medications is a neurokinin 1 antagonist (NK-1)

- a) Ondansetron
- ☒ b) Aprepitant
- c) Metoclopramide
- d) Palonosetron
- e) Prochlorperazine

12. Which of the following medications is a cytostatic

- a) Tamoxifen
- b) Paclitaxel
- ☒ c) Tramstuzumab
- d) Imatinib
- e) Exemestan

13. Bortezomib is

- a) Inhibitor of mTOR kinase
- b) Monoclonal antibody
- c) Tirokinase inhibitor
- d) A cytokine
- ☒ e) Proteosome inhibitor

14. CA 19-9 levels rise most frequently in

- a) Lung cancer
- b) Ovarian cancer
- c) Breast cancer
- d) Thyroid cancer
- ☒ e) Pancreatic cancer



15. Aromatase inhibitors

- ☒ a) Reduce the level of estrogen through the blockage of synthesis
- ☐ b) Reduce the level of estrogen through the binding of estrogen receptors
- ☐ c) Reduce the level of estrogen through the binding of estrogen-progesterone receptors
- ☒ d) Reduce the level of estrogen through chemical castration
- ☐ e) Reduce the level of estrogen through chemical castration and binding of estrogen receptors

MULTIPLE RESPONSES:

16. Carcinoembryonic antigens have frequent values ... raised in individuals

- ☒ a) with colorectal cancer
- ☒ b) with lung cancer
- ☒ c) with hepatic cirrhosis
- ☐ d) non-smokers
- ☐ e) with bacterial meningitis

17. The following affirmations are true about MRI

- ☐ a) Are based on the usage of x rays
- ☒ b) Are based on the property of hydrogen atoms in a magnetic field
- ☒ c) Is contraindicated in the presence of metallic deposits and pacemakers
- ☒ d) Is more effective than CT in evaluating brain or pelvic lesions
- ☐ e) The time of an MRI is very short

18. A series of criteria that microscopically suggests malignancy

18. A series of criteria that microscopically suggests malignancy

- ☒ a) Cellular pleomorphism
- ☐ b) Decreased mitotic activity
- ☒ c) Nuclear pleomorphism
- ☐ d) Conservation of tissue architecture

19. Immunohistochemistry

- ☐ a) Is utilized only for testicular tumors...
- ☒ b) Permits the detection of certain antigens expressed at the cytoplasmic or nuclear membrane level
- ☒ c) Permits the phenotyping of the tumor and its classification into a certain tumor type
- ☐ d) Is a complementary exam useful in tumor anaplasia with a non-precise point of exit
- ☐ e) Requires an initial scintigraphic exam

20. Grading of malignant tumors:

- ☒ a) Appreciates cellular anaplasia
- ☒ b) Appreciates the number of mitoses
- ☐ c) Appreciates the size of the primary tumor
- ☒ d) Appreciates the nuclear pleomorphism
- ☐ e) Appreciates the existence of distant metastasis

21. The following are examples of analytical epidemiological studies

- ☐ a) Incidence studies
- ☐ b) descriptive studies
- ☒ c) cohort studies
- ☒ d) case-control studies

22. Which of the following statements are true about dietary risk factors:

- a) Consumption of vegetables and fruits increases the risk of digestive cancers
- ☒ b) Vegetable glucan may increase the immunity of the intestinal wall
- ☒ c) Phytoestrogens have a protective effect for breast and prostate cancer
- d) Olive oil is an oncological risk factor
- e) Fungal aflatoxins are the main risk factor for lung cancer

23. Post-therapeutic follow-up

- a) is also called screening
- ☒ b) is also called follow-up
- c) is also called primary prevention
- d) is also called secondary prevention
- ☒ e) is also called tertiary prevention

24. The role of surgery in the diagnosis and staging of malignant tumors

- a) is lately increasingly important as new staging techniques develop
- ☒ b) is performed as the initial procedure in all digestive neoplasms
- c) allows surgical biopsy
- ☒ d) allows the subsequent tumorous excision to be performed in the same anesthetic session
- e) is the procedure that sets the stage of TNM in AML

25. Reconstructive surgery in the oncological patient

- ☒ a) improves survival
- ☒ b) repairs the defects
- ☒ c) Improves quality of life
- ☒ d) Restores functionality of affected organs or segments
- ☒ e) Reduces tumor mass

26. Which of the following statements are true about brachytherapy:

- ☒ a) In brachytherapy, radiation sources are in intimate contact with tumor tissue
- ☒ b) Endocavitary brachytherapy is especially used in prostate cancer
- ☒ c) Contact brachytherapy is used in cervical cancer
- ☒ d) Afterloading techniques protect medical staff from irradiation
- ☒ e) Interstitial brachytherapy is used in oral cavity cancer

27. The benefits of neoadjuvant chemotherapy are:

- ☒ a) reducing the likelihood of subsequent systemic recurrence
- ☒ b) is done after removal of the primary tumor and allows for obtaining microscopic information about the tumor
- ☒ c) allows for evaluation of treatment efficiency by measuring the primary tumor size
- ☒ d) reducing the mass of the primary tumor and facilitating surgical progression
- ☒ e) is done at low doses that do not cause immediate or late complications

28. The chemotherapy combinations have the following principles:

- ☒ a) substances that generate mechanisms differing in resistance

28. The chemotherapy combinations have the following principles:

- ☒ a) substances that generate mechanisms differing in resistance
- ☐ b) substances that act through the same mechanism
- ☒ c) substances with different adverse effects
- ☐ d) substances that act at different times in the cell cycle
- ☒ e) substances that act through different mechanisms

29. Thrombocytopenia during chemotherapy:

- ☒ a) is rarely severe or life-threatening
- ☒ b) Determine treatment if the platelets are $<100,000 / \text{mm}^3$
- ☐ c) Determine treatment if the platelets are $<10,000 / \text{mm}^3$
- ☐ d) Determine treatment if the platelets are $<50,000 / \text{mm}^3$
- ☒ e) Can be corrected with platelets

30. Which of the following are risk factors for breast cancer:

- ☐ a) menstrual cycle with late onset and early menopause
- ☒ b) Oral contraceptives
- ☒ c) Obesity
- ☒ d) Alcohol
- ☐ e) coming from a poor country

31. Trastuzumab:

- ☒ a) is an anti-HER2 monoclonal antibody
- ☐ b) is an anti-CD20 monoclonal antibody
- ☐ c) is an anti-VEGF monoclonal antibody
- ☒ d) may be associated with chemotherapy
- ☐ e) is indicated in metastatic colorectal cancer

32. Which of the following statements about the prognosis of

colorectal cancer are true:

32. Which of the following statements about the prognosis of esophageal cancer are true:

- a) Esophageal cancer has a good prognosis, survival at 5 years being $> 40\%$
- ☒ b) TNM status is the most important prognostic factor
- ☒ c) Decreasing of $> 10\%$ of body weight is associated with a worse prognosis
- d) The presence of liver metastases does not influence prognosis
- e) The location of the tumor at the lower and middle third level has a worse prognosis

33. Prognostic factors in colorectal cancer are:

- a) tumor grade, G1 and G2 tumors with a worse prognosis than G3 and G4
- b) histopathological type, carcinoma with Signet ring cell (?) and mucinous carcinoma having a more reserved prognosis
- ☒ c) the stage of TNM is the most important prognostic element
- ☒ d) lymphatic or venous invasion
- e) residual tumor

34. The risk factors for pancreatic cancer are:

- ☒ a) Smoking
- ☒ b) hyperlipidic diet
- ☒ c) partial gastrectomy
- ☒ d) chronic pancreatitis
- e) Excessive exposure to sunlight



35. Symptoms seen in gastric cancer are as follows:

35. Symptoms seen in gastric cancer are as follows:

- ☒ a) epigastric pain
- ☐ b) anorexia
- ☒ c) early satiety
- ☐ d) food vomiting
- ☐ e) hematuria

36. The negative prognostic factors in pancreatic cancer are:

- ☐ a) old age
- ☐ b) location at head of pancreas
- ☒ c) the advanced stage of the disease
- ☒ d) poorly differentiated tumor
- ☐ e) well-differentiated tumor

37. Which of the following is a risk factor in lung cancer

- ☐ a) liver cirrhosis
- ☒ b) exposure to asbestos
- ☒ c) exposure to radon
- ☐ d) HTA
- ☐ e) DZ

38. Which of the following statements are true of radiotherapy in lung cancer

- ☐ a) Cranial prophylactic CT is performed only in non-small cell cancer patients
- ☒ b) Radical radiotherapy radiates the primary tumor, hilar ganglions and mediastinum
- ☒ c) Palliative RT can be used in upper vena cava syndrome
- ☐ d) Concomitant chemo-radiotherapy brings a benefit to sequential

39. Chemotherapy in prostate cancer

- ☒ a) is usually a solution used late after hormone therapy
- ☒ b) is initial treatment in metastatic prostate cancer
- ☒ c) Docetaxel and Prednisone are used in the first line of chemotherapy
- d) Mitoxantrone can be used in patients who can not tolerate taxane therapy
- e) uses combinations of 3-4 cytostatics that give more effective responses

40. Which of the following histologies is characteristic of a testicular cancer

- a) cell carcinoma with Signet ring cell
- b) Yolk sac tumor
- ☒ c) Coriocarcinoma
- d) transitional carcinoma
- ☒ e) Embryonic carcinoma

41. Which of the following drugs are tyrosine kinase inhibitors used in kidney neoplasm?

- ☒ a) Sunitinib
- b) Temsirolimus
- c) Bevacizumab
- d) everolimus
- ☒ e) sorafenib

42. Which of the following statements are true about endometroid adenocarcinoma:

- a) occurs in 20% of cases of endometrial cancer
- b) has a better prognosis than clear cell adenocarcinoma

42. Which of the following statements are true about endometroid adenocarcinoma:

- a) occurs in 20% of cases of endometrial cancer
- ☒ b) has a better prognosis than clear cell adenocarcinoma
- ☒ c) cells have increased levels of progesterone receptors
- d) affects more often older people
- ☒ e) associated with obesity and diabetes

43. Which of the following statements about systemic therapy in endometrial cancer are true:

- ☒ a) Systemic chemotherapy is indicated in advanced or recurrent forms
- ☒ b) hormone therapy is particularly indicated in well differentiated tumors with positive hormone receptors
- c) hormone therapy is indicated in the adjuvant treatment of endometrial cancer
- ☒ d) hormone therapy is indicated in metastatic or recurrent forms
- e) hormone therapy uses bicalutamide or GnRH analogues

44. Which of the following statements about ovarian cancer are true:

- a) Most ovarian cancers are germ cell tumors
- b) Most ovarian cancers are stromal tumors
- ☒ c) Most ovarian cancers are of epithelial origin
- ☒ d) are usually chemosensitive tumors
- e) usually detected in advanced stages

45. Which of the following statements about ovarian cancer are true:

- a) The most frequent dissemination is peritoneal seeding



45. Which of the following statements about ovarian cancer are true:

- ☒ a) The most frequent dissemination is peritoneal seeding
- ☐ b) the most frequent dissemination is haematogene
- ☒ c) Pulmonary and cerebral metastases rarely occur
- ☐ d) Ovarian cancer is most often detected in early stages, limited to the ovary
- ☐ e) the bone is the most common place of metastasis

46. What surgical procedures can be used in cervical cancer:

- ☒ a) Conization
- ☐ b) Lumpectomy
- ☒ c) radical hysterectomy with bilateral pelvic lymphadenectomy
- ☐ d) sectorectomy
- ☒ e) pelvic exenteration

47. Tumor suppression genes:

- ☐ a) may give rise to protooncogenes by mutations
- ☒ b) control the cell cycle
- ☐ c) are also called protooncogenes
- ☐ d) play a role in carcinogenesis
- ☒ e) The most well-known p53 with the role of genomic guardian

48. Anemia in cancer may be due:

- ☐ a) most often an autoimmune mechanism
- ☒ b) direct inhibition of erythropoiesis by the tumor by cytokine production
- ☒ c) an iron deficiency mechanism through occult hemorrhages
- ☐ d) Edema (?)
- ☐ e) direct MOH (?) infiltration



e) The most well-known p53 with the role of genomic guardian

48. Anemia in cancer may be due:

- a) most often an autoimmune mechanism
- ☒ b) direct inhibition of erythropoiesis by the tumor by cytokine production
- ☒ c) an iron deficiency mechanism through occult hemorrhages
- d) Edema (?)
- e) direct MOH (?) infiltration

49. In hepatic metastases, changes may occur in:

- ☒ a) Transaminases
- ☒ b) Bilirubin
- c) PSA
- d) serum albumin
- ☒ e) clotting tests

50. The following statements regarding the treatment of colorectal cancer are true

- a) Chemotherapy is only useful in patients with unresectable tumors or metastatic disease
- ☒ b) Radiotherapy is useful in the treatment of rectal cancer
- ☒ c) the association of monoclonal antibodies with chemotherapy leads to prolonging the survival of patients with metastatic colorectal cancer
- ☒ d) Chemotherapy usually uses 5-FU or capecitabine-based combinations
- e) radiotherapy is useful in the treatment of cancer and proximal colon cancer

