

# ASSIGNMENT-2

B.ANUSHA

Download all python codes from

<https://github.com/BOJJAVOYINAANUSHA/Assignment-2/blob/main/ASSIGNMENT2/assignment2.py>

and latex-tikz codes from

<https://github.com/BOJJAVOYINAANUSHA/Assignment-2/blob/main/ASSIGNMENT2/main.tex>

## 1 QUESTION No. 2.42

Construct  $ABCD$  where  $AB = 4$ ,  $BC = 5$ ,  $CD = 6.5$ ,  $\angle B = 105^\circ$  and  $\angle C = 80^\circ$ .

## 2 SOLUTION

- 1) Let us assume vertices of given quadrilateral  $ABCD$  as  $\mathbf{A}, \mathbf{B}, \mathbf{C}$  and  $\mathbf{D}$ .
- 2) Let us generalize the given data:

$$\angle B = 105^\circ = \theta \quad (2.0.1)$$

$$\angle C = 80^\circ = \alpha \quad (2.0.2)$$

$$\|\mathbf{A} - \mathbf{B}\| = 4 = p \quad (2.0.3)$$

$$\|\mathbf{C} - \mathbf{B}\| = 5 = q \quad (2.0.4)$$

$$\|\mathbf{D} - \mathbf{C}\| = 6.5 = r \quad (2.0.5)$$

- For this quadrilateral  $ABCD$  we have,

$$\angle B + \angle C = 105^\circ + 80^\circ = 185^\circ \quad (2.0.6)$$

- Let,

$$\mathbf{B} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \mathbf{C} = \begin{pmatrix} 5 \\ 0 \end{pmatrix} \quad (2.0.7)$$

**Lemma 2.1.** *The coordinates of  $\mathbf{A}$  and  $\mathbf{D}$  can be written as follows:*

$$\mathbf{A} = p\mathbf{b} \quad \left( \because \mathbf{B} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right) \quad (2.0.8)$$

$$\mathbf{D} = \mathbf{C} + r\mathbf{c} \quad (2.0.9)$$

Let us define  $\mathbf{b}, \mathbf{c}$  as:

$$\mathbf{b} = \begin{pmatrix} \cos B \\ \sin B \end{pmatrix}, \mathbf{c} = \begin{pmatrix} \cos C \\ \sin C \end{pmatrix} \quad (2.0.10)$$

- For finding coordinates of  $\mathbf{A}$ :-

Putting (2.0.1) and (2.0.3) in (2.0.8) we get,

$$\Rightarrow \mathbf{A} = 4 \begin{pmatrix} \cos 105^\circ \\ \sin 105^\circ \end{pmatrix} \quad (2.0.11)$$

$$\Rightarrow \mathbf{A} = \begin{pmatrix} -1.03 \\ 3.86 \end{pmatrix} \quad (2.0.12)$$

- For finding coordinates of  $\mathbf{D}$ :-

Putting (2.0.2) and (2.0.5) in (2.0.9) we get,

$$\Rightarrow \mathbf{D} = \begin{pmatrix} 5 \\ 0 \end{pmatrix} + 6.5 \begin{pmatrix} \cos 80^\circ \\ \sin 80^\circ \end{pmatrix} \quad (2.0.13)$$

$$\Rightarrow \mathbf{D} = \begin{pmatrix} 5 \\ 0 \end{pmatrix} + \begin{pmatrix} 1.12 \\ 6.39 \end{pmatrix} \quad (2.0.14)$$

$$\Rightarrow \mathbf{D} = \begin{pmatrix} 6.12 \\ 6.39 \end{pmatrix} \quad (2.0.15)$$

- Now, the vertices of given Quadrilateral  $ABCD$  can be written as,

$$\mathbf{A} = \begin{pmatrix} -1.03 \\ 3.86 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \mathbf{C} = \begin{pmatrix} 5 \\ 0 \end{pmatrix}, \mathbf{D} = \begin{pmatrix} 6.12 \\ 6.39 \end{pmatrix} \quad (2.0.16)$$

- 3) On constructing the quadrilateral  $ABCD$  we get:

1




Fig. 2.1: Quadrilateral ABCD