

# Lagrange duality

## Mathematical Optimization

Georgiy Kormakov

CMC MSU

15 March 2024



# Dual function for norm

## Example

Let  $\|\cdot\|$  be the norm for  $\mathbb{R}^n$ , and  $\|\cdot\|_*$  be the dual norm. Prove that the conjugate function for  $f(x) = \|x\|$  is  $f^*(y) = 0$  with  $\text{dom } f^* = B_{\|\cdot\|_*}(0, 1)$ .

$$\|y\|_* > 1 :$$

$$\|y\|_* \leq 1 :$$

# Standart form

$$\begin{aligned} \min_x \quad & f_0(x) \\ \text{s.t.} \quad & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_j(x) = 0, \quad j = 1, \dots, n, \end{aligned} \tag{1}$$

# Standart form

$$\begin{aligned} \min_x & f_0(x) \\ \text{s.t. } & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_j(x) = 0, \quad j = 1, \dots, n, \end{aligned} \tag{1}$$

$x \in \mathbb{R}^d$ , but the whole domain is  $D = \cap_{i=1}^m \text{dom } f_i \cap \cap_{j=1}^n \text{dom } h_j$ .

# Standart form

$$\begin{aligned} \min_x & f_0(x) \\ \text{s.t. } & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_j(x) = 0, \quad j = 1, \dots, n, \end{aligned} \tag{1}$$

$x \in \mathbb{R}^d$ , but the whole domain is  $D = \cap_{i=1}^m \text{dom } f_i \cap \cap_{j=1}^n \text{dom } h_j$ .

## Feasible set

$$F = \{x \in D \mid f_i(x) \leq 0, i = \overline{1, m}, h_j(x) = 0, \forall j\}$$

# Standart form

$$\begin{aligned} \min_x \quad & f_0(x) \\ \text{s.t.} \quad & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_j(x) = 0, \quad j = 1, \dots, n, \end{aligned} \tag{1}$$

$x \in \mathbb{R}^d$ , but the whole domain is  $D = \cap_{i=1}^m \text{dom } f_i \cap \cap_{j=1}^n \text{dom } h_j$ .

## Feasible set

$$F = \{x \in D \mid f_i(x) \leq 0, i = \overline{1, m}, h_j(x) = 0, \forall j\}$$

## Lagrangian

For the optimisation problem 1, the *Lagrangian* is equal to

$$L(x, \lambda, \nu) = f_0(x) + \sum_{i=1}^m \lambda_i f_i(x) + \sum_{j=1}^n \nu_j h_j(x).$$

# Lagrangian Dual Function

## Definition

Let's define the *Lagrangian dual function* (or just the dual function)  $g : \mathbb{R}^m \times \mathbb{R}^n \rightarrow \mathbb{R}$  as follows:

$$g(\lambda, \nu) = \inf_{x \in D} \left( f_0(x) + \sum_{i=1}^m \lambda_i f_i(x) + \sum_{j=1}^n \nu_j h_j(x) \right) = \inf_{x \in D} L(x, \lambda, \nu) \quad (2)$$

# Lagrangian Dual Function

## Definition

Let's define the *Lagrangian dual function* (or just the dual function)  $g : \mathbb{R}^m \times \mathbb{R}^n \rightarrow \mathbb{R}$  as follows:

$$g(\lambda, \nu) = \inf_{x \in D} \left( f_0(x) + \sum_{i=1}^m \lambda_i f_i(x) + \sum_{j=1}^n \nu_j h_j(x) \right) = \inf_{x \in D} L(x, \lambda, \nu) \quad (2)$$

## Property 1

$g(\lambda, \nu)$  is always concave function of  $(\lambda, \nu)$



# Lagrangian Dual Function. Properties

## Property 2

$g(\lambda, \nu)$  is smaller than optimum  $p^*$  of 1 for  $\lambda \geq 0$  and any  $\nu$ :

$$\forall \lambda \geq 0, \nu \in \mathbb{R}^n \quad g(\lambda, \nu) \leq p^*$$

# Geometrical interpretation of Lagrangian

$$l_-(x) = \begin{cases} 0, & x \leq 0 \\ \infty, & \text{else} \end{cases} \quad l_0(x) = \begin{cases} 0, & x = 0 \\ \infty, & \text{else} \end{cases}$$

# Geometrical interpretation of Lagrangian

$$l_{-}(x) = \begin{cases} 0, & x \leq 0 \\ \infty, & \text{else} \end{cases} \quad l_0(x) = \begin{cases} 0, & x = 0 \\ \infty, & \text{else} \end{cases}$$

$$\min_x f_0(x) + \sum_{i=1}^m l_{-}(f_i(x)) + \sum_{j=1}^n l_0(h_j(x))$$

# Dual problem

## Definition

The dual problem for 1 has the following form

$$\begin{aligned} \max_{\lambda, \nu} \quad & g(\lambda, \nu) \\ \text{s.t.} \quad & \lambda \geq 0. \end{aligned} \tag{3}$$

# Linear equations with minimal norm

Consider the following optimization problem

$$\begin{aligned} \min_x \quad & x^T x \\ \text{s.t.} \quad & Ax = b, \end{aligned}$$

where  $A \in \mathbb{R}^{p \times d}$ ,  $b \in \mathbb{R}^p$ . Formulate the dual problem.

# Linear equations with minimal norm

Consider the following optimization problem

$$\begin{aligned} \min_x \quad & x^T x \\ \text{s.t.} \quad & Ax = b, \end{aligned}$$

where  $A \in \mathbb{R}^{p \times d}$ ,  $b \in \mathbb{R}^p$ . Formulate the dual problem.

$$x^* = -\frac{1}{2}A^T \nu \Rightarrow q(\nu) =$$

# Linear equations with minimal norm

Consider the following optimization problem

$$\begin{aligned} \min_x \quad & x^T x \\ \text{s.t.} \quad & Ax = b, \end{aligned}$$

where  $A \in \mathbb{R}^{p \times d}$ ,  $b \in \mathbb{R}^p$ . Formulate the dual problem.

$$x^* = -\frac{1}{2}A^T\nu \Rightarrow q(\nu) = -\frac{1}{4}\nu^T AA^T\nu - \nu^T b$$

# Linear Programming

$$\begin{aligned} \min_x \quad & c^T x \\ \text{s.t.} \quad & Ax = b, \\ & x \geq 0, \end{aligned}$$

where  $A \in \mathbb{R}^{p \times d}$ ,  $b \in \mathbb{R}^p$ .



# Linear Programming

$$\begin{aligned} \min_x \quad & c^T x \\ \text{s.t.} \quad & Ax = b, \\ & x \geq 0, \end{aligned}$$

where  $A \in \mathbb{R}^{p \times d}$ ,  $b \in \mathbb{R}^p$ .

$$g(\lambda, \nu) = \begin{cases} -\nu^T b, & c + A^T \nu - \lambda = 0 \\ -\infty, & \text{else} \end{cases}$$

# Linear Programming

$$\begin{aligned} \min_x \quad & c^T x \\ \text{s.t.} \quad & Ax = b, \\ & x \geq 0, \end{aligned}$$

where  $A \in \mathbb{R}^{p \times d}$ ,  $b \in \mathbb{R}^p$ .

$$g(\lambda, \nu) = \begin{cases} -\nu^T b, & c + A^T \nu - \lambda = 0 \\ -\infty, & \text{else} \end{cases}$$

$$\begin{aligned} \max_{\nu, \lambda} \quad & -\nu^T b \\ \text{s.t.} \quad & c + A^T \nu - \lambda = 0, \\ & \lambda \geq 0. \end{aligned}$$

# Splitting problem

$$\begin{aligned} \min_x & x^T W x \\ \text{s.t. } & x_j^2 = 1, \quad j = 1, \dots, d, \end{aligned}$$

where  $W \in \mathcal{S}^d$ .

# Splitting problem

$$\begin{aligned} \min_x & x^T W x \\ \text{s.t. } & x_j^2 = 1, \quad j = 1, \dots, d, \end{aligned}$$

where  $W \in \mathcal{S}^d$ .

$$g(\nu) = \begin{cases} -1^T \nu, & W + \text{diag}(\nu) \geq 0, \\ -\infty, & \text{else,} \end{cases}$$

# Splitting problem

$$\begin{aligned} \min_x & x^T W x \\ \text{s.t. } & x_j^2 = 1, \quad j = 1, \dots, d, \end{aligned}$$

where  $W \in \mathcal{S}^d$ .

$$g(\nu) = \begin{cases} -1^T \nu, & W + \text{diag}(\nu) \geq 0, \\ -\infty, & \text{else,} \end{cases}$$

$$\begin{aligned} \max_{\nu} & -1^T \nu \\ \text{s.t. } & W + \text{diag}(\nu) \geq 0. \end{aligned}$$

# Illustration

$$f^*(y) = \sup_{x \in \text{dom } f} (y^T x - f(x)).$$

## The simplest problem

$$\begin{aligned} \min_x \quad & f(x) \\ \text{s.t.} \quad & x = 0 \end{aligned}$$

# Illustration

$$f^*(y) = \sup_{x \in \text{dom } f} (y^T x - f(x)).$$

## The simplest problem

$$\begin{aligned} \min_x \quad & f(x) \\ \text{s.t.} \quad & x = 0 \end{aligned}$$

$$g(\nu) = \inf_x (f(x) + \nu^T x) = -\sup_x (-\nu^T x - f(x)) = -f^*(-\nu).$$

# Generalisation

$$\begin{aligned} \min_x \quad & f(x) \\ \text{s.t.} \quad & Ax \leq b, \\ & Cx = d. \end{aligned}$$



# Generalisation

$$\begin{aligned} \min_x \quad & f(x) \\ \text{s.t.} \quad & Ax \leq b, \\ & Cx = d. \end{aligned}$$

$$\begin{aligned} g(\lambda, \nu) &= \inf_x (f(x) + \lambda^T (Ax - b) + \nu^T (Cx - d)) \\ &= -\lambda^T b - \nu^T d + \inf_x (f(x) + (A^T \lambda + C^T \nu)^T x) \\ &= -\lambda^T b - \nu^T d - f^*(-A^T \lambda - C^T \nu). \end{aligned}$$

# System of equations. Again

$$\begin{aligned} \min_x \quad & \|x\| \\ \text{s.t.} \quad & Ax = b, \end{aligned}$$

where  $\|\cdot\|$  - any norm.

# System of equations. Again

$$\begin{aligned} \min_x \quad & \|x\| \\ \text{s.t.} \quad & Ax = b, \end{aligned}$$

where  $\|\cdot\|$  - any norm.

$$g(\nu) = -\nu^T b - f^*(-A^T \nu) = \begin{cases} -\nu^T b, & \|A^T \nu\|_* \leq 1 \\ -\infty, & \text{else.} \end{cases}$$

# System of equations. Again

$$\begin{aligned} \min_x \quad & \|x\| \\ \text{s.t.} \quad & Ax = b, \end{aligned}$$

where  $\|\cdot\|$  - any norm.

$$g(\nu) = -\nu^T b - f^*(-A^T \nu) = \begin{cases} -\nu^T b, & \|A^T \nu\|_* \leq 1 \\ -\infty, & \text{else.} \end{cases}$$

$$\begin{aligned} \max_{\nu} \quad & -\nu^T b \\ \text{s.t.} \quad & \|A^T \nu\|_* \leq 1. \end{aligned}$$

# Max Entropy

$$\begin{aligned} \min_x \quad & \sum_{i=1}^d x_i \log x_i \\ \text{s.t.} \quad & Ax \leq b, \\ & 1^T x = 1, \end{aligned}$$

where  $\text{dom } f = \mathbb{R}_{++}^d$ .

# Max Entropy

$$\begin{aligned} \min_x \quad & \sum_{i=1}^d x_i \log x_i \\ \text{s.t.} \quad & Ax \leq b, \\ & 1^T x = 1, \end{aligned}$$

where  $\text{dom } f = \mathbb{R}_{++}^d$ .

$$f^*(y) = \sum_{i=1}^d e^{y_i - 1}.$$

# Max Entropy

$$\begin{aligned} \min_x \quad & \sum_{i=1}^d x_i \log x_i \\ \text{s.t.} \quad & Ax \leq b, \\ & 1^T x = 1, \end{aligned}$$

where  $\text{dom } f = \mathbb{R}_{++}^d$ .

$$f^*(y) = \sum_{i=1}^d e^{y_i - 1}.$$

$$g(\lambda, \nu) = -\lambda^T b - \nu - \sum_{i=1}^d e^{-a_i^T \lambda - \nu - 1},$$

# Max Entropy

$$\begin{aligned} \min_x \quad & \sum_{i=1}^d x_i \log x_i \\ \text{s.t.} \quad & Ax \leq b, \\ & 1^T x = 1, \end{aligned}$$

where  $\text{dom } f = \mathbb{R}_{++}^d$ .

$$f^*(y) = \sum_{i=1}^d e^{y_i - 1}.$$

$$g(\lambda, \nu) = -\lambda^T b - \nu - \sum_{i=1}^d e^{-a_i^T \lambda - \nu - 1},$$

$$\max_{\lambda, \nu} \left[ -\lambda^T b - \nu - \sum_{i=1}^d e^{-a_i^T \lambda - \nu - 1} \right] \text{ s.t. } \lambda \geq 0.$$



# Change of variables

$$\min_x \sum_{i=1}^N \|A_i x + b_i\|_2 + \frac{1}{2} \|x - x_0\|_2^2.$$

# Change of variables

$$\min_x \sum_{i=1}^N \|A_i x + b_i\|_2 + \frac{1}{2} \|x - x_0\|_2^2.$$

$$\min_x \sum_{i=1}^N \|y_i\|_2 + \frac{1}{2} \|x - x_0\|_2^2$$

$$\text{s.t. } A_j x + b_j = y_j \quad \forall j$$

# Change of variables

$$\min_x \sum_{i=1}^N \|A_i x + b_i\|_2 + \frac{1}{2} \|x - x_0\|_2^2.$$

$$\min_x \sum_{i=1}^N \|y_i\|_2 + \frac{1}{2} \|x - x_0\|_2^2$$

$$\text{s.t. } A_j x + b_j = y_j \quad \forall j$$

$$g(\nu_1, \dots, \nu_N) = \begin{cases} \sum_{i=1}^N (A_i x_0 + b_i)^T \nu_i - \frac{1}{2} \left\| \sum_{j=1}^N A_j^T \nu_j \right\|_2^2, & \|\nu_j\|_2 \leq 1 \quad \forall i \\ -\infty, & \text{else} \end{cases}$$

# Piece-wise linear

$$\min_x \max_{i=1,\dots,m} (a_i^T x + b_i)$$

# Piece-wise linear

$$\min_x \max_{i=1,\dots,m} (a_i^T x + b_i)$$

$$\min_x \max_{i=1,\dots,m} y_i$$

$$\text{s.t. } a_j^T x + b_j = y_j \quad \forall j.$$

# Piece-wise linear

$$\min_x \max_{i=1,\dots,m} (a_i^T x + b_i)$$

$$\min_x \max_{i=1,\dots,m} y_i$$

$$\text{s.t. } a_j^T x + b_j = y_j \quad \forall j.$$

$$\inf_y (\max_i y_i - \nu^T y) = \begin{cases} 0, & \nu \geq 0, \quad 1^T \nu = 1 \\ -\infty, & \text{else} \end{cases}$$

# Piece-wise linear

$$\min_x \max_{i=1,\dots,m} (a_i^T x + b_i)$$

$$\min_x \max_{i=1,\dots,m} y_i$$

$$\text{s.t. } a_j^T x + b_j = y_j \quad \forall j.$$

$$\inf_y (\max_i y_i - \nu^T y) = \begin{cases} 0, & \nu \geq 0, \quad 1^T \nu = 1 \\ -\infty, & \text{else} \end{cases}$$

$$g(\nu) = \begin{cases} b^T \nu, & \sum_{i=1}^m \nu_i a_i = 0, \quad \nu \geq 0, \quad 1^T \nu = 1 \\ -\infty, & \text{else} \end{cases}$$