

Linear Equations




Figure 1




Figure 2

A linear equation has the following general form:

$$y = ax + b$$

In this equation, a and b are constants. This equation is called linear because the graph of y versus x is a straight line, as shown in **Figure 1**. The constant b , called the *intercept*, represents the value of y where the straight line intersects the y -axis. The constant a is equal to the *slope* of the straight line and is also equal to the tangent of the angle that the line makes with the x -axis (θ). If any two points on the straight line are specified by the coordinates (x_1, y_1) and (x_2, y_2) , as in **Figure 1**, then the slope of the straight line can be expressed as follows:

$$\text{slope} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{\Delta y}{\Delta x}$$

For example, if the two points shown in **Figure 1** are $(2, 4)$ and $(6, 9)$, then the slope of the line is as follows:

$$\text{slope} = \frac{(9 - 4)}{(6 - 2)} = \frac{5}{4}$$

Note that a and b can have either positive or negative values. If $a > 0$, the straight line has a *positive* slope, as in **Figure 1**. If $a < 0$, the straight line has a *negative* slope. Furthermore, if $b > 0$, the y intercept is positive (above the x -axis), while if $b < 0$, the y intercept is negative (below the x -axis). **Figure 2** gives an example of each of these four possible cases, which are summarized in **Table 4**.

Table 4 Linear Equations

Constants	Slope	y intercept
$a > 0, b > 0$	positive slope	positive y intercept
$a > 0, b < 0$	positive slope	negative y intercept
$a < 0, b > 0$	negative slope	positive y intercept
$a < 0, b < 0$	negative slope	negative y intercept

Solving Simultaneous Linear Equations

Consider the following equation:

$$3x + 5y = 15$$

This equation has two unknowns, x and y . Such an equation does not have a unique solution. That is, $(x = 0, y = 3)$, $(x = 5, y = 0)$, and $(x = 2, y = \frac{9}{5})$ are all solutions to this equation.

If a problem has two unknowns, a unique solution is possible only if there are two independent equations. In general, if a problem has n unknowns, its solution requires n independent equations. There are three basic methods that can be used to solve simultaneous equations. Each of these methods is discussed below, and an example is given for each.