Metallic Bonding

C hemical bonding is different in metals than it is in ionic, molecular, or covalent-network compounds. This difference is reflected in the unique properties of metals. They are excellent electrical conductors in the solid state—much better conductors than even molten ionic compounds. This property is due to the highly mobile valence electrons of the atoms that make up a metal. Such mobility is not possible in molecular compounds, in which valence electrons are localized in electron-pair bonds between neutral atoms. Nor is it possible in solid ionic compounds, in which electrons are bound to individual ions that are held in place in crystal structures.

The Metallic-Bond Model

The highest energy levels of most metal atoms are occupied by very few electrons. In s-block metals, for example, one or two valence electrons occupy the outermost orbital, and all three outermost p orbitals, which can hold a total of six electrons, are vacant. In addition to completely vacant outer p orbitals, d-block metals also possess many vacant d orbitals in the energy level just below their highest energy level.

Within a metal, the vacant orbitals in the atoms' outer energy levels overlap. This overlapping of orbitals allows the outer electrons of the atoms to roam freely throughout the entire metal. The electrons are *delocalized*, which means that they do not belong to any one atom but move freely about the metal's network of empty atomic orbitals. These mobile electrons form a *sea of electrons* around the metal atoms, which are packed together in a crystal lattice (see **Figure 18**). The chemical bonding that results from the attraction between metal atoms and the surrounding sea of electrons is called **metallic bonding**.

Metallic Properties

The freedom of electrons to move in a network of metal atoms accounts for the high electrical and thermal conductivity characteristic of all metals. In addition, metals are both strong absorbers and reflectors of light. Because they contain many orbitals separated by extremely small energy differences, metals can absorb a wide range of light frequencies. This absorption of light results in the excitation of the metal atoms' electrons to higher energy levels. However, in metals the electrons immediately fall back down to lower levels, emitting energy in the form of light at a frequency similar to the absorbed frequency. This re-radiated (or reflected) light is responsible for the metallic appearance or luster of metal surfaces.

SECTION 4

OBJECTIVES

- Describe the electron-sea model of metallic bonding, and explain why metals are good electrical conductors.
- Explain why metal surfaces are shiny.
- Explain why metals are malleable and ductile but ioniccrystalline compounds are not.






FIGURE 18 The model shows a portion of the crystal structure of solid sodium. The atoms are arranged so that each sodium atom is surrounded by eight other sodium atoms. The atoms are relatively fixed in position, while the electrons are free to move throughout the crystal, forming an electron sea.