CHAPTER HIGHLIGHTS

The Atom: From Philosophical Idea to Scientific Theory

Vocabulary

law of conservation of mass law of definite proportions law of multiple proportions

- The idea of atoms has been around since the time of the ancient Greeks. In the nineteenth century, John Dalton proposed a scientific theory of atoms that can still be used to explain properties of most chemicals today.
- Matter and its mass cannot be created or destroyed in chemical reactions.
- The mass ratios of the elements that make up a given compound are always the same, regardless of how much of the compound there is or how it was formed.
- If two or more different compounds are composed of the same two elements, then the ratio of the masses of the second element combined with a certain mass of the first element can be expressed as a ratio of small whole numbers.

The Structure of the Atom

Vocabulary

atom nuclear forces

- Cathode-ray tubes supplied evidence of the existence of electrons, which are negatively charged subatomic particles that have relatively little mass.
- Rutherford found evidence for the existence of the atomic nucleus by bombarding metal foil with a beam of positively charged particles.
- Atomic nuclei are composed of protons, which have an electric charge of +1, and (in all but one case) neutrons, which have no electric charge.
- Atomic nuclei have radii of about 0.001 pm (pm = picometers; $1 \text{ pm} \times 10^{-12} \text{ m}$), and atoms have radii of about 40–270 pm.

Counting Atoms

Vocabulary

atomic number isotope mass number nuclide atomic mass unit average atomic mass mole Avogadro's number molar mass

- The atomic number of an element is equal to the number of protons of an atom of that element.
- The mass number is equal to the total number of protons and neutrons that make up the nucleus of an atom of that element.
- The relative atomic mass unit (amu) is based on the carbon-12 atom and is a convenient unit for measuring the mass of atoms. It equals $1.660\ 540 \times 10^{-24}\ g$.
- The average atomic mass of an element is found by calculating the weighted average of the atomic masses of the naturally occurring isotopes of the element.
- Avogadro's number is equal to approximately 6.022 1415 × 10²³.
 A sample that contains a number of particles equal to Avogadro's number contains a mole of those particles.