

M083 Stroke: Ischemic - Integrated Admission Rule Hierarchy

Admission only. Domain paths are shown with linked shared definitions embedded under each linked rule.

Definition

Composite Definitions

- [altered_mental_status_condition_present](#)
- [cardiac_arrhythmia_of_immediate_concern_condition_present](#)
- [dangerous_arrhythmia_condition_present](#)
- [hemodynamic_instability_condition_present](#)
- [hypotension_condition_present](#)
- [hypoxemia_condition_present](#)
- [orthostatic_hypotension_condition_present](#)
- [prolonged_cardiac_telemetry_monitoring_required](#)
- [respiratory_abnormality_condition_present](#)
- [tachycardia_condition_present](#)

Atomic Numeric Definitions

- [arterial_or_venous_ph](#)
- [arterial_po2_mmhg](#)
- [diastolic_blood_pressure_mmhg](#)
- [lactate_mmol_l](#)
- [mean_arterial_pressure_mmhg](#)
- [nihss_score](#)
- [orthostatic_diastolic_blood_pressure_drop_mmhg](#)
- [orthostatic_systolic_blood_pressure_drop_mmhg](#)
- [shock_index](#)
- [spo2_percent](#)

Atomic Boolean / Leaf Definitions

- [acute_ischemic_stroke_condition_present](#)
- [acute_myocarditis_condition_present](#)
- [aphasia_condition_present](#)
- [automatic_icd_firing_malfunction_or_setting_adjustment_needed_condition_present](#)
- [baseline_sbp_decrease_with_symptoms_condition_present](#)
- [brain_imaging_finding_requiring_inpatient_care_condition_present](#)
- [cerebral_venous_thrombosis_condition_present](#)
- [clinically_active_atrial_fibrillation_condition_present](#)
- [clinically_significant_cardiac_or_vascular_disorder_condition_present](#)
- [coma_condition_present](#)
- [confusional_state_condition_present](#)
- [continuous_long_term_ecg_monitoring_needed_condition_present](#)
- [dangerous_arrhythmia_associated_significant_symptoms_condition_present](#)
- [dysphagia_evaluation_required](#)
- [end_organ_dysfunction_due_to_hypotension_condition_present](#)
- [gait_impairment_condition_present](#)
- [hemorrhagic_transformation_condition_present](#)
- [hypotension_in_dangerous_arrhythmia_context_condition_present](#)
- [increased_supplemental_oxygen_from_baseline_condition_present](#)
- [lethargy_condition_present](#)
- [myocardial_ischemia_condition_present](#)
- [neurologic_worsening_monitoring_required](#)

- new_onset_lbbb_with_suspected_myocardial_ischemia_condition_present
- nonsustained_ventricular_tachycardia_condition_present
- obtundation_condition_present
- persistence_despite_appropriate_treatment_context_present
- prolonged_telemetry_beyond_observation_required
- respiratory_distress_in_dangerous_arrhythmia_context_condition_present
- resuscitated_ventricular_fibrillation_or_cardiac_arrest_condition_present
- significant_limb_weakness_condition_present
- stupor_condition_present
- supplemental_oxygen_required_to_maintain_targets_condition_present
- suspected_cardiac_ischemia_as_cause_or_consequence_of_vt_condition_present
- suspected_vasculitis_condition_present
- sustained_ventricular_tachycardia_condition_present
- third_degree_av_block_condition_present
- thrombolysis_or_thrombectomy_performed_or_planned_condition_present
- type_ii_second_degree_av_block_condition_present
- vasopressor_or_inotrope_required
- ventricular_escape_rhythm_condition_present

Needs Review / Pediatric Placeholders

- atrial_fibrillation_detection_only_context_absent
- clinical_stability_unclear_condition_present
- heart_rate_bpm
- impaired_airway_protection_condition_present
- pediatric_hypoperfusion_due_to_hypotension_condition_present
- pediatric_severe_hypertension_threshold_condition_present
- pediatric_shock_index_threshold_condition_present
- pediatric_systolic_blood_pressure_threshold_condition_present
- pediatric_vasopressor_or_inotrope_required
- systolic_blood_pressure_mmhg

M083

Admission is indicated for 1 or more of the following

Path 1: Acute ischemic stroke with neurologic findings that warrant inpatient care, as indicated by 1 or more of the following

- [composite / AND] Acute ischemic stroke with neurologic findings that warrant inpatient care, as indicated by 1 or more of the following
 - [context / IS_TRUE / acute_ischemic_stroke_condition_present] Acute ischemic stroke

Linked shared atomic definition: acute_ischemic_stroke_condition_present -> atom.M083.admission.acute_ischemic_stroke_condition_present

IS_TRUE

Acute ischemic stroke

review_status=ok source_strategy=domain_source_atomic_leaf

- [composite / OR] Acute ischemic stroke [A] with neurologic findings that warrant inpatient care, as indicated by 1 or more of the following:
 - [atomic / > / nihss_score] National Institutes of Health Stroke Scale (NIHSS) score greater than 2 [B] (2) (9)

Linked shared atomic definition: nihss_score -> atom.M083.admission.nihss_score

NIHSS > 2 score

National Institutes of Health Stroke Scale (NIHSS) score greater than 2 [B] (2) (9)

review_status=ok source_strategy=domain_source_numeric_leaf

- [atomic / IS_TRUE / hemorrhagic_transformation_condition_present] Evidence of hemorrhagic transformation (4) (9)

Linked shared atomic definition: hemorrhagic_transformation_condition_present -> atom.M083.admission.hemorrhagic_transformation_condition_present

IS_TRUE

Evidence of hemorrhagic transformation (4) (9)

review_status=ok source_strategy=domain_source_atomic_leaf

- [atomic / IS_TRUE / altered_mental_status_condition_present] Altered mental status (10)

Linked shared definition: altered_mental_status_condition_present -> comp.altered_mental_status.root

Source: Definition - Altered mental status

- OR altered_mental_status_condition_present

Altered mental status (ie, different from baseline), as indicated by 1 or more of the following

- confusional_state_condition_present

Confusional state (eg, disorientation, difficulty following commands, deficit in attention)

- lethargy_condition_present

Lethargy (eg, awake or arousable, but with drowsiness; reduced awareness of self and environment)

- obtundation_condition_present

Obtundation (ie, arousable only with strong stimuli, lessened interest in environment, slowed responses to stimulation)

- stupor_condition_present

Stupor (ie, may be arousable but patient does not return to normal baseline level of awareness)

- coma_condition_present

Coma (ie, not arousable)

- [atomic / IS_TRUE / dysphagia_evaluation_required] Dysphagia that warrants evaluation (eg, aspiration risk suspected) (2) (9)

Linked shared atomic definition: dysphagia_evaluation_required -> atom.M083.admission.dysphagia_evaluation_required

IS_TRUE

Dysphagia that warrants evaluation (eg, aspiration risk suspected) (2) (9)

review_status=ok source_strategy=domain_source_atomic_leaf

- [atomic / IS_TRUE / significant_limb_weakness_condition_present] Significant arm or leg weakness (eg, limiting movement against gravity) (4)

Linked shared atomic definition: significant_limb_weakness_condition_present -> atom.M083.admission.significant_limb_weakness_condition_present

IS_TRUE

Significant arm or leg weakness (eg, limiting movement against gravity) (4)

review_status=ok source_strategy=domain_source_atomic_leaf

- [atomic / IS_TRUE / aphasia_condition_present] Aphasia (4)

Linked shared atomic definition: aphasia_condition_present -> atom.M083.admission.aphasia_condition_present

IS_TRUE

Aphasia (4)

review_status=ok source_strategy=domain_source_atomic_leaf

- [atomic / IS_TRUE / gait_impairment_condition_present] Gait impairment (4)

Linked shared atomic definition: gait_impairment_condition_present -> atom.M083.admission.gait_impairment_condition_present
IS_TRUE
Gait impairment (4)
review_status=ok source_strategy=domain_source_atomic_leaf

- [atomic / IS_TRUE / brain_imaging_finding_requiring_inpatient_care_condition_present] Finding on brain imaging that requires inpatient care (eg, edema, threatened herniation) (2) (9)

Linked shared atomic definition: brain_imaging_finding_requiring_inpatient_care_condition_present -> atom.M083.admission.brain_imaging_finding_requiring_inpatient_care_condition_present
IS_TRUE
Finding on brain imaging that requires inpatient care (eg, edema, threatened herniation) (2) (9)
review_status=ok source_strategy=domain_source_atomic_leaf

- [atomic / IS_TRUE / neurologic_worsening_monitoring_required] Clinical monitoring for neurologic worsening needed (eg, large infarct, posterior circulation infarct) [C] (2) (9) (11)

Linked shared atomic definition: neurologic_worsening_monitoring_required -> atom.M083.admission.neurologic_worsening_monitoring_required
IS_TRUE
Clinical monitoring for neurologic worsening needed (eg, large infarct, posterior circulation infarct) [C] (2) (9) (11)
review_status=ok source_strategy=domain_source_atomic_leaf

- [atomic / IS_TRUE / clinical_stability_unclear_condition_present] Clinical stability unclear (eg, recurrent or worsening findings)

Linked shared atomic definition: clinical_stability_unclear_condition_present -> atom.M083.admission.clinical_stability_unclear_condition_present
IS_TRUE
Clinical stability unclear (eg, recurrent or worsening findings)
review_status=needs_review source_strategy>manual_review_placeholder

Path 2: Acute ischemic stroke with clinical need for inpatient care or monitoring, as indicated by 1 or more of the following

- [composite / AND] Acute ischemic stroke with clinical need for inpatient care or monitoring, as indicated by 1 or more of the following
 - [context / IS_TRUE / acute_ischemic_stroke_condition_present] Acute ischemic stroke

Linked shared atomic definition: acute_ischemic_stroke_condition_present -> atom.M083.admission.acute_ischemic_stroke_condition_present

IS_TRUE

Acute ischemic stroke

review_status=ok source_strategy=domain_source_atomic_leaf

- [composite / OR] Acute ischemic stroke [A] with clinical need for inpatient care or monitoring, as indicated by 1 or more of the following:
 - [atomic / IS_TRUE / hemodynamic_instability_condition_present] Hemodynamic instability (2) (9) (12)

Linked shared definition: hemodynamic_instability_condition_present -> comp.hemodynamic_instability.root

Source: Definition - Hemodynamic instability

- **OR hemodynamic_instability_condition_present**

Hemodynamic instability, as indicated by 1 or more of the following

- **OR vital_sign_abnormality_not_corrected_condition_present**

Vital sign abnormality not corrected by appropriate treatment, as indicated by 1 or more of the following:

- **AND persistent_tachycardia_condition_present**

Tachycardia that persists despite appropriate treatment (eg, treatment of underlying cause, despite observation care)

- **OR tachycardia_condition_present**

Tachycardia,[A][B] as indicated by 1 or more of the following:

- **heart_rate > 100 bpm**

Heart rate greater than 100 beats per minute in adult{A}[B](1)

- **heart_rate > 85 bpm (needs_review)**

Heart rate greater than 85 beats per minute in child 13 to 17 years of age{A}[C](2)

- **heart_rate > 95 bpm (needs_review)**

Heart rate greater than 95 beats per minute in child 6 to 12 years of age{A}[C](2)

- **heart_rate > 110 bpm (needs_review)**

Heart rate greater than 110 beats per minute in child 1 to 5 years of age{A}[C](2)

- **heart_rate > 120 bpm (needs_review)**

Heart rate greater than 120 beats per minute in infant 3 to 11 months of age{A}[C](2)

- **heart_rate > 150 bpm (needs_review)**

Heart rate greater than 150 beats per minute in infant 1 or 2 months of age{A}[C](2)

- **persistence_despite_appropriate_treatment_context_present**

persists despite appropriate treatment

- **AND persistent_hypotension_condition_present**

Hypotension that persists despite appropriate treatment (eg, treatment of underlying cause, despite observation care)

- **OR hypotension_condition_present**

Hypotension, as indicated by 1 or more of the following:

- **OR hypotension_in_adult_patient_condition_present**

Hypotension in adult patient, as indicated by 1 or more of the following:

- **SBP < 90 mmHg**

Systolic blood pressure less than 90 mm Hg{A}[B](1)

- **MAP < 70 mmHg**

Mean arterial pressure{C} less than 70 mm Hg{A}[B] MAP Calculator(1)(2)

- **baseline_sbp_decrease_with_symptoms_condition_present**

Decrease in baseline systolic blood pressure of 30 mm Hg or more, with significant signs or symptoms due to lower blood pressure (eg, near syncope, syncope, chest pain){A}[B](1)

- **OR hypotension_in_pediatric_patient_condition_present**

Hypotension in pediatric patient, as indicated by 1 or more of the following:

- **SBP < 110 mmHg (needs_review)**

Systolic blood pressure less than 110 mm Hg in child 13 to 17 years of age{A}[D](3)

- **SBP < 100 mmHg (needs_review)**

Systolic blood pressure less than 100 mm Hg in child 6 to 12 years of age{A}[D](3)

- **SBP < 95 mmHg (needs_review)**

Systolic blood pressure less than 95 mm Hg in child 3 to 5 years of age{A}[D](3)

- **SBP < 90 mmHg (needs_review)**

Systolic blood pressure less than 90 mm Hg in child 1 or 2 years of age{A}[D](3)

- **SBP < 80 mmHg (needs_review)**

- Systolic blood pressure less than 80 mm Hg in infant 6 to 11 months of age[A][D](3)
 - **SBP < 70 mmHg (needs_review)**
Systolic blood pressure less than 70 mm Hg in infant 3 to 5 months of age[A][D](3)
 - **SBP < 65 mmHg (needs_review)**
Systolic blood pressure less than 65 mm Hg in infant 1 or 2 months of age[A][D](3)
- **persistence_despite_appropriate_treatment_context_present**
persists despite appropriate treatment
- **AND persistent_orthostatic_hypotension_condition_present**
Orthostatic hypotension that persists despite appropriate treatment (eg, treatment of underlying cause, despite observation care)
 - **OR orthostatic_hypotension_condition_present**
Orthostatic hypotension,[A][B] as indicated by 1 or more of the following(1)(2)(3):
 - **orthostatic_SBP_drop >= 20 mmHg**
Fall in SBP of 20 mm Hg or more 1 to 3 minutes after patient sits or stands from recumbent position
 - **orthostatic_DBP_drop >= 10 mmHg**
Fall in DBP of 10 mm Hg or more 1 to 3 minutes after patient sits or stands from recumbent position
 - **persistence_despite_appropriate_treatment_context_present**
persists despite appropriate treatment
- **OR severe_adult_hypotension_condition_present**
Severe hypotension in adult patient, as indicated by 1 or more of the following:
 - **SBP < 80 mmHg**
Systolic blood pressure less than 80 mm Hg
 - **shock_index > 1.0 ratio**
Shock index greater than 1.0 (shock index is the heart rate divided by systolic blood pressure)
 - **MAP < 65 mmHg**
Mean arterial pressure[B] less than 65 mm Hg MAP Calculator[A](1)(6)
 - **vasopressor_or_inotrope_required**
IV inotropic or vasopressor medication required to maintain adequate blood pressure or perfusion
 - **AND hypoperfusion_due_to_hypotension_condition_present**
Hypoperfusion due to hypotension, as indicated by ALL of the following:
 - **OR hypotension_condition_present**
Hypotension, as indicated by 1 or more of the following:
 - **OR hypotension_in_adult_patient_condition_present**
Hypotension in adult patient, as indicated by 1 or more of the following:
 - **SBP < 90 mmHg**
Systolic blood pressure less than 90 mm Hg[A][B](1)
 - **MAP < 70 mmHg**
Mean arterial pressure[C] less than 70 mm Hg[A][B] MAP Calculator(1)(2)
 - **baseline_sbp_decrease_with_symptoms_condition_present**
Decrease in baseline systolic blood pressure of 30 mm Hg or more, with significant signs or symptoms due to lower blood pressure (eg, near syncope, syncope, chest pain)[A][B](1)
 - **OR hypotension_in_pediatric_patient_condition_present**
Hypotension in pediatric patient, as indicated by 1 or more of the following:
 - **SBP < 110 mmHg (needs_review)**
Systolic blood pressure less than 110 mm Hg in child 13 to 17 years of age[A][D](3)
 - **SBP < 100 mmHg (needs_review)**
Systolic blood pressure less than 100 mm Hg in child 6 to 12 years of age[A][D](3)
 - **SBP < 95 mmHg (needs_review)**
Systolic blood pressure less than 95 mm Hg in child 3 to 5 years of age[A][D](3)
 - **SBP < 90 mmHg (needs_review)**
Systolic blood pressure less than 90 mm Hg in child 1 or 2 years of age[A][D](3)
 - **SBP < 80 mmHg (needs_review)**
Systolic blood pressure less than 80 mm Hg in infant 6 to 11 months of age[A][D](3)
 - **SBP < 70 mmHg (needs_review)**
Systolic blood pressure less than 70 mm Hg in infant 3 to 5 months of age[A][D](3)
 - **SBP < 65 mmHg (needs_review)**
Systolic blood pressure less than 65 mm Hg in infant 1 or 2 months of age[A][D](3)
 - **OR hypoperfusion_evidence_condition_present**
Evidence of hypoperfusion, as indicated by 1 or more of the following:
 - **lactate >= 2.0 mmol/L**
Lactate of 2.0 mmol/L (18 mg/dL) or more secondary to hypotension (ie, hypoperfusion)
 - **pH < 7.35 pH**
Metabolic acidosis (arterial or venous pH less than 7.35) not otherwise explained

Any heart rhythm and 1 or more of the following(5)(8)(10)(11)(12)(13):

- **suspected_cardiac_ischemia_as_cause_or_consequence_of_vt_condition_present**
Suspected cardiac ischemia as cause or consequence of ventricular tachycardia
- **continuous_long_term_ecg_monitoring_needed_condition_present**
Continuous long-term ECG monitoring needed (eg, initiation of drug requiring monitoring beyond observation care)
- **automatic_icd_firing_malfunction_or_setting_adjustment_needed_condition_present**
Patient has automatic implantable cardioverter-defibrillator that is repeatedly firing, malfunctioning, or in need of immediate adjustment of settings beyond scope of observation care.

- **[atomic / IS_TRUE / clinically_significant_cardiac_or_vascular_disorder_condition_present]** Clinically significant cardiac or vascular disorder identified (eg, on echocardiogram, imaging) that requires inpatient care (eg, severe valvular disease, atrial myxoma, cervical artery dissection) (15)

Linked shared atomic definition: **clinically_significant_cardiac_or_vascular_disorder_condition_present** -> atom.M083.admission.clinically_significant_cardiac_or_vascular_disorder_condition_present

IS_TRUE

Clinically significant cardiac or vascular disorder identified (eg, on echocardiogram, imaging) that requires inpatient care (eg, severe valvular disease, atrial myxoma, cervical artery dissection) (15)

review_status=ok source_strategy=domain_source_atomic_leaf

- **[atomic / IS_TRUE / cerebral_venous_thrombosis_condition_present]** Ischemic stroke due to cerebral venous thrombosis [D] (1) (16)

Linked shared atomic definition: **cerebral_venous_thrombosis_condition_present** -> atom.M083.admission.cerebral_venous_thrombosis_condition_present

IS_TRUE

Ischemic stroke due to cerebral venous thrombosis [D] (1) (16)

review_status=ok source_strategy=domain_source_atomic_leaf

- **[atomic / IS_TRUE / respiratory_abnormality_condition_present]** Respiratory abnormalities (eg, Hypoxemia , impaired airway protection) (2)

Linked shared definition: **respiratory_abnormality_condition_present** -> comp.respiratory_abnormality.root

Source: Definition - Hypoxemia

- **OR respiratory_abnormality_condition_present**
Respiratory abnormalities (eg, Hypoxemia, impaired airway protection)
 - **OR hypoxemia_condition_present**
Hypoxemia, as indicated by 1 or more of the following(1):
 - **OR hypoxemia_room_air_saturation_or_pao2_criteria_condition_present**
Patient without baseline need for supplemental oxygen with oxygen saturation (SpO2) of less than 90% or arterial blood gas partial pressure of oxygen (PaO2) of less than 60 mm Hg (8.0 kPa) on room air[A][B]
 - **SpO2 < 90 %**
Patient without baseline need for supplemental oxygen with oxygen saturation (SpO2) of less than 90% or arterial blood gas partial pressure of oxygen (PaO2) of less than 60 mm Hg (8.0 kPa) on room air[A][B]
 - **PaO2 < 60 mmHg**
Patient without baseline need for supplemental oxygen with oxygen saturation (SpO2) of less than 90% or arterial blood gas partial pressure of oxygen (PaO2) of less than 60 mm Hg (8.0 kPa) on room air[A][B]
 - **supplemental_oxygen_required_to_maintain_targets_condition_present**
Patient without baseline need for supplemental oxygen who now requires supplemental oxygen to keep SpO2 greater than 89% or PaO2 greater than 59 mm Hg (7.9 kPa)[A]
 - **increased_supplemental_oxygen_from_baseline_condition_present**
Patient with baseline need for supplemental oxygen who now requires increased supplemental oxygen to maintain oxygenation at baseline or acceptable level
 - **impaired_airway_protection_condition_present (needs_review)**
impaired airway protection

- **[atomic / IS_TRUE / prolonged_cardiac_telemetry_monitoring_required]** Prolonged cardiac telemetry monitoring needed (eg, beyond observation care time frame) due to concern for Dangerous arrhythmia (ie, not in effort to detect atrial fibrillation) [G]

Linked shared definition: **prolonged_cardiac_telemetry_monitoring_required** -> comp.prolonged_cardiac_telemetry_monitoring.root

Source: Definition - Cardiac arrhythmias of immediate concern

- **AND prolonged_cardiac_telemetry_monitoring_required**
Prolonged cardiac telemetry monitoring needed (eg, beyond observation care time frame) due to concern for Dangerous arrhythmia (ie, not in effort to detect atrial fibrillation) [G]
 - **prolonged_telemetry_beyond_observation_required**
Prolonged cardiac telemetry monitoring needed (eg, beyond observation care time frame)
 - **OR dangerous_arrhythmia_condition_present**
Dangerous arrhythmia, as indicated by 1 or more of the following(1)(2)(3)(4):
 - **OR dangerous_inherently_unstable_rhythm_mechanisms_condition_present**
Heart rhythms that are inherently dangerous or unstable, as indicated by 1 or more of the following(5)(6):
 - **resuscitated_ventricular_fibrillation_or_cardiac_arrest_condition_present**
Resuscitated ventricular fibrillation or cardiac arrest
 - **ventricular_escape_rhythm_condition_present**
Ventricular escape rhythm
 - **sustained_ventricular_tachycardia_condition_present**

Sustained ventricular tachycardia (30 seconds or more of ventricular rhythm at greater than 100 beats per minute)

- **AND nonsustained_vt_with_comorbidity_condition_present** (needs_review)

Nonsustained ventricular tachycardia and 1 or more of the following:

- **nonsustained_ventricular_tachycardia_condition_present**

Nonsustained ventricular tachycardia and 1 or more of the following:

- **OR nsvt_with_acute_myocarditis_or_myocardial_ischemia_condition_present**

Nonsustained ventricular tachycardia and 1 or more of the following:

- **acute_myocarditis_condition_present**

Acute myocarditis

- **myocardial_ischemia_condition_present**

Myocardial ischemia

- **OR unstable_cardiac_conduction_defects_condition_present**

Unstable cardiac conduction defects, as indicated by 1 or more of the following(7):

- **type_ii_second_degree_av_block_condition_present**

Type II second-degree atrioventricular block

- **third_degree_av_block_condition_present**

Third-degree atrioventricular block

- **new_onset_lbbb_with_suspected_myocardial_ischemia_condition_present**

New-onset left bundle branch block with suspected myocardial ischemia

- **OR dangerous_rhythm_hemodynamic_or_symptom_concern_condition_present**

Heart rhythms of concern due to 1 or more of the following(8):

- **hypotension_in_dangerous_arrhythmia_context_condition_present**

Hypotension

- **respiratory_distress_in_dangerous_arrhythmia_context_condition_present**

Respiratory distress

- **dangerous_arrhythmia_associated_significant_symptoms_condition_present**

Association with other significant symptoms (eg, syncope)(7)(9)(10)(11)

- **atrial_fibrillation_detection_only_context_absent** (needs_review)

due to concern for Dangerous arrhythmia (ie, not in effort to detect atrial fibrillation)

- **[atomic / IS_TRUE / suspected_vasculitis_condition_present]** Suspected vasculitis (21) (22)

Linked shared atomic definition: suspected_vasculitis_condition_present -> atom.M083.admission.suspected_vasculitis_condition_present

IS_TRUE

Suspected vasculitis (21) (22)

review_status=ok source_strategy=domain_source_atomic_leaf

- **[composite / OR]** Severe hypertension (SBP greater than 180 mm Hg or DBP greater than 120 mm Hg, or greater than the 95th percentile for age, sex, and height plus 30 mm Hg in pediatric patients [E] [F]) (2) (4) (17) (18)

- **[atomic / > / systolic_blood_pressure_mmhg]** Systolic blood pressure greater than 180 mm Hg

Linked shared atomic definition: systolic_blood_pressure_mmhg -> atom.M083.admission.systolic_blood_pressure_mmhg

SBP > 180 mmHg

Systolic blood pressure greater than 180 mm Hg

review_status=ok source_strategy=domain_source_numeric_leaf

- **[atomic / > / diastolic_blood_pressure_mmhg]** Diastolic blood pressure greater than 120 mm Hg

Linked shared atomic definition: diastolic_blood_pressure_mmhg -> atom.M083.admission.diastolic_blood_pressure_mmhg

DBP > 120 mmHg

Diastolic blood pressure greater than 120 mm Hg

review_status=ok source_strategy=domain_source_numeric_leaf

- **[atomic / IS_TRUE / pediatric_severe_hypertension_threshold_condition_present]** Severe hypertension (SBP greater than 180 mm Hg or DBP greater than 120 mm Hg, or greater than the 95th percentile for age, sex, and height plus 30 mm Hg in pediatric patients [E] [F]) (2) (4) (17) (18)

Linked shared atomic definition: pediatric_severe_hypertension_threshold_condition_present -> atom.M083.admission.pediatric_severe_hypertension_threshold_condition_present

IS_TRUE

Severe hypertension (SBP greater than 180 mm Hg or DBP greater than 120 mm Hg, or greater than the 95th percentile for age, sex, and height plus 30 mm Hg in pediatric patients [E] [F]) (2) (4) (17) (18)

review_status=needs_review source_strategy=pediatric_placeholder

•

Path 3: Thrombolysis or thrombectomy performed or planned

- [atomic / IS_TRUE / thrombolysis_or_thrombectomy_performed_or_planned_condition_present] Thrombolysis or thrombectomy performed or planned [H] (2) (23) (24)

Linked shared atomic definition: thrombolysis_or_thrombectomy_performed_or_planned_condition_present -> atom.M083.admission.thrombolysis_or_thrombectomy_performed_or_planned_condition_present

IS_TRUE

Thrombolysis or thrombectomy performed or planned [H] (2) (23) (24)

review_status=ok source_strategy=domain_source_atomic_leaf