

MA 201: Partial Differential Equations

Lecture - 7

Change of coordinates: Canonical Transformations

Change of coordinates: Canonical Transformations

- Consider the following equation:

$$Au_{xx} + Bu_{xy} + Cu_{yy} + Du_x + Eu_y + Fu + G = 0. \quad (1)$$

Change of coordinates: Canonical Transformations

- Consider the following equation:

$$Au_{xx} + Bu_{xy} + Cu_{yy} + Du_x + Eu_y + Fu + G = 0. \quad (1)$$

- Equation (1) can be reduced to a simpler equation by introducing new coordinates:

Change of coordinates: Canonical Transformations

- Consider the following equation:

$$Au_{xx} + Bu_{xy} + Cu_{yy} + Du_x + Eu_y + Fu + G = 0. \quad (1)$$

- Equation (1) can be reduced to a simpler equation by introducing new coordinates:

$$\xi = \xi(x, y), \quad \eta = \eta(x, y).$$

Change of coordinates: Canonical Transformations

- Consider the following equation:

$$Au_{xx} + Bu_{xy} + Cu_{yy} + Du_x + Eu_y + Fu + G = 0. \quad (1)$$

- Equation (1) can be reduced to a simpler equation by introducing new coordinates:

$$\xi = \xi(x, y), \quad \eta = \eta(x, y).$$

ξ and η are known as **characteristic coordinates**.

- Compute the partial derivatives appearing in (1):

Change of coordinates: Canonical Transformations

- Consider the following equation:

$$Au_{xx} + Bu_{xy} + Cu_{yy} + Du_x + Eu_y + Fu + G = 0. \quad (1)$$

- Equation (1) can be reduced to a simpler equation by introducing new coordinates:

$$\xi = \xi(x, y), \quad \eta = \eta(x, y).$$

ξ and η are known as **characteristic coordinates**.

- Compute the partial derivatives appearing in (1):

$$u_x = u_\xi \xi_x + u_\eta \eta_x, \quad u_y = u_\xi \xi_y + u_\eta \eta_y,$$

$$u_{xx} = u_{\xi\xi} \xi_x^2 + 2u_{\xi\eta} \xi_x \eta_x + u_{\eta\eta} \eta_x^2 + u_\xi \xi_{xx} + u_\eta \eta_{xx},$$

$$u_{yy} = u_{\xi\xi} \xi_y^2 + 2u_{\xi\eta} \xi_y \eta_y + u_{\eta\eta} \eta_y^2 + u_\xi \xi_{yy} + u_\eta \eta_{yy},$$

$$u_{xy} = u_{\xi\xi} \xi_x \xi_y + u_{\xi\eta} (\xi_x \eta_y + \xi_y \eta_x) + u_{\eta\eta} \eta_x \eta_y + u_\xi \xi_{xy} + u_\eta \eta_{xy}.$$

- Substitute these values into (1) so as to obtain a new form of the PDE in the variables ξ and η :

$$\tilde{A}u_{\xi\xi} + \tilde{B}u_{\xi\eta} + \tilde{C}u_{\eta\eta} + \tilde{D}u_{\xi} + \tilde{E}u_{\eta} + Fu = G$$

In above, the new coefficients are as follows:

$$\tilde{A} = A\xi_x^2 + B\xi_x\xi_y + C\xi_y^2, \quad \tilde{B} = 2A\xi_x\eta_x + B(\xi_x\eta_y + \xi_y\eta_x) + 2C\xi_y\eta_y,$$

$$\tilde{C} = A\eta_x^2 + B\eta_x\eta_y + C\eta_y^2, \quad \tilde{D} = A\xi_{xx} + B\xi_{xy} + C\xi_{yy} + D\xi_x + E\xi_y,$$

$$\tilde{E} = A\eta_{xx} + B\eta_{xy} + C\eta_{yy} + D\eta_x + E\eta_y.$$

- Substitute these values into (1) so as to obtain a new form of the PDE in the variables ξ and η :

$$\tilde{A}u_{\xi\xi} + \tilde{B}u_{\xi\eta} + \tilde{C}u_{\eta\eta} + \tilde{D}u_{\xi} + \tilde{E}u_{\eta} + Fu = G$$

In above, the new coefficients are as follows:

$$\tilde{A} = A\xi_x^2 + B\xi_x\xi_y + C\xi_y^2, \quad \tilde{B} = 2A\xi_x\eta_x + B(\xi_x\eta_y + \xi_y\eta_x) + 2C\xi_y\eta_y,$$

$$\tilde{C} = A\eta_x^2 + B\eta_x\eta_y + C\eta_y^2, \quad \tilde{D} = A\xi_{xx} + B\xi_{xy} + C\xi_{yy} + D\xi_x + E\xi_y,$$

$$\tilde{E} = A\eta_{xx} + B\eta_{xy} + C\eta_{yy} + D\eta_x + E\eta_y.$$

- Whether the form of the PDE (1) remains invariant even after coordinate transformation?

- It can be observed that

$$\begin{pmatrix} 2\tilde{A} & \tilde{B} \\ \tilde{B} & 2\tilde{C} \end{pmatrix} = \begin{pmatrix} \xi_x & \xi_y \\ \eta_x & \eta_y \end{pmatrix} \begin{pmatrix} 2A & B \\ B & 2C \end{pmatrix} \begin{pmatrix} \xi_x & \eta_x \\ \xi_y & \eta_y \end{pmatrix}^t. \quad (2)$$

- It can be observed that

$$\begin{pmatrix} 2\tilde{A} & \tilde{B} \\ \tilde{B} & 2\tilde{C} \end{pmatrix} = \begin{pmatrix} \xi_x & \xi_y \\ \eta_x & \eta_y \end{pmatrix} \begin{pmatrix} 2A & B \\ B & 2C \end{pmatrix} \begin{pmatrix} \xi_x & \eta_x \\ \xi_y & \eta_y \end{pmatrix}^t. \quad (2)$$

- Taking the determinant on both sides gives

- It can be observed that

$$\begin{pmatrix} 2\tilde{A} & \tilde{B} \\ \tilde{B} & 2\tilde{C} \end{pmatrix} = \begin{pmatrix} \xi_x & \xi_y \\ \eta_x & \eta_y \end{pmatrix} \begin{pmatrix} 2A & B \\ B & 2C \end{pmatrix} \begin{pmatrix} \xi_x & \eta_x \\ \xi_y & \eta_y \end{pmatrix}^t. \quad (2)$$

- Taking the determinant on both sides gives

$$\tilde{B}^2 - 4\tilde{A}\tilde{C} = (\xi_x\eta_y - \xi_y\eta_x)^2(B^2 - 4AC) = J^2(B^2 - 4AC). \quad (3)$$

- It can be observed that

$$\begin{pmatrix} 2\tilde{A} & \tilde{B} \\ \tilde{B} & 2\tilde{C} \end{pmatrix} = \begin{pmatrix} \xi_x & \xi_y \\ \eta_x & \eta_y \end{pmatrix} \begin{pmatrix} 2A & B \\ B & 2C \end{pmatrix} \begin{pmatrix} \xi_x & \eta_x \\ \xi_y & \eta_y \end{pmatrix}^t. \quad (2)$$

- Taking the determinant on both sides gives

$$\tilde{B}^2 - 4\tilde{A}\tilde{C} = (\xi_x\eta_y - \xi_y\eta_x)^2(B^2 - 4AC) = J^2(B^2 - 4AC). \quad (3)$$

- J is the Jacobian of the transformation and we select the transformation (ξ, η) such that $J \neq 0$.

- It can be observed that

$$\begin{pmatrix} 2\tilde{A} & \tilde{B} \\ \tilde{B} & 2\tilde{C} \end{pmatrix} = \begin{pmatrix} \xi_x & \xi_y \\ \eta_x & \eta_y \end{pmatrix} \begin{pmatrix} 2A & B \\ B & 2C \end{pmatrix} \begin{pmatrix} \xi_x & \eta_x \\ \xi_y & \eta_y \end{pmatrix}^t. \quad (2)$$

- Taking the determinant on both sides gives

$$\tilde{B}^2 - 4\tilde{A}\tilde{C} = (\xi_x\eta_y - \xi_y\eta_x)^2(B^2 - 4AC) = J^2(B^2 - 4AC). \quad (3)$$

- J is the Jacobian of the transformation and we select the transformation (ξ, η) such that $J \neq 0$.
- Transformation given by $\xi = \xi(x, y)$ and $\eta = \eta(x, y)$ is called canonical transformation or characteristics and the reduced form of the PDE is called the canonical form.

Canonical Transformations: Hyperbolic PDE

- **Hyperbolic PDEs:** $B^2 - 4AC > 0$, which for the transformed equation is $\tilde{B}^2 - 4\tilde{A}\tilde{C} > 0$. (Look at (3))
 - **Natural choice:**

Canonical Transformations: Hyperbolic PDE

- **Hyperbolic PDEs:** $B^2 - 4AC > 0$, which for the transformed equation is $\tilde{B}^2 - 4\tilde{A}\tilde{C} > 0$. (Look at (3))
 - Natural choice: $\tilde{A} = 0$ and $\tilde{C} = 0$.

Canonical Transformations: Hyperbolic PDE

- **Hyperbolic PDEs:** $B^2 - 4AC > 0$, which for the transformed equation is $\tilde{B}^2 - 4\tilde{A}\tilde{C} > 0$. (Look at (3))
 - **Natural choice:** $\tilde{A} = 0$ and $\tilde{C} = 0$.
- That is $A\xi_x^2 + B\xi_x\xi_y + C\xi_y^2 = 0$ and $A\eta_x^2 + B\eta_x\eta_y + C\eta_y^2 = 0$.

Canonical Transformations: Hyperbolic PDE

- **Hyperbolic PDEs:** $B^2 - 4AC > 0$, which for the transformed equation is $\tilde{B}^2 - 4\tilde{A}\tilde{C} > 0$. (Look at (3))
 - Natural choice: $\tilde{A} = 0$ and $\tilde{C} = 0$.
- That is $A\xi_x^2 + B\xi_x\xi_y + C\xi_y^2 = 0$ and $A\eta_x^2 + B\eta_x\eta_y + C\eta_y^2 = 0$.
 - Leads to the algebraic equations:

$$A\lambda^2 + B\lambda + C = 0, \quad \lambda_1 = \frac{\xi_x}{\xi_y}, \quad \lambda_2 = \frac{\eta_x}{\eta_y}.$$

Canonical Transformations: Hyperbolic PDE

- **Hyperbolic PDEs:** $B^2 - 4AC > 0$, which for the transformed equation is $\tilde{B}^2 - 4\tilde{A}\tilde{C} > 0$. (Look at (3))
 - **Natural choice:** $\tilde{A} = 0$ and $\tilde{C} = 0$.
- That is $A\xi_x^2 + B\xi_x\xi_y + C\xi_y^2 = 0$ and $A\eta_x^2 + B\eta_x\eta_y + C\eta_y^2 = 0$.
 - **Leads to the algebraic equations:**

$$A\lambda^2 + B\lambda + C = 0, \quad \lambda_1 = \frac{\xi_x}{\xi_y}, \quad \lambda_2 = \frac{\eta_x}{\eta_y}.$$

- **Observations:** (What are ξ and η ?)

Canonical Transformations: Hyperbolic PDE

- **Hyperbolic PDEs:** $B^2 - 4AC > 0$, which for the transformed equation is $\tilde{B}^2 - 4\tilde{A}\tilde{C} > 0$. (Look at (3))
 - **Natural choice:** $\tilde{A} = 0$ and $\tilde{C} = 0$.
- That is $A\xi_x^2 + B\xi_x\xi_y + C\xi_y^2 = 0$ and $A\eta_x^2 + B\eta_x\eta_y + C\eta_y^2 = 0$.
 - **Leads to the algebraic equations:**

$$A\lambda^2 + B\lambda + C = 0, \quad \lambda_1 = \frac{\xi_x}{\xi_y}, \quad \lambda_2 = \frac{\eta_x}{\eta_y}.$$

► **Observations:** (What are ξ and η ?)

- $\xi(x, y) = c$

Canonical Transformations: Hyperbolic PDE

- **Hyperbolic PDEs:** $B^2 - 4AC > 0$, which for the transformed equation is $\tilde{B}^2 - 4\tilde{A}\tilde{C} > 0$. (Look at (3))
 - **Natural choice:** $\tilde{A} = 0$ and $\tilde{C} = 0$.
- That is $A\xi_x^2 + B\xi_x\xi_y + C\xi_y^2 = 0$ and $A\eta_x^2 + B\eta_x\eta_y + C\eta_y^2 = 0$.
 - **Leads to the algebraic equations:**

$$A\lambda^2 + B\lambda + C = 0, \quad \lambda_1 = \frac{\xi_x}{\xi_y}, \quad \lambda_2 = \frac{\eta_x}{\eta_y}.$$

► **Observations:** (What are ξ and η ?)

$$\begin{aligned} & \bullet \quad \xi(x, y) = c \qquad \bullet \text{For } \xi, \text{ solve } \frac{dy}{dx} = -\lambda_1(x, y) \\ \Rightarrow \quad & \xi_x + \xi_y \frac{dy}{dx} = 0, \end{aligned}$$

Canonical Transformations: Hyperbolic PDE

- **Hyperbolic PDEs:** $B^2 - 4AC > 0$, which for the transformed equation is $\tilde{B}^2 - 4\tilde{A}\tilde{C} > 0$. (Look at (3))
 - **Natural choice:** $\tilde{A} = 0$ and $\tilde{C} = 0$.
- That is $A\xi_x^2 + B\xi_x\xi_y + C\xi_y^2 = 0$ and $A\eta_x^2 + B\eta_x\eta_y + C\eta_y^2 = 0$.
 - **Leads to the algebraic equations:**

$$A\lambda^2 + B\lambda + C = 0, \quad \lambda_1 = \frac{\xi_x}{\xi_y}, \quad \lambda_2 = \frac{\eta_x}{\eta_y}.$$

► **Observations:** (What are ξ and η ?)

- $\xi(x, y) = c$
 - **For ξ , solve** $\frac{dy}{dx} = -\lambda_1(x, y)$
- $$\Rightarrow \quad \xi_x + \xi_y \frac{dy}{dx} = 0,$$
- **For η , solve** $\frac{dy}{dx} = -\lambda_2(x, y)$.

Canonical Transformations: Hyperbolic PDE

- **Hyperbolic PDEs:** $B^2 - 4AC > 0$, which for the transformed equation is $\tilde{B}^2 - 4\tilde{A}\tilde{C} > 0$. (Look at (3))
 - **Natural choice:** $\tilde{A} = 0$ and $\tilde{C} = 0$.
- That is $A\xi_x^2 + B\xi_x\xi_y + C\xi_y^2 = 0$ and $A\eta_x^2 + B\eta_x\eta_y + C\eta_y^2 = 0$.
 - **Leads to the algebraic equations:**

$$A\lambda^2 + B\lambda + C = 0, \quad \lambda_1 = \frac{\xi_x}{\xi_y}, \quad \lambda_2 = \frac{\eta_x}{\eta_y}.$$

► **Observations:** (What are ξ and η ?)

- $\xi(x, y) = c$ • **For ξ , solve** $\frac{dy}{dx} = -\lambda_1(x, y)$
- $\Rightarrow \quad \xi_x + \xi_y \frac{dy}{dx} = 0,$
- **For η , solve** $\frac{dy}{dx} = -\lambda_2(x, y).$
- Canonical form for hyperbolic equation can be written as

$$u_{\xi\eta} = \phi(\xi, \eta, u, u_\xi, u_\eta).$$

Hyperbolic Equation with Constant Coefficients

- Assume that either $A \neq 0$ or $C \neq 0$.

Hyperbolic Equation with Constant Coefficients

- Assume that either $A \neq 0$ or $C \neq 0$.
- What can you say when A and C vanish simultaneously?

Hyperbolic Equation with Constant Coefficients

- Assume that either $A \neq 0$ or $C \neq 0$.
- What can you say when A and C vanish simultaneously?
- For $A \neq 0$, canonical transformation (ξ, η) takes the form

$$\xi = y + \lambda_1 x, \quad \eta = y + \lambda_2 x, \quad \lambda_{1,2} = \frac{-B \pm \sqrt{B^2 - 4AC}}{2A}.$$

Hyperbolic Equation with Constant Coefficients

- Assume that either $A \neq 0$ or $C \neq 0$.
- What can you say when A and C vanish simultaneously?
- For $A \neq 0$, canonical transformation (ξ, η) takes the form

$$\xi = y + \lambda_1 x, \quad \eta = y + \lambda_2 x, \quad \lambda_{1,2} = \frac{-B \pm \sqrt{B^2 - 4AC}}{2A}.$$

► For $A \neq 0$ and $C = 0$, we have

$$\xi = y, \quad \eta = y - \frac{B}{A}x.$$

Hyperbolic Equation with Constant Coefficients

- Assume that either $A \neq 0$ or $C \neq 0$.
- What can you say when A and C vanish simultaneously?
- For $A \neq 0$, canonical transformation (ξ, η) takes the form

$$\xi = y + \lambda_1 x, \quad \eta = y + \lambda_2 x, \quad \lambda_{1,2} = \frac{-B \pm \sqrt{B^2 - 4AC}}{2A}.$$

► For $A \neq 0$ and $C = 0$, we have

$$\xi = y, \quad \eta = y - \frac{B}{A}x.$$

- What are the characteristics when $A = 0$ and $C \neq 0$?

Example

Find the characteristics of the following equation and reduce it to the appropriate standard form and then obtain the general solution:

$$3u_{xx} + 10u_{xy} + 3u_{yy} = 0.$$

Example

Find the characteristics of the following equation and reduce it to the appropriate standard form and then obtain the general solution:

$$3u_{xx} + 10u_{xy} + 3u_{yy} = 0.$$

- We have $\mathcal{D} = B^2 - 4CA = 100 - 36 = 64 > 0$.

Example

Find the characteristics of the following equation and reduce it to the appropriate standard form and then obtain the general solution:

$$3u_{xx} + 10u_{xy} + 3u_{yy} = 0.$$

- We have $\mathcal{D} = B^2 - 4CA = 100 - 36 = 64 > 0$.
- The equation is of hyperbolic type.

Example

Find the characteristics of the following equation and reduce it to the appropriate standard form and then obtain the general solution:

$$3u_{xx} + 10u_{xy} + 3u_{yy} = 0.$$

- We have $\mathcal{D} = B^2 - 4CA = 100 - 36 = 64 > 0$.
- The equation is of hyperbolic type.
- The characteristics are given by

Example

Find the characteristics of the following equation and reduce it to the appropriate standard form and then obtain the general solution:

$$3u_{xx} + 10u_{xy} + 3u_{yy} = 0.$$

- We have $\mathcal{D} = B^2 - 4CA = 100 - 36 = 64 > 0$.
- The equation is of hyperbolic type.
- The characteristics are given by

$$\begin{aligned}\xi &= y + \left(\frac{-B + \sqrt{\mathcal{D}}}{2A} \right) x = y - (1/3)x, \\ \eta &= y + \left(\frac{-B - \sqrt{\mathcal{D}}}{2A} \right) x = y - 3x.\end{aligned}$$

- Recall the canonical form

$$\tilde{A}u_{\xi\xi} + \tilde{B}u_{\xi\eta} + \tilde{C}u_{\eta\eta} + \tilde{D}u_{\xi} + \tilde{E}u_{\eta} + Fu = G$$

with

$$\tilde{A} = A\xi_x^2 + B\xi_x\xi_y + C\xi_y^2, \quad \tilde{B} = 2A\xi_x\eta_x + B(\xi_x\eta_y + \xi_y\eta_x) + 2C\xi_y\eta_y,$$

$$\tilde{C} = A\eta_x^2 + B\eta_x\eta_y + C\eta_y^2, \quad \tilde{D} = A\xi_{xx} + B\xi_{xy} + C\xi_{yy} + D\xi_x + E\xi_y,$$

$$\tilde{E} = A\eta_{xx} + B\eta_{xy} + C\eta_{yy} + D\eta_x + E\eta_y.$$

- Due to the choice of canonical transformation, we have

- Recall the canonical form

$$\tilde{A}u_{\xi\xi} + \tilde{B}u_{\xi\eta} + \tilde{C}u_{\eta\eta} + \tilde{D}u_{\xi} + \tilde{E}u_{\eta} + Fu = G$$

with

$$\tilde{A} = A\xi_x^2 + B\xi_x\xi_y + C\xi_y^2, \quad \tilde{B} = 2A\xi_x\eta_x + B(\xi_x\eta_y + \xi_y\eta_x) + 2C\xi_y\eta_y,$$

$$\tilde{C} = A\eta_x^2 + B\eta_x\eta_y + C\eta_y^2, \quad \tilde{D} = A\xi_{xx} + B\xi_{xy} + C\xi_{yy} + D\xi_x + E\xi_y,$$

$$\tilde{E} = A\eta_{xx} + B\eta_{xy} + C\eta_{yy} + D\eta_x + E\eta_y.$$

- Due to the choice of canonical transformation, we have $\tilde{A} = 0 = \tilde{C}$.
- Further, it is easy to see that

- Recall the canonical form

$$\tilde{A}u_{\xi\xi} + \tilde{B}u_{\xi\eta} + \tilde{C}u_{\eta\eta} + \tilde{D}u_{\xi} + \tilde{E}u_{\eta} + Fu = G$$

with

$$\tilde{A} = A\xi_x^2 + B\xi_x\xi_y + C\xi_y^2, \quad \tilde{B} = 2A\xi_x\eta_x + B(\xi_x\eta_y + \xi_y\eta_x) + 2C\xi_y\eta_y,$$

$$\tilde{C} = A\eta_x^2 + B\eta_x\eta_y + C\eta_y^2, \quad \tilde{D} = A\xi_{xx} + B\xi_{xy} + C\xi_{yy} + D\xi_x + E\xi_y,$$

$$\tilde{E} = A\eta_{xx} + B\eta_{xy} + C\eta_{yy} + D\eta_x + E\eta_y.$$

- Due to the choice of canonical transformation, we have $\tilde{A} = 0 = \tilde{C}$.
- Further, it is easy to see that $\tilde{D} = 0 = \tilde{E} = F$ and $\tilde{B} = -18$.
- Thus, the desired canonical form is

$$u_{\xi\eta} = 0.$$

- Recall the canonical form

$$\tilde{A}u_{\xi\xi} + \tilde{B}u_{\xi\eta} + \tilde{C}u_{\eta\eta} + \tilde{D}u_{\xi} + \tilde{E}u_{\eta} + Fu = G$$

with

$$\tilde{A} = A\xi_x^2 + B\xi_x\xi_y + C\xi_y^2, \quad \tilde{B} = 2A\xi_x\eta_x + B(\xi_x\eta_y + \xi_y\eta_x) + 2C\xi_y\eta_y,$$

$$\tilde{C} = A\eta_x^2 + B\eta_x\eta_y + C\eta_y^2, \quad \tilde{D} = A\xi_{xx} + B\xi_{xy} + C\xi_{yy} + D\xi_x + E\xi_y,$$

$$\tilde{E} = A\eta_{xx} + B\eta_{xy} + C\eta_{yy} + D\eta_x + E\eta_y.$$

- Due to the choice of canonical transformation, we have $\tilde{A} = 0 = \tilde{C}$.
- Further, it is easy to see that $\tilde{D} = 0 = \tilde{E} = F$ and $\tilde{B} = -18$.
- Thus, the desired canonical form is

$$u_{\xi\eta} = 0.$$

- Integrating the above first with respect to ξ , we get $u_{\eta} = f'(\eta)$. Integrating now with respect to η , we get

$$u = \mathcal{F}(\eta) + \mathcal{G}(\xi) = \mathcal{F}(y - 3x) + \mathcal{G}(3y - x),$$

where \mathcal{F} and \mathcal{G} are arbitrary functions.

Canonical Transformations: Parabolic PDE

- Parabolic PDEs: $B^2 - 4AC = 0$

Canonical Transformations: Parabolic PDE

- Parabolic PDEs: $B^2 - 4AC = 0$

► $A\lambda^2 + B\lambda + C = 0$ has two equal roots

$$\lambda = \lambda_1 = \frac{\xi_x}{\xi_y} = \lambda_2 = \frac{\eta_x}{\eta_y}.$$

Canonical Transformations: Parabolic PDE

- Parabolic PDEs: $B^2 - 4AC = 0$

► $A\lambda^2 + B\lambda + C = 0$ has two equal roots

$$\lambda = \lambda_1 = \frac{\xi_x}{\xi_y} = \lambda_2 = \frac{\eta_x}{\eta_y}.$$

► Thus, either $\tilde{A} = 0$ or $\tilde{C} = 0$

Canonical Transformations: Parabolic PDE

- Parabolic PDEs: $B^2 - 4AC = 0$

► $A\lambda^2 + B\lambda + C = 0$ has two equal roots

$$\lambda = \lambda_1 = \frac{\xi_x}{\xi_y} = \lambda_2 = \frac{\eta_x}{\eta_y}.$$

► Thus, either $\tilde{A} = 0$ or $\tilde{C} = 0$

► Necessary Requirement $\tilde{B} = 0$

Canonical Transformations: Parabolic PDE

- Parabolic PDEs: $B^2 - 4AC = 0$

► $A\lambda^2 + B\lambda + C = 0$ has two equal roots

$$\lambda = \lambda_1 = \frac{\xi_x}{\xi_y} = \lambda_2 = \frac{\eta_x}{\eta_y}.$$

► Thus, either $\tilde{A} = 0$ or $\tilde{C} = 0$

► Necessary Requirement $\tilde{B} = 0$

Canonical Transformations: Parabolic PDE

- Parabolic PDEs: $B^2 - 4AC = 0$

► $A\lambda^2 + B\lambda + C = 0$ has two equal roots

$$\lambda = \lambda_1 = \frac{\xi_x}{\xi_y} = \lambda_2 = \frac{\eta_x}{\eta_y}.$$

► Thus, either $\tilde{A} = 0$ or $\tilde{C} = 0$

► Necessary Requirement $\tilde{B} = 0$

- We obtain only a single characteristic/canonical transformation as

$$\tilde{A} = 0 \text{ gives } \lambda = \frac{\xi_x}{\xi_y} = -\frac{dy}{dx}, \text{ i.e., choice of } \xi \text{ OR}$$

Canonical Transformations: Parabolic PDE

- Parabolic PDEs: $B^2 - 4AC = 0$

► $A\lambda^2 + B\lambda + C = 0$ has two equal roots

$$\lambda = \lambda_1 = \frac{\xi_x}{\xi_y} = \lambda_2 = \frac{\eta_x}{\eta_y}.$$

► Thus, either $\tilde{A} = 0$ or $\tilde{C} = 0$

► Necessary Requirement $\tilde{B} = 0$

- We obtain only a single characteristic/canonical transformation as

$$\tilde{A} = 0 \text{ gives } \lambda = \frac{\xi_x}{\xi_y} = -\frac{dy}{dx}, \text{ i.e., choice of } \xi \text{ **OR**}$$

$$\tilde{C} = 0 \text{ gives } \lambda = \frac{\eta_x}{\eta_y} = -\frac{dy}{dx}, \text{ i.e., choice of } \eta.$$

- Suppose ξ is calculated. What could be the possible choice for η ?

- Suppose ξ is calculated. What could be the possible choice for η ?
- We select η as any arbitrary function of x and y such that η is independent of ξ , i.e.,

$$J = \begin{vmatrix} \xi_x & \xi_y \\ \eta_x & \eta_y \end{vmatrix} \neq 0.$$

- Suppose ξ is calculated. What could be the possible choice for η ?
- We select η as any arbitrary function of x and y such that η is independent of ξ , i.e.,

$$J = \begin{vmatrix} \xi_x & \xi_y \\ \eta_x & \eta_y \end{vmatrix} \neq 0.$$

- For example, if $\xi = x + y$, we may select $\eta = y$ or $\eta = x$.

- Suppose ξ is calculated. What could be the possible choice for η ?
- We select η as any arbitrary function of x and y such that η is independent of ξ , i.e.,

$$J = \begin{vmatrix} \xi_x & \xi_y \\ \eta_x & \eta_y \end{vmatrix} \neq 0.$$

- For example, if $\xi = x + y$, we may select $\eta = y$ or $\eta = x$.
- Canonical form for Parabolic Equation:

- Suppose ξ is calculated. What could be the possible choice for η ?
- We select η as any arbitrary function of x and y such that η is independent of ξ , i.e.,

$$J = \begin{vmatrix} \xi_x & \xi_y \\ \eta_x & \eta_y \end{vmatrix} \neq 0.$$

- For example, if $\xi = x + y$, we may select $\eta = y$ or $\eta = x$.

► Canonical form for Parabolic Equation:

$$u_{\eta\eta} = \phi(\xi, \eta, u, u_\xi, u_\eta), \text{ OR}$$

- Suppose ξ is calculated. What could be the possible choice for η ?
- We select η as any arbitrary function of x and y such that η is independent of ξ , i.e.,

$$J = \begin{vmatrix} \xi_x & \xi_y \\ \eta_x & \eta_y \end{vmatrix} \neq 0.$$

- For example, if $\xi = x + y$, we may select $\eta = y$ or $\eta = x$.

► Canonical form for Parabolic Equation:

$$\begin{aligned} u_{\eta\eta} &= \phi(\xi, \eta, u, u_\xi, u_\eta), \text{ OR} \\ u_{\xi\xi} &= \phi(\xi, \eta, u, u_\xi, u_\eta). \end{aligned}$$

- Suppose ξ is calculated. What could be the possible choice for η ?
- We select η as any arbitrary function of x and y such that η is independent of ξ , i.e.,

$$J = \begin{vmatrix} \xi_x & \xi_y \\ \eta_x & \eta_y \end{vmatrix} \neq 0.$$

- For example, if $\xi = x + y$, we may select $\eta = y$ or $\eta = x$.

► Canonical form for Parabolic Equation:

$$\begin{aligned} u_{\eta\eta} &= \phi(\xi, \eta, u, u_\xi, u_\eta), \text{ OR} \\ u_{\xi\xi} &= \phi(\xi, \eta, u, u_\xi, u_\eta). \end{aligned}$$

Example

Find the characteristics of the following equation and reduce it to the appropriate standard form and then obtain the general solution:

$$u_{xx} + 4u_{xy} + 4u_{yy} = 0.$$

- We have $B^2 - 4CA = 0$. Hence the equation is parabolic.

- We have $B^2 - 4CA = 0$. Hence the equation is parabolic.
- The characteristics are given by

$$\xi = y + \left(\frac{-B + \sqrt{\mathcal{D}}}{2A} \right) x = y - 2x, \quad \eta = y.$$

- We have $B^2 - 4CA = 0$. Hence the equation is parabolic.
- The characteristics are given by

$$\xi = y + \left(\frac{-B + \sqrt{D}}{2A} \right) x = y - 2x, \quad \eta = y.$$

- Recall the canonical form

$$\tilde{A}u_{\xi\xi} + \tilde{B}u_{\xi\eta} + \tilde{C}u_{\eta\eta} + \tilde{D}u_{\xi} + \tilde{E}u_{\eta} + Fu = G$$

with

$$\tilde{A} = A\xi_x^2 + B\xi_x\xi_y + C\xi_y^2, \quad \tilde{B} = 2A\xi_x\eta_x + B(\xi_x\eta_y + \xi_y\eta_x) + 2C\xi_y\eta_y,$$

$$\tilde{C} = A\eta_x^2 + B\eta_x\eta_y + C\eta_y^2, \quad \tilde{D} = A\xi_{xx} + B\xi_{xy} + C\xi_{yy} + D\xi_x + E\xi_y,$$

$$\tilde{E} = A\eta_{xx} + B\eta_{xy} + C\eta_{yy} + D\eta_x + E\eta_y.$$

- We have $B^2 - 4CA = 0$. Hence the equation is parabolic.
- The characteristics are given by

$$\xi = y + \left(\frac{-B + \sqrt{D}}{2A} \right) x = y - 2x, \quad \eta = y.$$

- Recall the canonical form

$$\tilde{A}u_{\xi\xi} + \tilde{B}u_{\xi\eta} + \tilde{C}u_{\eta\eta} + \tilde{D}u_{\xi} + \tilde{E}u_{\eta} + Fu = G$$

with

$$\tilde{A} = A\xi_x^2 + B\xi_x\xi_y + C\xi_y^2, \quad \tilde{B} = 2A\xi_x\eta_x + B(\xi_x\eta_y + \xi_y\eta_x) + 2C\xi_y\eta_y,$$

$$\tilde{C} = A\eta_x^2 + B\eta_x\eta_y + C\eta_y^2, \quad \tilde{D} = A\xi_{xx} + B\xi_{xy} + C\xi_{yy} + D\xi_x + E\xi_y,$$

$$\tilde{E} = A\eta_{xx} + B\eta_{xy} + C\eta_{yy} + D\eta_x + E\eta_y.$$

- Due to the choice of canonical transformation, we have $\tilde{A} = 0 = \tilde{B}$.

- Due to the choice of canonical transformation, we have $\tilde{A} = 0 = \tilde{B}$.
- Further, it is easy to see that $\tilde{D} = 0 = \tilde{E} = F$ and $\tilde{C} = 4$.

- Due to the choice of canonical transformation, we have $\tilde{A} = 0 = \tilde{B}$.
- Further, it is easy to see that $\tilde{D} = 0 = \tilde{E} = F$ and $\tilde{C} = 4$.
- Thus, the desired canonical form is

$$u_{\eta\eta} = 0.$$

- Due to the choice of canonical transformation, we have $\tilde{A} = 0 = \tilde{B}$.
- Further, it is easy to see that $\tilde{D} = 0 = \tilde{E} = F$ and $\tilde{C} = 4$.
- Thus, the desired canonical form is

$$u_{\eta\eta} = 0.$$

- Integrating it twice in succession with respect to η , we will get

$$u = \eta\mathcal{F}(\xi) + \mathcal{G}(\xi) = y\mathcal{F}(y - 2x) + \mathcal{G}(y - 2x),$$

where \mathcal{F} and \mathcal{G} are arbitrary functions.

Canonical Transformations: Elliptic PDE

- Elliptic PDEs: $B^2 - 4AC < 0$
 - $A\lambda^2 + B\lambda + C = 0$ has distinct roots λ_1 and λ_2 .
 - Leads to complex conjugate canonical transformation ξ and η .

Canonical Transformations: Elliptic PDE

- Elliptic PDEs: $B^2 - 4AC < 0$
 - $A\lambda^2 + B\lambda + C = 0$ has distinct roots λ_1 and λ_2 .
 - Leads to complex conjugate canonical transformation ξ and η .
- Since ξ and η are complex, we introduce new real variables

$$\alpha = \frac{1}{2}(\xi + \eta), \quad \beta = \frac{1}{2i}(\xi - \eta),$$

so that

$$\xi = \alpha + i\beta, \quad \eta = \alpha - i\beta.$$

- Under the transformation $(x, y) \rightarrow (\alpha, \beta)$, the canonical form is given by

$$u_{\alpha\alpha} + u_{\beta\beta} = \Phi(\alpha, \beta, u, u_{\alpha}, u_{\beta}),$$

which still contains two second-order derivatives.

- Under the transformation $(x, y) \rightarrow (\alpha, \beta)$, the canonical form is given by

$$u_{\alpha\alpha} + u_{\beta\beta} = \Phi(\alpha, \beta, u, u_{\alpha}, u_{\beta}),$$

which still contains two second-order derivatives.

- Note that method of characteristics is not found suitable for elliptic equations since even after using the transformation, that is, the characteristics (in new variables), the equation gets reduced to Laplacian form only.

- Under the transformation $(x, y) \rightarrow (\alpha, \beta)$, the canonical form is given by

$$u_{\alpha\alpha} + u_{\beta\beta} = \Phi(\alpha, \beta, u, u_{\alpha}, u_{\beta}),$$

which still contains two second-order derivatives.

- Note that method of characteristics is not found suitable for elliptic equations since even after using the transformation, that is, the characteristics (in new variables), the equation gets reduced to Laplacian form only.
- In other words: the given equation gets reduced marginally only with two double derivatives still remaining.

- Under the transformation $(x, y) \rightarrow (\alpha, \beta)$, the canonical form is given by

$$u_{\alpha\alpha} + u_{\beta\beta} = \Phi(\alpha, \beta, u, u_{\alpha}, u_{\beta}),$$

which still contains two second-order derivatives.

- Note that method of characteristics is not found suitable for elliptic equations since even after using the transformation, that is, the characteristics (in new variables), the equation gets reduced to Laplacian form only.
- In other words: the given equation gets reduced marginally only with two double derivatives still remaining.
- For this reason, the method of characteristics is usually not applied to elliptic equations.

Example

Find the characteristics of the following equation and reduce it to the appropriate standard form and then obtain the general solution:

$$u_{xx} - 4u_{xy} + 4u_{yy} = \cos(2x + y).$$

Example

Find the characteristics of the following equation and reduce it to the appropriate standard form and then obtain the general solution:

$$u_{xx} - 4u_{xy} + 4u_{yy} = \cos(2x + y).$$

- The given equation is a parabolic one.

Example

Find the characteristics of the following equation and reduce it to the appropriate standard form and then obtain the general solution:

$$u_{xx} - 4u_{xy} + 4u_{yy} = \cos(2x + y).$$

- The given equation is a parabolic one.
- The characteristics are:

$$\xi = x, \quad \eta = x - \frac{B}{2C}y = x + (1/2)y$$

Example

Find the characteristics of the following equation and reduce it to the appropriate standard form and then obtain the general solution:

$$u_{xx} - 4u_{xy} + 4u_{yy} = \cos(2x + y).$$

- The given equation is a parabolic one.
- The characteristics are:

$$\xi = x, \quad \eta = x - \frac{B}{2C}y = x + (1/2)y$$

- The canonical form will be $u_{\xi\xi} = \cos \eta$.

Example

Find the characteristics of the following equation and reduce it to the appropriate standard form and then obtain the general solution:

$$u_{xx} - 4u_{xy} + 4u_{yy} = \cos(2x + y).$$

- The given equation is a parabolic one.
- The characteristics are:

$$\xi = x, \quad \eta = x - \frac{B}{2C}y = x + (1/2)y$$

- The canonical form will be $u_{\xi\xi} = \cos \eta$.
- Integrating partially with respect to ξ :

Example

Find the characteristics of the following equation and reduce it to the appropriate standard form and then obtain the general solution:

$$u_{xx} - 4u_{xy} + 4u_{yy} = \cos(2x + y).$$

- The given equation is a parabolic one.
- The characteristics are:

$$\xi = x, \quad \eta = x - \frac{B}{2C}y = x + (1/2)y$$

- The canonical form will be $u_{\xi\xi} = \cos \eta$.
- Integrating partially with respect to ξ :

$$u_{\xi} = \xi \cos \eta + f(\eta).$$

- Again integrating w.r.t. ξ

$$u = \frac{\xi^2}{2} \cos \eta + \xi f(\eta) + g(\eta) = \frac{x^2}{2} \cos(2x+y) + xf(2x+y) + g(2x+y).$$

Classification

- Finding the nature of a PDE depends on finding an algebraic equation

$$A\lambda^2 + B\lambda + C = 0$$

and examining the nature of the roots.

Classification

- Finding the nature of a PDE depends on finding an algebraic equation

$$A\lambda^2 + B\lambda + C = 0$$

and examining the nature of the roots.

- When the roots λ are real and distinct, we get specific characteristics and the equation is called **hyperbolic**.

Classification

- Finding the nature of a PDE depends on finding an algebraic equation

$$A\lambda^2 + B\lambda + C = 0$$

and examining the nature of the roots.

- When the roots λ are real and distinct, we get specific characteristics and the equation is called **hyperbolic**.
- When the roots λ are real but equal, we get specific characteristics and the equation is called **parabolic**.

Classification

- Finding the nature of a PDE depends on finding an algebraic equation

$$A\lambda^2 + B\lambda + C = 0$$

and examining the nature of the roots.

- When the roots λ are real and distinct, we get specific characteristics and the equation is called **hyperbolic**.
- When the roots λ are real but equal, we get specific characteristics and the equation is called **parabolic**.
- When the roots λ are real complex (appearing as complex conjugates), we get specific characteristics and the equation is called **elliptic**.

Classification

- Finding the nature of a PDE depends on finding an algebraic equation

$$A\lambda^2 + B\lambda + C = 0$$

and examining the nature of the roots.

- When the roots λ are real and distinct, we get specific characteristics and the equation is called **hyperbolic**.
- When the roots λ are real but equal, we get specific characteristics and the equation is called **parabolic**.
- When the roots λ are real complex (appearing as complex conjugates), we get specific characteristics and the equation is called **elliptic**.
- That is, the classification of such PDEs is simply based on the roots of an algebraic equation - the way we do for conic section.