# **Genome Editing and Engineering Course No: BT-637**



### **LECTURE-4**

Dr. Kusum K. Singh
Department of Biosciences and Bioengineering
Indian Institute of Technology Guwahati

### Introduction

- Fokl (Flavobacterium okeanokoites)
- Functional domains of Fokl (N and C-terminus)
- Mutate the Fokl to study its modular nature

Gene, 133 (1993) 79-84 © 1993 Elsevier Science Publishers B.V. All rights reserved. 0378-1119/93/\$06.00

**GENE 07304** 

#### **Short Communications**

### C-terminal deletion mutants of the FokI restriction endonuclease

(Catalytic domain; Flavobacterium okeanokoites; hydroxyl radical footprinting; protein-DNA interaction; recognition domain)

#### Lin Li, Louisa P. Wu, Robert Clarke and Srinivasan Chandrasegaran

Department of Environmental Health Sciences, The Johns Hopkins University, School of Hygiene and Public Health, Baltimore, MD 21205, USA

Received by A.J. Podhajska: 4 April 1993; Revised/Accepted: 18 May/19 May 1993; Received at publishers: 27 May 1993

### **C-terminal deletion mutants**





Fig. 1. Construction and analysis of MP. (A) Sequences of the 5' and 3' primers used to construct the C-terminal deletion mutants of FokI. All primers are flanked by BamHI sites. New translational signals were introduced to achieve high expression of the mutant proteins. A 7-bp spacer separates the RBS for E. coli from the ATG start codon. (B) 0.1% SDS-12% PAGE profile of the purified 41-kDa and 30-kDa MP. Lanes: 1, protein standards (from BRL, Gaithersburg, MD, USA); 2, wt FokI ENase; 3, the 41-kDa MP; 4, the 30-kDa MP. The procedures for cell growth and purification of the MP are similar to the one described elsewhere (Li and Chandrasegaran, 1993).

## **Trypsin digestion pattern**




Trypsin digestion of the 41-kDa protein in the absence


Trypsin digestion of the 41-kDa MP in the presence

# **Trypsin Map**


II II



FokI endonuclease




41-kDa mutant protein




30 kDa &11 kDa associate to reconstitute the Fokl recognition domain

### **Affinities of wt & mt for DNA**



Deletion of C-terminal, 25 kDa portion has little or no effect

### Mode of interaction of wt & mt



- 1, untreated DNA substrate
- 2, DNA substrate cleaved by hydroxyl radicals;
- 3 and 4, presence of 7.5 pmoles and 15 pmoles FokI;
- 5 and 6, presence of 9 pmoles and 18 pmoles 41-kDa MP.

- Wt & mt interact only parts of recognition site
- No protection at cleavage site

### **Conclusions of C-ter. article**

- C-terminal mutants were generated
- 41 kDa binds seq. specific manner (wt)
- 30 kDa no longer bind DNA
- 30+11 reconstitutes to restore specificity
- No protection at the cleavage site

Proc. Natl. Acad. Sci. USA Vol. 90, pp. 9596–9600, October 1993 Biochemistry

# Single amino acid substitutions uncouple the DNA binding and strand scission activities of *Fok* I endonuclease

DAVID S. WAUGH\* AND ROBERT T. SAUER


Department of Biology, Massachusetts Institute of Technology, Cambridge, MA 02139

Communicated by Hamilton O. Smith, July 22, 1993 (received for review May 18, 1993)

D450A; D467A

### Structures comparison


(Pro-Asp-Xaa<sub>15-19</sub>-Glu/Asp-Xaa-Lys).



D450A; D467A

FIG. 2. Segments of amino acid sequences of *EcoRI* (residues 88–113), *EcoRV* (residues 71–92), and *Fok I* (residues 447–469) endonucleases are aligned for comparison. Identical residues are indicated by solid circles; conservative substitutions are identified by open circles. Gaps are indicated by dashes. The consensus sequence of the proposed active site motif (16) is shown on the top line.

# **Protection Experiments**




# Known facts till (1993)

- Fokl (Flavobacterium okeanokoites), Type IIS
- Recognizes the asymmetric sequence GGATG
- Staggered DNA cuts 9/13 bases away
- It appear to function as monomers
- By contrast other type II (EcoRI & EcoRV) acts as dimers
- How both strands are cleaved by Fokl?

# Hypothesis about cleavage

- Single molecule = two active sites
- Single molecule = one active site (use twice
- Mutational analysis = one catalytic center

- How monomeric Fokl
- Single catalytic center
- Cleave both strands of DNA?



#### Insertion and Deletion Mutants of FokI Restriction Endonuclease\*

(Received for publication, August 5, 1994, and in revised form, October 11, 1994)

#### Yang-Gyun Kim, Lin Li, and Srinivasan Chandrasegaran‡

From the Department of Environmental Health Sciences, The Johns Hopkins University, School of Hygiene and Public Health, Baltimore, Maryland 21205-2179

However, compared with the wild-type enzyme, the insertion mutants cleave predominantly one nucleotide further away from the recognition site on both strands of the DNA substrate. 9/13 10/14

The DNA binding and cleavage domains of FokI appear to be linked by a relatively malleable linker. No simple linear relationship exists between the linker length and the distance of the cut site from the recognition site.

#### Insertion and Deletion Mutants of FokI Restriction Endonuclease\*

(Received for publication, August 5, 1994, and in revised form, October 11, 1994)

#### Yang-Gyun Kim, Lin Li, and Srinivasan Chandrasegaran‡

From the Department of Environmental Health Sciences, The Johns Hopkins University, School of Hygiene and Public Health, Baltimore, Maryland 21205-2179


Amino acid sequence at the insertion/deletion site <sup>a</sup> areQLVKSELRHKLK	amino acid inser	amino acid insertion $Activity^b$	
	-7	-	
QLVEEKKSELRHKLK	-4	+	
QLVKSELEEKKSELRHKLK	wt	+	
QLVTAELKSELEEKKSELRHKLK	+4	+	
QLVTAELTAELKSELEEKKSELRHKLK	+8	+	
QLVTAELTAELKSELEEKKSELRHKLK	+12	+	
QLVTAELTAELKSELKSELEEKKSELEEKKSELRHKLK	+19	+	
QLVTAELTAELTAELKSELKSELEEKKSELEEKKSELRHKLK	. +23	+	
QLRGGGGGGGGGGGGQLVKSELEEKKSELRHKLK	+18°	+	


#### Insertion and Deletion Mutants of FokI Restriction Endonuclease\*

(Received for publication, August 5, 1994, and in revised form, October 11, 1994)

#### Yang-Gyun Kim, Lin Li, and Srinivasan Chandrasegaran‡

From the Department of Environmental Health Sciences, The Johns Hopkins University, School of Hygiene and Public Health, Baltimore, Maryland 21205-2179





### **Conclusions of Lecture-4**

- Uncouple the recognition & cleavage domain
- Domains protect recognition site
- Single catalytic center
- How both strands are cleaved??

# Thank You!