



# Session 3.2 - SNP matrix and phylogenetics

#### <u>BU-ISCIII</u> <u>Unidades Comunes Científico Técnicas - SGSAFI-ISCIII</u>

28-31 Octubre 2024, 6ª Edición Programa Formación Continua, ISCIII





#### Index


#### SNP matrix and Phylogenetics:

- SNP Matrix build
- Phylogeny
  - Maximum Likelihood
  - Parsimony
- WGS-Outbreaker
- iTOL phylogenetic tree visualization.
- How to interpret SNP-based outbreak analysis
- Examples: SnapperDB and GenomeTrakr





## Building a SNP matrix



Sample n





## Building a SNP matrix

• Once we have our multisample vcf:

#CHROM	POS	ID	REF	ALT	QUAL	FILTER	INFO	FORMAT	RA-L2073	RA-2805
NC_021827.1	276	9.	С	Α	291.68	PASS	AC=1;	GT:AD:DP:	0:13,0:13:	1:0,50:30
NC_021827.1	731		Α	G	2313.68	PASS	AC=1;	GT:AD:DP:	0:23,0:23:	1:0,10:10
NC_021827.1	921		С	T	1841.68	PASS	AC=1;	GT:AD:DP:	0:53,0:53:	0:20,0:20

We can generate the genotype for each sample


#CHROM	POS	RA-L2073	RA-2805
NC_021827.1	276	С	Α
NC_021827.1	731	A	G
NC_021827.1	921	С	С





## Building a SNP matrix

- So... now we have a simple multifasta, where each nucleotide represents a SNP.
- This means that even the nucleotide positions are sequentially in the fasta, they don't have to be near each other in the genome!
- The SNP matrix file will look like this:








## Phylogeny


#### **SNP** matrix







## Maximum parsimony




- Search the most parsimonious tree
- The most simple hipothesis must be the correct.
- Search the tree that explains the relationships with the less changes possible.






### Maximum Likelihood

- Searchs the most likely tree given the data and based in a evolutionary model.
- More sofisticated.
- Not prepared a priori for snp matrix.
- RAxML
  - Heterogenity rate disabled.
  - Branchs indicate the expected number of substitution per site.





- 0,1 = differences along that branch
- Which hypothesis is more likely, given that the change is rare?





#### SNP distance


- SNP distance is calculated with N model. Simply the number of sites that differ between each pair of sequences.
- By default sites with at least one missing data is deleted for all sequences in R
  (complete deletion option in MEGA).

dis_matrix.names	RA.L2073	RA.L2281	RA.L2327	RA.L2391	RA.L2450	RA.L2677	RA.L2701
RA-L2073	0	9403	9028	80	46	46	49
RA-L2281	9403	0	8777	9415	9397	9397	9402
RA-L2327	9028	8777	0	9040	9022	9022	9027
RA-L2391	80	9415	9040	0	74	74	79
RA-L2450	46	9397	9022	74	0	38	45
RA-L2677	46	9397	9022	74	38	0	45
RA-L2701	49	9402	9027	79	45	45	0
RA-L2782	9120	9183	4277	9132	9114	9114	9119
RA-L2805	4	9403	9028	80	46	46	49
RA-L2978	2	9401	9026	78	44	44	47





## WGS-Outbreaker










#### iTOL



https://itol.embl.de/





#### Combination of:

- SNP counts
- Tree topologies
- Bootstrap support






Organism	Maximum SNP count (number)	М	aximum SNP count	Reference	
		<21	21-100	>100	
E. coli	4	X			Underwood et al., 2013
E. coli	15	X			Eppinger et al., 2011
L. monocytogenes	9	X			Chen et al., 2017c
monocytogenes	12	X			Chen et al., 2017a
monocytogenes	18	X			Li et al., 2017
monocytogenes	20	X			Wang et al., 2015
monocytogenes	21		X		Nielsen et al., 2017
L. monocytogenes	28		X		Gilmour et al., 2010
L. monocytogenes	29		X		Chen et al., 2017b
L. monocytogenes	42		X		Chen et al., 2016
L. monocytogenes	67		X		Jackson et al., 2016
S. enterica	2	X			Wuyts et al., 2015
S. enterica	3	X			Allard et al., 2016
S. enterica	3	X			Taylor et al., 2015
S. enterica	6	X			Hoffmann et al., 2016
S. enterica	12	X			Octavia et al., 2015
S. enterica	30		X		Leekitcharoenphon et al.,

The maximum SNP counts for isolates that were traced back to the same source in the original study are presented. Whether the maximum SNP counts are less than 21 SNPs, 21 to 100 SNP, or greater than 100 SNPs is also indicated.

Pightling et al. Frontiers in Microbiology. 2018







together to the exclusion of all others. A paraphyletic topology is one in which isolates of interest (e.g., C and E) group together but not to the exclusion of all others (e.g., D). A polyphyletic topology exists when isolates of interest do not form a group (e.g., A and E).

Pightling et al. Frontiers in Microbiology. 2018






**TABLE 2** Conditions used to determine whether whole-genome sequence analyses support a match between two or more genomes.

	Supports	Neutral	Does not support	
SNP distance	<21	21-100	>100	
Bootstrap support	>0.89	0.80-0.89	< 0.80	
Tree topology	Monophyletic	Paraphyletic	Polyphyletic	

Pightling et al. Frontiers in Microbiology. 2018



- FDA00010369

FDA00010370

PNUSAE003499

FDA00010276 FDA00010258 FDA00010298 FDA00010297

DA00010284

FDA00010310 — FDA00010259 FDA00010296

FDA00010285

FDA00010256 — FDA00010281

- Bootstrap support
- SNP count support
- Topology support
- Epidemiology support


Median 1 SNP Range 0 - 4 SNPs

E. coli Clinical isolates - source


Clinical isolates

Flour isolates

#### L. monocytogenes ingredient supplier – Ice cream producer




- Bootstrap support
- SNP count support
- Topology support
- Epidemiology support



- Clinical isolates
- Environmental isolates

- Bootstrap support
- SNP count support
- Topology support
- Epidemiology not support




inspection.

Clinical isolates

Environmental isolates


- Bootstrap support
- SNP count neutral
- Topology support



- Bootstrap neutral
- SNP count support
- Topology neutral
- Epidemiology Not support

Median 3 SNPs Range 0 - 7 SNPs

- Clinical isolates
- Food isolates
- Environmental isolates



- Bootstrap supports
- SNP count supports
- Topology supports
- Epidemiology does not support








Example	SNP distance	Bootstrap support	Tree topology	Epidemiology, traceback, or compliance findings	Conclusion
Identifying the source of an E. coli outbreak	Supports	Supports	Supports	Supports	Match
Matching food isolates from one firm to environmental isolates from another firm	Supports	Supports	Supports	Supports	Match
Identifying a resident pathogen	Supports	Supports	Supports	Not applicable	Not applicable
Populations of environmental isolates can be very diverse	Neutral	Supports	Supports	Not applicable	Not applicable
Analyzing paraphyletic relationships	Supports	Neutral	Neutral	Does not support	No match
Evidence that isolates arose from the same source by WGS does not necessarily mean that they are linked	Supports	Supports	Supports	Does not support	No match

Pightling et al. Frontiers in Microbiology. 2018





## SnapperDB



- Hierarchical single linkage clustering of pairwise SNP distances
- Performed at 7 snp thresholds.
- If two strains have the same "address" they have 0 SNP differences.
- If two strains have the same address till 50 threshold, they have less than 50 SNP differences.


Dallman et al. Genome anaysis. 2018





#### GenomeTrakr

https://bit.ly/2NNGo37








#### GenomeTrakr

https://bit.ly/2NNGo37








#### GenomeTrakr

#### https://www.ncbi.nlm.nih.gov/pathogens/







# Thank you for your attention!