

## Lecture 3: Random Variable, Part I

Yi, Yung (이웅)

EE210: Probability and Introductory Random Processes  
KAIST EE

June 12, 2021

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

June 12, 2021 1 / 1


June 12, 2021 2 / 1

## Roadmap

## Random Variable: Idea

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

- In reality, many outcomes are **numerical**, e.g., stock price.
- Even if not, very convenient if we map numerical values to random outcomes, e.g., '0' for male and '1' for female.



(b) Two rolls of tetrahedral dice

- Mathematically, a random variable  $X$  is a **function** which maps from  $\Omega$  to  $\mathbb{R}$ .
- Notation.** Random variable  $X$ , numerical value  $x$ .
- Different random variables can be defined on the same sample space.
- For a fixed value  $x$ , we can associate an **event** that a random variable  $X$  has the value  $x$ , i.e.,  $\{\omega \in \Omega \mid X(\omega) = x\}$
- Assume that values  $x$  are discrete<sup>1</sup> such as  $1, 2, 3, \dots$ .  
For notational convenience,  

$$p_X(x) \triangleq \mathbb{P}(X = x) \triangleq \mathbb{P}(\{\omega \in \Omega \mid X(\omega) = x\})$$
- For a discrete random variable  $X$ , we call  $p_X(x)$  **probability mass function** (PMF).

<sup>1</sup>Finite or countably infinite.

- Rolls a dice,  $\Omega = \{1, 2, 3, 4, 5, 6\}$
- Define a random variable  $X = 1$  for even numbers and  $X = 0$  for odd numbers
- Event  $A_1 = \{\omega \in \Omega \mid X(\omega) = 1\} = \{2, 4, 6\} \subset \Omega$ , but simply  $A_1 = \{X = 1\}$
- Event  $A_0 = \{\omega \in \Omega \mid X(\omega) = 0\} = \{1, 3, 5\} \subset \Omega$ , but simply  $A_0 = \{X = 0\}$
- Remember that the random variable  $X$  is a **function** from  $\Omega$  to  $\mathbb{R}$

- (1) Random variable: Idea and formal definition
- (2) **Popular discrete random variables**
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

- Only **binary** values

$$X = \begin{cases} 0, & \text{w.p. } 1 - p, \\ 1, & \text{w.p. } p \end{cases}$$


In other words,  $p_X(0) = 1 - p$  and  $p_X(1) = p$  from our PMF notation.

- Models a trial that results in binary results, e.g., success/failure, head/tail
- Very useful for an **indicator rv** of an event  $A$ . Define a rv  $\mathbf{1}_A$  as:

$$\mathbf{1}_A = \begin{cases} 1, & \text{if } A \text{ occurs,} \\ 0, & \text{otherwise} \end{cases}$$

<sup>1</sup>w.p.: with probability

- integers  $a, b$ , where  $a \leq b$
- Choose a number out of  $\Omega = \{a, a+1, \dots, b\}$  uniformly at random.
- $p_X(i) = \frac{1}{b-a+1}$ ,  $i \in \Omega$




- Models complete **ignorance** (I don't know anything about  $X$ )

L3(2)

June 12, 2021 9 / 1

- Models the number of **successes** in a given number of **independent** trials
- $n$  independent trials, where one trial has the success probability  $p$ .

$$p_X(k) = \binom{n}{k} p^k (1-p)^{n-k}$$




$$\binom{n}{k} = \frac{n!}{k!(n-k)!}, \text{ which we read 'n choose k'.$$

June 12, 2021 10 / 1

- Infinitely many independent Bernoulli trials, where each trial has success probability  $p$
- Random variable: number of trials until the **first success**.

$$p_X(k) = (1-p)^{k-1} p$$

- Models **waiting** times until something happens.



- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) **Summarizing random variables: Expectation and Variance**
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

L3(2)

June 12, 2021 11 / 1

L3(3)

June 12, 2021 12 / 1

- Average

## Definition

$$\mathbb{E}[X] = \sum_x x p_X(x)$$

- $p_X(x)$ : relative frequency of value  $x$  (trials with  $x$ /total trials)

- **Example.** Bernoulli rv with  $p$

$$\mathbb{E}[X] = 1 \times p + 0 \times (1 - p) = p = p_X(1)$$


Not very surprising. Easy to prove using the definition.

- If  $X \geq 0$ ,  $\mathbb{E}[X] \geq 0$ .

- If  $a \leq X \leq b$ ,  $a \leq \mathbb{E}[X] \leq b$ .

- For a constant  $c$ ,  $\mathbb{E}[c] = c$ .

- For a rv  $X$ ,  $Y = g(X)$  is also a r.v.
- $\mathbb{E}[Y] = \mathbb{E}[g(X)] = \sum_x g(x) p_X(x)$
- Compute  $\mathbb{E}[Y]$  for the following:



## Linearity of Expectation

$$\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$$

- Measures how much the **spread** of a PMF is.
- What about  $\mathbb{E}[X - \mu]$ , where  $\mu = \mathbb{E}[X]$ ? Zero
- Then, what about  $\mathbb{E}[(X - \mu)^2]$ ?

## Variance, Standard Deviation

$$\text{var}[X] = \mathbb{E}[(X - \mu)^2]$$

$$\sigma_X = \sqrt{\text{var}[X]}$$

- $\text{var}[X] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$   
 $\text{var}[X] = \mathbb{E}[X^2 - 2\mu X + \mu^2]$   
 $= \mathbb{E}[X^2] - 2\mu\mathbb{E}[X] + \mu^2 = \mathbb{E}[X^2] - \mu^2$
- $Y = X + b, \text{var}[Y] = \text{var}[X]$   
 $\text{var}[Y] = \mathbb{E}[(X + b)^2] - (\mathbb{E}[X + b])^2$
- $Y = aX, \text{var}[Y] = a^2\text{var}[X]$   
 $\text{var}[Y] = \mathbb{E}[a^2X^2] - (a\mathbb{E}[X])^2$

Example: Variance of a Bernoulli rv ( $p$ )

$$\begin{aligned}\mu &= \mathbb{E}[X] = 1 \times p + 0 \times (1 - p) = p \\ \mathbb{E}[X^2] &= 1 \times p + 0 \times (1 - p) = p \\ \text{var}[X] &= \mathbb{E}[X^2] - \mu^2 = p - p^2 \\ &= p(1 - p)\end{aligned}$$

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

- **Joint PMF.** For two random variables  $X, Y$ , consider two events  $\{X = x\}$  and  $\{Y = y\}$ , and  
 $p_{X,Y}(x, y) \triangleq \mathbb{P}(\{X = x\} \cap \{Y = y\})$
- $\sum_x \sum_y p_{X,Y}(x, y) = 1$
- **Marginal PMF.**

$$p_X(x) = \sum_y p_{X,Y}(x, y),$$

$$p_Y(y) = \sum_x p_{X,Y}(x, y)$$

Example.

VIDEO PAUSE

|   |      |      |      |      |
|---|------|------|------|------|
|   |      |      |      |      |
| y |      |      |      |      |
| 4 | 1/20 | 2/20 | 2/20 |      |
| 3 | 2/20 | 4/20 | 1/20 | 2/20 |
| 2 |      | 1/20 | 3/20 | 1/20 |
| 1 |      | 1/20 |      |      |
|   | 1    | 2    | 3    | 4    |

$$p_{X,Y}(1, 3) = 2/20$$

$$p_X(4) = 2/20 + 1/20 = 3/20$$

$$\mathbb{P}(X = Y) = 1/20 + 4/20 + 3/20 = 8/20$$

- Consider a rv  $Z = g(X, Y)$ . (Ex)  $X + Y, X^2 + Y^2$ . Then, PMF of  $Z$  is:

$$p_Z(z) = \mathbb{P}(g(X, Y) = z) = \sum_{(x,y): g(x,y)=z} p_{X,Y}(x, y)$$

- Similarly,

$$\mathbb{E}[Z] = \mathbb{E}[g(X, Y)] = \sum_x \sum_y g(x, y) p_{X,Y}(x, y)$$

- Remember:  $\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$
- Similarly,
 
$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$
 (easy to prove, using the definition.)
- $\mathbb{E}[X_1 + \dots + X_n] = \mathbb{E}[X_1] + \dots + \mathbb{E}[X_n]$
- $\mathbb{E}[2X + 3Y - Z] = 2\mathbb{E}[X] + 3\mathbb{E}[Y] - \mathbb{E}[Z]$

- Example.** Mean of a binomial rv  $Y$  with  $(n, p)$
  - $Y$ : number of successes in  $n$  Bernoulli trials with  $p$
  - $Y = X_1 + \dots + X_n$ , where  $X_i$  is a Bernoulli rv.
  - $\mathbb{E}[Y] = n\mathbb{E}[X_i] = n\mathbb{P}(X_i = 1) = np$
- Message.** When some rv  $X$  is written as a linear combination of other rvs,  $X$  becomes easy to handle.

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) **Conditioning for random variables**
- (6) Independence for random variables

L3(4)

June 12, 2021 21 / 1


L3(5)

June 12, 2021 22 / 1

Remember two probability laws:  $\mathbb{P}(\cdot)$  and  $\mathbb{P}(\cdot|A)$  for an event  $A$ .


- $p_X(x) \triangleq \mathbb{P}(X = x)$
  - $\mathbb{E}[X] = \sum_x x p_X(x)$
  - $\mathbb{E}[g(X)] = \sum_x g(x) p_X(x)$
  - $\text{var}[X] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$
- $p_{X|A}(x) \triangleq \mathbb{P}(X = x|A)$
  - $\mathbb{E}[X|A] \triangleq \sum_x x p_{X|A}(x)$
  - $\mathbb{E}[g(X)|A] \triangleq \sum_x g(x) p_{X|A}(x)$
  - $\text{var}[X|A] \triangleq \mathbb{E}[X^2|A] - (\mathbb{E}[X|A])^2$
  - (Note)**  $p_{X|A}(x)$ ,  $\mathbb{E}[X|A]$ ,  $\mathbb{E}[g(X)|A]$ , and  $\text{var}[X|A]$  are all just notations!

$$A = \{X \geq 2\}$$



$$\mathbb{E}[X] = \frac{1}{4}(1 + 2 + 3 + 4) = 2.5$$

$$\begin{aligned} \text{var}[X] &= \mathbb{E}[X^2] - (\mathbb{E}[X])^2 \\ &= \frac{1}{4}(1 + 2^2 + 3^2 + 4^2) - 2.5^2 \end{aligned}$$



$$\mathbb{E}[X|A] = \frac{1}{3}(2 + 3 + 4) = 3$$

$$\begin{aligned} \text{var}[X|A] &= \mathbb{E}[X^2|A] - (\mathbb{E}[X|A])^2 \\ &= \frac{1}{3}(2^2 + 3^2 + 4^2) - 3^2 = 2/3 \end{aligned}$$

L3(5)

June 12, 2021 23 / 1

L3(5)

June 12, 2021 24 / 1

What do we mean by “conditioning on a rv”? Consider  $A = \{Y = y\}$  for a rv  $Y$ .

- $p_{X|A}(x) \triangleq \mathbb{P}(X = x|A)$
  - $\mathbb{E}[X|A] \triangleq \sum_x x p_{X|A}(x)$
  - $\mathbb{E}[g(X)|A] \triangleq \sum_x g(x) p_{X|A}(x)$
  - $\text{var}[X|A] \triangleq \mathbb{E}[X^2|A] - (\mathbb{E}[X|A])^2$
- $p_{X|Y}(x|y) \triangleq \mathbb{P}(X = x|Y = y)$
  - $\mathbb{E}[X|Y = y] \triangleq \sum_x x p_{X|Y}(x|y)$
  - $\mathbb{E}[g(X)|Y = y] \triangleq \sum_x g(x) p_{X|Y}(x|y)$
  - $\text{var}[X|Y = y] \triangleq \mathbb{E}[X^2|Y = y] - (\mathbb{E}[X|Y = y])^2$

L3(5)

June 12, 2021 25 / 1

VIDEO PAUSE

- Conditional PMF

$$p_{X|Y}(x|y) \triangleq \mathbb{P}(X = x|Y = y) = \frac{p_{X,Y}(x,y)}{p_Y(y)}$$

for  $y$  such that  $p_Y(y) > 0$ .

- $\sum_x p_{X|Y}(x|y) = 1$

- Multiplication rule

$$p_{X,Y}(x,y) = p_Y(y) p_{X|Y}(x|y) \\ = p_X(x) p_{Y|X}(y|x)$$

- $p_{X,Y,Z}(x,y,z) = p_X(x) p_{Y|X}(y|x) p_{Z|X,Y}(z|x,y)$

|   |      |      |      |      |
|---|------|------|------|------|
|   |      |      |      |      |
|   |      |      |      |      |
| 4 | 1/20 | 2/20 | 2/20 |      |
| 3 | 2/20 | 4/20 | 1/20 | 2/20 |
| 2 |      | 1/20 | 3/20 | 1/20 |
| 1 |      | 1/20 |      |      |
|   | 1    | 2    | 3    | 4    |

$$p_{X|Y}(2|2) = \frac{1}{1+3+1}$$

$$p_{X|Y}(3|2) = \frac{3}{1+3+1}$$

$$\mathbb{E}[X|Y = 3] = 1(2/9) + 2(4/9) + 3(1/9) + 4(2/9)$$


L3(5)

June 12, 2021 26 / 1

- Partition of  $\Omega$  into  $A_1, A_2, A_3$
- Known:  $\mathbb{P}(A_i)$  and  $\mathbb{P}(B|A_i)$
- What is  $\mathbb{P}(B)$ ?

## Total Probability Theorem

$$\mathbb{P}(B) = \sum_i \mathbb{P}(A_i) \mathbb{P}(B|A_i)$$




L3(5)

June 12, 2021 27 / 1

- Partition of  $\Omega$  into  $A_1, A_2, A_3$

## Total Probability Theorem

$$p_X(x) = \sum_i \mathbb{P}(A_i) \mathbb{P}(X = x|A_i) = \sum_i \mathbb{P}(A_i) p_{X|A_i}(x)$$



L3(5)

June 12, 2021 28 / 1

## Total Expectation Theorem for $\{A_i\}$

- Partition of  $\Omega$  into  $A_1, A_2, A_3$


### Total Probability Theorem

$$p_X(x) = \sum_i \mathbb{P}(A_i) \mathbb{P}(X = x | A_i) = \sum_i \mathbb{P}(A_i) p_{X|A_i}(x)$$

### Total Expectation Theorem

$$\mathbb{E}[X] = \sum_i \mathbb{P}(A_i) \mathbb{E}[X | A_i]$$

- Weighted average of expectations from  $A_i$ 's perspective



L3(5)

June 12, 2021 29 / 1

## Total Expectation Theorem for $\{Y = y\}$


- Partition of  $\Omega$  into  $A_1, A_2, A_3$

### Total Expectation Theorem

$$\mathbb{E}[X] = \sum_i \mathbb{P}(A_i) \mathbb{E}[X | A_i]$$

### Total Expectation Theorem

$$\mathbb{E}[X] = \sum_y \mathbb{P}(Y = y) \mathbb{E}[X | Y = y] = \sum_y p_Y(y) \mathbb{E}[X | Y = y]$$



L3(5)

June 12, 2021 30 / 1

## Example 1: Total Expectation Theorem

- Question.** What is  $\mathbb{E}(X)$ ?


- Just using the definition of expectation,

$$\begin{aligned} \mathbb{E}[X] &= \frac{1}{9}(0 + 1 + 2) + \frac{2}{9}(6 + 7 + 8) \\ &= \frac{3 + 12 + 14 + 16}{9} = 5 \end{aligned}$$

- Let's use TET, for which consider

$$A_1 = \{X \in \{0, 1, 2\}\}, A_2 = \{X \in \{6, 7, 8\}\}$$

$$\begin{aligned} \mathbb{E}[X] &= \sum_{i=1,2} \mathbb{P}(A_i) \mathbb{E}[X | A_i] \\ &= 1/3 \cdot 1 + 2/3 \cdot 7 = 5 \end{aligned}$$



L3(5)

June 12, 2021 31 / 1

## Example 2: Mean of Geometric rv

- Write softwares over and over, and each time w.p.  $p$  of working correctly (independent from previous programs).

- $X$ : number of trials until the program works correctly.

- (Q)**  $\mathbb{E}(X)$ ?

- $X$  is a geometric rv

- Direct computation is boring.


$$\mathbb{E}[X] = \sum_{k=1}^{\infty} k(1-p)^{k-1}p = p + 2(1-p)p + 3(1-p)^2p + \dots$$

- Total expectation theorem and a notion of **memorylessness** helps a lot.

L3(5)

June 12, 2021 32 / 1





L3(5)

June 12, 2021 33 / 1

L3(5)

June 12, 2021 34 / 1

- Some random variable often does not have **memory**.
- Definition.** A random variable  $X$  is called **memoryless** if, for any  $n, m \geq 0$ ,
 
$$\mathbb{P}(X > n + m | X > m) = \mathbb{P}(X > n)$$
- Meaning.** Conditioned on  $X > m$ ,  $X - m$ 's distribution is the same as the original  $X$ .
 
$$\mathbb{P}(X - m > n | X > m) = \mathbb{P}(X > n)$$

- Theorem.** Any **geometric** random variable is **memoryless**.
- Remind.** Geometric rv  $X$  with parameter  $p$

$$\mathbb{P}(X = k) = (1 - p)^{k-1}p, \quad \mathbb{P}(X > k) = \sum_{i=k+1}^{\infty} (1 - p)^{i-1}p = (1 - p)^k$$

- Proof.**

$$\begin{aligned} \mathbb{P}(X > n + m | X > m) &= \frac{\mathbb{P}(X > n + m \text{ and } X > m)}{\mathbb{P}(X > m)} = \frac{\mathbb{P}(X > n + m)}{\mathbb{P}(X > m)} \\ &= \frac{(1 - p)^{n+m}}{(1 - p)^m} = (1 - p)^n = \mathbb{P}(X > n) \end{aligned}$$

- Meaning.** Conditioned on  $X > m$ ,  $X - m$  is geometric with the same parameter.

L3(5)

June 12, 2021 35 / 1

L3(5)

June 12, 2021 36 / 1

- $A_1 = \{X = 1\}$  (first try is success),  $A_2 = \{X > 1\}$  (first try is failure).
 
$$\begin{aligned} \mathbb{E}[X] &= 1 + \mathbb{E}[X - 1] \\ &= 1 + \mathbb{P}(A_1)\mathbb{E}[X - 1 | X = 1] + \mathbb{P}(A_2)\mathbb{E}[X - 1 | X > 1] \quad (\text{from TET}) \\ &= 1 + (1 - p)\mathbb{E}[X] \quad (\text{from memorylessness}) \end{aligned}$$
- Thus,  $\mathbb{E}[X] = \frac{1}{p}$

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

L3(6)

June 12, 2021 37 / 1

- Two events

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$$

$$\mathbb{P}(A \cap B | C) = \mathbb{P}(A | C) \cdot \mathbb{P}(B | C)$$

- A rv and an event

$$\mathbb{P}(\{X = x\} \cap B) = \mathbb{P}(X = x) \cdot \mathbb{P}(B), \quad \text{for all } x$$

$$\mathbb{P}(\{X = x\} \cap B | C) = \mathbb{P}(X = x | C) \cdot \mathbb{P}(B | C), \quad \text{for all } x$$

- Two rvs

$$\mathbb{P}(\{X = x\} \cap \{Y = y\}) = \mathbb{P}(X = x) \cdot \mathbb{P}(Y = y), \quad \text{for all } x, y$$

$$p_{X,Y}(x, y) = p_X(x) \cdot p_Y(y)$$

$$\mathbb{P}(\{X = x\} \cap \{Y = y\} | Z = z) = \mathbb{P}(X = x | Z = z) \cdot \mathbb{P}(Y = y | Z = z), \quad \text{for all } x, y$$

$$p_{X,Y|Z}(x, y) = p_{X|Z}(x) \cdot p_{Y|Z}(y)$$

L3(6)

June 12, 2021 38 / 1

## Example

- $X \perp\!\!\!\perp Y$ ?

$$p_{X,Y}(1, 1) = 0$$

$$p_X(1) = 3/20$$


$$p_Y(1) = 1/20$$

- $X \perp\!\!\!\perp Y | \{X \leq 2 \text{ and } Y \geq 3\}$ ?

VIDEO PAUSE

|               |               |               |
|---------------|---------------|---------------|
| $Y = 4$ (1/3) | 1/9           | 2/9           |
| $Y = 3$ (2/3) | 2/9           | 4/9           |
|               | $X = 1$ (1/3) | $X = 2$ (2/3) |

- Yes.



L3(6)

June 12, 2021 39 / 1

## Expectation and Variance

- Always true.

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

- Generally,  $\mathbb{E}[g(X, Y)] \neq g(\mathbb{E}[X], \mathbb{E}[Y])$

- However, if  $X \perp\!\!\!\perp Y$ ,

$$\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$$

$$\mathbb{E}[g(X)h(Y)] = \mathbb{E}[g(X)]\mathbb{E}[h(Y)]$$

- Proof.

$$\begin{aligned} \mathbb{E}[g(X)h(Y)] &= \sum_x \sum_y g(x)h(y)p_{X,Y}(x, y) \\ &= \sum_x g(x)p_X(x) \sum_y h(y)p_Y(y) \end{aligned}$$

- Always true.

$$\text{var}[aX] = a^2\text{var}[X], \text{var}[X + a] = \text{var}[X]$$

- Generally,  $\text{var}[X + Y] \neq \text{var}[X] + \text{var}[Y]$  (next slide)

- However, if  $X \perp\!\!\!\perp Y$ ,

$$\text{var}[X + Y] = \text{var}[X] + \text{var}[Y]$$

- Practice.

- $X = Y \implies \text{var}[X + Y] = 4\text{var}[X]$
- $X = -Y \implies \text{var}[X + Y] = 0$
- $X \perp\!\!\!\perp Y \implies \text{var}[X - 3Y] = \text{var}[X] + 9\text{var}[Y]$

L3(6)

June 12, 2021 40 / 1

- Why not generally true?

$$\begin{aligned}\text{var}[X + Y] &= \mathbb{E}[(X + Y)^2] - (\mathbb{E}[X + Y])^2 \\ &= \mathbb{E}[X^2 + Y^2 + 2XY] - ((\mathbb{E}[X])^2 + (\mathbb{E}[Y])^2 + 2\mathbb{E}[X]\mathbb{E}[Y]) \\ &= \text{var}[X] + \text{var}[Y] + 2(\mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y])\end{aligned}$$

- $X \perp\!\!\!\perp Y$  is a sufficient condition for  $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$
- Also, a necessary condition? we will see later, when we study **covariance**.

L3(6)

June 12, 2021 41 / 1

- $n$  people throw their hats in a box and then pick one at random

- $X$ : number of people with their own hat

- $\mathbb{E}[X]$ ?  $\text{var}[X]$ ?

- All permutations are equally likely as  $1/n!$ . Thus, this equals to picking one hat at a time.

- Key step 1.** Define a rv  $X_i = 1$  if  $i$  selects its own hat and 0 otherwise.

$$X = \sum_{i=1}^n X_i.$$

- $\{X_i\}, i = 1, 2, \dots, n$ : identically distributed (from symmetry)

L3(6)

June 12, 2021 42 / 1

- $\mathbb{E}[X] = n\mathbb{E}[X_1] = n\mathbb{P}(X_1 = 1) = n \times \frac{1}{n} = 1$ .
- Key step 2.** Are  $X_i$ s are independent? If yes, easy to get  $\text{var}(X)$ .
- Assume  $n = 2$ . Then,  $X_1 = 1 \rightarrow X_2 = 1$ , and  $X_1 = 0 \rightarrow X_2 = 0$ . Thus, **dependent**.

$$\text{var}(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2 = \mathbb{E}\left[\sum_i X_i^2 + \sum_{i,j:i \neq j} X_i X_j\right] - (\mathbb{E}[X])^2$$

$$\mathbb{E}[X_i^2] = \mathbb{E}[X_1^2] = 1 \times \frac{1}{n} + 0 \times \frac{n-1}{n} = \frac{1}{n}$$

$$\mathbb{E}[X_i X_j] = \mathbb{E}[X_1 X_2] = 1 \times \mathbb{P}(X_1 X_2 = 1) = \mathbb{P}(X_1 = 1)\mathbb{P}(X_2 = 1|X_1 = 1), \quad (i \neq j)$$

- $\mathbb{E}[X^2] = n\mathbb{E}[X_1^2] + n(n-1)\mathbb{E}[X_1 X_2] = n\frac{1}{n} + n(n-1)\frac{1}{n(n-1)} = 2$
- $\text{var}(X) = 2 - 1 = 1$

Questions?

L3(6)

June 12, 2021 43 / 1

L3(6)

June 12, 2021 44 / 1

- 1) What is Random Variable? Why is it useful?
- 2) What is PMF (Probability Mass Function)?
- 3) Explain Bernoulli, Binomial, Poisson, Geometric rvs, when they are used and what their PMFs are.
- 4) What are joint and marginal PMFs?
- 5) Describe and explain the total probability/expectation theorem for random variables?
- 6) When is it useful to use total probability/expectation theorem?
- 7) What is conditional independence?