

## Lecture 3: Random Variable, Part I

Yi, Yung (이웅)

EE210: Probability and Introductory Random Processes  
KAIST EE

May 5, 2021

May 5, 2021 1 / 45

### Roadmap

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

May 5, 2021 2 / 45


- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

L3(1)

May 5, 2021 3 / 45

## Random Variable: Idea

- In reality, many outcomes are **numerical**, e.g., stock price.
- Even if not, very convenient if we map numerical values to random outcomes, e.g., '0' for male and '1' for female.



(b) Two rolls of tetrahedral dice

L3(1)

May 5, 2021 4 / 45

- Mathematically, a random variable  $X$  is a **function** which maps from  $\Omega$  to  $\mathbb{R}$ .
- Notation.** Random variable  $X$ , numerical value  $x$ .
- Different random variables can be defined on the same sample space.
- For a fixed value  $x$ , we can associate an **event** that a random variable  $X$  has the value  $x$ , i.e.,  $\{\omega \in \Omega \mid X(\omega) = x\}$
- Assume that values  $x$  are discrete<sup>1</sup> such as  $1, 2, 3, \dots$ .  
For notational convenience,

$$p_X(x) \triangleq \mathbb{P}(X = x) \triangleq \mathbb{P}(\{\omega \in \Omega \mid X(\omega) = x\})$$

- For a discrete random variable  $X$ , we call  $p_X(x)$  **probability mass function** (PMF).

<sup>1</sup>Finite or countably infinite.

## Example

- Rolls a dice,  $\Omega = \{1, 2, 3, 4, 5, 6\}$
- Define a random variable  $X = 1$  for even numbers and  $X = 0$  for odd numbers
- Event  $A_1 = \{\omega \in \Omega \mid X(\omega) = 1\} = \{2, 4, 6\} \subset \Omega$ , but simply  $A_1 = \{X = 1\}$
- Event  $A_0 = \{\omega \in \Omega \mid X(\omega) = 0\} = \{1, 3, 5\} \subset \Omega$ , but simply  $A_0 = \{X = 0\}$
- Remember that the random variable  $X$  is a **function** from  $\Omega$  to  $\mathbb{R}$

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

L3(2)

May 5, 2021 7 / 45

Bernoulli  $X$  with parameter  $p \in [0, 1]$ 

- Only **binary** values

$$X = \begin{cases} 0, & \text{w.p. } 1 - p, \\ 1, & \text{w.p. } p \end{cases}$$

In other words,  $p_X(0) = 1 - p$  and  $p_X(1) = p$  from our PMF notation.


- Models a trial that results in binary results, e.g., success/failure, head/tail
- Very useful for an **indicator rv** of an event  $A$ . Define a rv  $\mathbf{1}_A$  as:

$$\mathbf{1}_A = \begin{cases} 1, & \text{if } A \text{ occurs,} \\ 0, & \text{otherwise} \end{cases}$$

---

<sup>1</sup>w.p.: with probability  
L3(2)

- integers  $a, b$ , where  $a \leq b$
- Choose a number out of  $\Omega = \{a, a + 1, \dots, b\}$  uniformly at random.
- $p_X(i) = \frac{1}{b-a+1}$ ,  $i \in \Omega$




- Models complete **ignorance** (I don't know anything about  $X$ )

L3(2)

May 5, 2021 9 / 45

- Models the number of **successes** in a given number of **independent** trials
- $n$  independent trials, where one trial has the success probability  $p$ .

$$p_X(k) = \binom{n}{k} p^k (1-p)^{n-k}$$



<sup>1</sup> $\binom{n}{k} = \frac{n!}{k!(n-k)!}$ , which we read ' $n$  choose  $k$ '.


L3(2)

May 5, 2021 10 / 45

- Infinitely many independent Bernoulli trials, where each trial has success probability  $p$
- Random variable: number of trials until the **first success**.

$$p_X(k) = (1 - p)^{k-1}p$$

- Models **waiting** times until something happens.



- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) **Summarizing random variables: Expectation and Variance**
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

- Average

## Definition

$$\mathbb{E}[X] = \sum_x x p_X(x)$$

- $p_X(x)$ : relative frequency of value  $x$  (trials with  $x$ /total trials)
- **Example.** Bernoulli rv with  $p$

$$\mathbb{E}[X] = 1 \times p + 0 \times (1 - p) = p = p_X(1)$$

L3(3)

May 5, 2021

13 / 45

## Properties of Expectation

Not very surprising. Easy to prove using the definition.


- If  $X \geq 0$ ,  $\mathbb{E}[X] \geq 0$ .
- If  $a \leq X \leq b$ ,  $a \leq \mathbb{E}[X] \leq b$ .
- For a constant  $c$ ,  $\mathbb{E}[c] = c$ .

L3(3)

May 5, 2021

14 / 45

- For a rv  $X$ ,  $Y = g(X)$  is also a r.v.
- $\mathbb{E}[Y] = \mathbb{E}[g(X)] = \sum_x g(x)p_X(x)$
- Compute  $\mathbb{E}[Y]$  for the following:



$$4 \times (0.4 + 0.3) + 3 \times (0.1 + 0.2) = 2.8 + 0.9 = 3.7$$

## Linearity of Expectation

$$\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$$

L3(3)

May 5, 2021

15 / 45

# Variance

- Measures how much the **spread** of a PMF is.
- What about  $\mathbb{E}[X - \mu]$ , where  $\mu = \mathbb{E}[X]$ ? Zero
- Then, what about  $\mathbb{E}[(X - \mu)^2]$ ?

## Variance, Standard Deviation

$$\text{var}[X] = \mathbb{E}[(X - \mu)^2]$$

$$\sigma_X = \sqrt{\text{var}[X]}$$

L3(3)

May 5, 2021

16 / 45



- $\text{var}[X] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$

$$\begin{aligned}\text{var}[X] &= \mathbb{E}[X^2 - 2\mu X + \mu^2] \\ &= \mathbb{E}[X^2] - 2\mu\mathbb{E}[X] + \mu^2 = \mathbb{E}[X^2] - \mu^2\end{aligned}$$

- $Y = X + b, \text{var}[Y] = \text{var}[X]$

$$\text{var}[Y] = \mathbb{E}[(X + b)^2] - (\mathbb{E}[X + b])^2$$

- $Y = aX, \text{var}[Y] = a^2\text{var}[X]$

$$\text{var}[Y] = \mathbb{E}[a^2X^2] - (a\mathbb{E}[X])^2$$

Example: Variance of a Bernoulli rv ( $p$ )

$$\mu = \mathbb{E}[X] = 1 \times p + 0 \times (1 - p) = p$$

$$\mathbb{E}[X^2] = 1 \times p + 0 \times (1 - p) = p$$

$$\begin{aligned}\text{var}[X] &= \mathbb{E}[X^2] - \mu^2 = p - p^2 \\ &= p(1 - p)\end{aligned}$$

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

- Joint PMF.** For two random variables  $X, Y$ , consider two events  $\{X = x\}$  and  $\{Y = y\}$ , and

$$p_{X,Y}(x,y) \triangleq \mathbb{P}(\{X = x\} \cap \{Y = y\})$$

- $\sum_x \sum_y p_{X,Y}(x,y) = 1$


- Marginal PMF.**

$$p_X(x) = \sum_y p_{X,Y}(x,y),$$

$$p_Y(y) = \sum_x p_{X,Y}(x,y)$$

Example.

VIDEO PAUSE



$$p_{X,Y}(1,3) = 2/20$$

$$p_X(4) = 2/20 + 1/20 = 3/20$$

$$\mathbb{P}(X = Y) = 1/20 + 4/20 + 3/20 = 8/20$$

L3(4)

May 5, 2021 19 / 45

- Consider a rv  $Z = g(X, Y)$ . (Ex)  $X + Y, X^2 + Y^2$ . Then, PMF of  $Z$  is:

$$p_Z(z) = \mathbb{P}(g(X, Y) = z) = \sum_{(x,y): g(x,y)=z} p_{X,Y}(x,y)$$

- Similarly,

$$\mathbb{E}[Z] = \mathbb{E}[g(X, Y)] = \sum_x \sum_y g(x,y) p_{X,Y}(x,y)$$

L3(4)

May 5, 2021 20 / 45

- Remember:  $\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$

- Similarly,

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

(easy to prove, using the definition.)

- $\mathbb{E}[X_1 + \dots + X_n] = \mathbb{E}[X_1] + \dots + \mathbb{E}[X_n]$
- $\mathbb{E}[2X + 3Y - Z] = 2\mathbb{E}[X] + 3\mathbb{E}[Y] - \mathbb{E}[Z]$

- Example.** Mean of a binomial rv  $Y$  with  $(n, p)$

- $Y$ : number of successes in  $n$  Bernoulli trials with  $p$

- $Y = X_1 + \dots + X_n$ , where  $X_i$  is a Bernoulli rv.

- $\mathbb{E}[Y] = n\mathbb{E}[X_i] = n\mathbb{P}(X_i = 1) = np$

**Message.** When some rv  $X$  is written as a linear combination of other rvs,  $X$  becomes easy to handle.

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) **Conditioning for random variables**
- (6) Independence for random variables

Remember two probability laws:  $\mathbb{P}(\cdot)$  and  $\mathbb{P}(\cdot|A)$  for an event  $A$ .


- $p_X(x) \triangleq \mathbb{P}(X = x)$
  - $\mathbb{E}[X] = \sum_x x p_X(x)$
  - $\mathbb{E}[g(X)] = \sum_x g(x) p_X(x)$
  - $\text{var}[X] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$
- $p_{X|A}(x) \triangleq \mathbb{P}(X = x|A)$
  - $\mathbb{E}[X|A] \triangleq \sum_x x p_{X|A}(x)$
  - $\mathbb{E}[g(X)|A] \triangleq \sum_x g(x) p_{X|A}(x)$
  - $\text{var}[X|A] \triangleq \mathbb{E}[X^2|A] - (\mathbb{E}[X|A])^2$
  - (Note)  $p_{X|A}(x)$ ,  $\mathbb{E}[X|A]$ ,  $\mathbb{E}[g(X)|A]$ , and  $\text{var}[X|A]$  are all just notations!

L3(5)

May 5, 2021 23 / 45


## Example: Conditional PMF

$$A = \{X \geq 2\}$$



$$\mathbb{E}[X] = \frac{1}{4}(1 + 2 + 3 + 4) = 2.5$$

$$\begin{aligned} \text{var}[X] &= \mathbb{E}[X^2] - (\mathbb{E}[X])^2 \\ &= \frac{1}{4}(1 + 2^2 + 3^2 + 4^2) - 2.5^2 \end{aligned}$$



$$\mathbb{E}[X|A] = \frac{1}{3}(2 + 3 + 4) = 3$$

$$\begin{aligned} \text{var}[X|A] &= \mathbb{E}[X^2|A] - (\mathbb{E}[X|A])^2 \\ &= \frac{1}{3}(2^2 + 3^2 + 4^2) - 3^2 = 2/3 \end{aligned}$$

L3(5)

May 5, 2021 24 / 45

What do we mean by “conditioning on a rv”? Consider  $A = \{Y = y\}$  for a rv  $Y$ .

- $p_{X|A}(x) \triangleq \mathbb{P}(X = x|A)$
  - $\mathbb{E}[X|A] \triangleq \sum_x x p_{X|A}(x)$
  - $\mathbb{E}[g(X)|A] \triangleq \sum_x g(x) p_{X|A}(x)$
  - $\text{var}[X|A] \triangleq \mathbb{E}[X^2|A] - (\mathbb{E}[X|A])^2$
- $p_{X|Y}(x|y) \triangleq \mathbb{P}(X = x|Y = y)$
  - $\mathbb{E}[X|Y = y] \triangleq \sum_x x p_{X|Y}(x|y)$
  - $\mathbb{E}[g(X)|Y = y] \triangleq \sum_x g(x) p_{X|Y}(x|y)$
  - $\text{var}[X|Y = y] \triangleq \mathbb{E}[X^2|Y = y] - (\mathbb{E}[X|Y = y])^2$

L3(5)

May 5, 2021 25 / 45

## Conditional PMF

### Conditional PMF

$$p_{X|Y}(x|y) \triangleq \mathbb{P}(X = x|Y = y) = \frac{p_{X,Y}(x,y)}{p_Y(y)}$$

for  $y$  such that  $p_Y(y) > 0$ .


- $\sum_x p_{X|Y}(x|y) = 1$

### Multiplication rule

$$\begin{aligned} p_{X,Y}(x,y) &= p_Y(y) p_{X|Y}(x|y) \\ &= p_X(x) p_{Y|X}(y|x) \end{aligned}$$

- $p_{X,Y,Z}(x,y,z) = p_X(x) p_{Y|X}(y|x) p_{Z|X,Y}(z|x,y)$

VIDEO PAUSE



$$p_{X|Y}(2|2) = \frac{1}{1+3+1}$$

$$p_{X|Y}(3|2) = \frac{3}{1+3+1}$$

$$\mathbb{E}[X|Y = 3] = 1(2/9) + 2(4/9) + 3(1/9) + 4(2/9)$$



L3(5)

May 5, 2021 26 / 45

- Partition of  $\Omega$  into  $A_1, A_2, A_3$
- Known:  $\mathbb{P}(A_i)$  and  $\mathbb{P}(B|A_i)$
- What is  $\mathbb{P}(B)$ ?

## Total Probability Theorem

$$\mathbb{P}(B) = \sum_i \mathbb{P}(A_i) \mathbb{P}(B|A_i)$$



L3(5)

May 5, 2021



27 / 45

Total Probability Theorem:  $B = \{X = x\}$ 

- Partition of  $\Omega$  into  $A_1, A_2, A_3$

## Total Probability Theorem

$$p_X(x) = \sum_i \mathbb{P}(A_i) \mathbb{P}(X = x | A_i) = \sum_i \mathbb{P}(A_i) p_{X|A_i}(x)$$



L3(5)

May 5, 2021

28 / 45

## Total Expectation Theorem for $\{A_i\}$

- Partition of  $\Omega$  into  $A_1, A_2, A_3$


### Total Probability Theorem

$$p_X(x) = \sum_i \mathbb{P}(A_i) \mathbb{P}(X = x | A_i) = \sum_i \mathbb{P}(A_i) p_{X|A_i}(x)$$

### Total Expectation Theorem

$$\mathbb{E}[X] = \sum_i \mathbb{P}(A_i) \mathbb{E}[X | A_i]$$

- Weighted average of expectations from  $A_i$ 's perspective



L3(5)

May 5, 2021 29 / 45

## Total Expectation Theorem for $\{Y = y\}$


- Partition of  $\Omega$  into  $A_1, A_2, A_3$

### Total Expectation Theorem

$$\mathbb{E}[X] = \sum_i \mathbb{P}(A_i) \mathbb{E}[X | A_i]$$

### Total Expectation Theorem

$$\mathbb{E}[X] = \sum_y \mathbb{P}(Y = y) \mathbb{E}[X | Y = y] = \sum_y p_Y(y) \mathbb{E}[X | Y = y]$$



L3(5)

May 5, 2021 30 / 45

- **Question.** What is  $\mathbb{E}(X)$ ?


(1) Just using the definition of expectation,

$$\begin{aligned}\mathbb{E}[X] &= \frac{1}{9}(0 + 1 + 2) + \frac{2}{9}(6 + 7 + 8) \\ &= \frac{3 + 12 + 14 + 16}{9} = 5\end{aligned}$$

(2) Let's use TET, for which consider

$$A_1 = \{X \in \{0, 1, 2\}\}, A_2 = \{X \in \{6, 7, 8\}\}$$

$$\begin{aligned}\mathbb{E}[X] &= \sum_{i=1,2} \mathbb{P}(A_i) \mathbb{E}[X|A_i] \\ &= 1/3 \cdot 1 + 2/3 \cdot 7 = 5\end{aligned}$$



L3(5)

May 5, 2021 31 / 45

- Write softwares over and over, and each time w.p.  $p$  of working correctly (independent from previous programs).
- $X$ : number of trials until the program works correctly.
- **(Q)**  $\mathbb{E}(X)$ ?
- $X$  is a geometric rv
- Direct computation is boring.


$$\mathbb{E}[X] = \sum_{k=1}^{\infty} k(1-p)^{k-1}p = p + 2(1-p)p + 3(1-p)^2p + \dots$$

- Total expectation theorem and a notion of **memorylessness** helps a lot.

L3(5)

May 5, 2021 32 / 45





L3(5)

May 5, 2021 33 / 45

- Some random variable often does not have **memory**.
- **Definition.** A random variable  $X$  is called **memoryless** if, for any  $n, m \geq 0$ ,  

$$\mathbb{P}(X > n + m | X > m) = \mathbb{P}(X > n)$$
- **Meaning.** Conditioned on  $X > m$ ,  $X - m$ 's distribution is the same as the original  $X$ .  

$$\mathbb{P}(X - m > n | X > m) = \mathbb{P}(X > n)$$

L3(5)

May 5, 2021 34 / 45

- **Theorem.** Any **geometric** random variable is **memoryless**.
- **Remind.** Geometric rv  $X$  with parameter  $p$

$$\mathbb{P}(X = k) = (1 - p)^{k-1}p, \quad \mathbb{P}(X > k) = \sum_{i=k+1}^{\infty} (1 - p)^{i-1}p = (1 - p)^k$$

- **Proof.**

$$\begin{aligned} \mathbb{P}(X > n + m | X > m) &= \frac{\mathbb{P}(X > n + m \text{ and } X > m)}{\mathbb{P}(X > m)} = \frac{\mathbb{P}(X > n + m)}{\mathbb{P}(X > m)} \\ &= \frac{(1 - p)^{n+m}}{(1 - p)^m} = (1 - p)^n = \mathbb{P}(X > n) \end{aligned}$$

- **Meaning.** Conditioned on  $X > m$ ,  $X - m$  is geometric with the same parameter.

L3(5)

May 5, 2021 35 / 45

- $A_1 = \{X = 1\}$  (first try is success),  $A_2 = \{X > 1\}$  (first try is failure).

$$\begin{aligned} \mathbb{E}[X] &= 1 + \mathbb{E}[X - 1] \\ &= 1 + \mathbb{P}(A_1)\mathbb{E}[X - 1 | X = 1] + \mathbb{P}(A_2)\mathbb{E}[X - 1 | X > 1] && \text{(from TET)} \\ &= 1 + (1 - p)\mathbb{E}[X] && \text{(from memorylessness)} \end{aligned}$$

- Thus,  $\mathbb{E}[X] = \frac{1}{p}$

L3(5)

May 5, 2021 36 / 45

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

L3(6)

May 5, 2021 37 / 45

## Independence, Conditional Independence

- Two events

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$$

$$\mathbb{P}(A \cap B | C) = \mathbb{P}(A | C) \cdot \mathbb{P}(B | C)$$

- A rv and an event

$$\mathbb{P}(\{X = x\} \cap B) = \mathbb{P}(X = x) \cdot \mathbb{P}(B), \quad \text{for all } x$$

$$\mathbb{P}(\{X = x\} \cap B | C) = \mathbb{P}(X = x | C) \cdot \mathbb{P}(B | C), \quad \text{for all } x$$

- Two rvs

$$\mathbb{P}(\{X = x\} \cap \{Y = y\}) = \mathbb{P}(X = x) \cdot \mathbb{P}(Y = y), \quad \text{for all } x, y$$

$$p_{X,Y}(x, y) = p_X(x) \cdot p_Y(y)$$

$$\mathbb{P}(\{X = x\} \cap \{Y = y\} | Z = z) = \mathbb{P}(X = x | Z = z) \cdot \mathbb{P}(Y = y | Z = z), \quad \text{for all } x, y$$

$$p_{X,Y|Z}(x, y) = p_{X|Z}(x) \cdot p_{Y|Z}(y)$$

L3(6)

May 5, 2021 38 / 45

- $X \perp\!\!\!\perp Y$ ?

$$p_{X,Y}(1,1) = 0$$

$$p_X(1) = 3/20$$


$$p_Y(1) = 1/20$$

- $X \perp\!\!\!\perp Y | \{X \leq 2 \text{ and } Y \geq 3\}$ ?

VIDEO PAUSE

$Y = 4$ (1/3)	1/9	2/9
$Y = 3$ (2/3)	2/9	4/9
	$X = 1$ (1/3)	$X = 2$ (2/3)

- Yes.



## Expectation and Variance

- Always true.

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

- Generally,  $\mathbb{E}[g(X, Y)] \neq g(\mathbb{E}[X], \mathbb{E}[Y])$

- However, if  $X \perp\!\!\!\perp Y$ ,

$$\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$$

$$\mathbb{E}[g(X)h(Y)] = \mathbb{E}[g(X)]\mathbb{E}[h(Y)]$$

- **Proof.**

$$\begin{aligned} \mathbb{E}[g(X)h(Y)] &= \sum_x \sum_y g(x)h(y)p_{X,Y}(x,y) \\ &= \sum_x g(x)p_X(x) \sum_y h(y)p_Y(y) \end{aligned}$$

- Always true.

$$\text{var}[aX] = a^2\text{var}[X], \text{var}[X + a] = \text{var}[X]$$

- Generally,  $\text{var}[X + Y] \neq \text{var}[X] + \text{var}[Y]$  (next slide)

- However, if  $X \perp\!\!\!\perp Y$ ,

$$\text{var}[X + Y] = \text{var}[X] + \text{var}[Y]$$

- **Practice.**

- $X = Y \implies \text{var}[X + Y] = 4\text{var}[X]$
- $X = -Y \implies \text{var}[X + Y] = 0$
- $X \perp\!\!\!\perp Y \implies \text{var}[X - 3Y] = \text{var}[X] + 9\text{var}[Y]$

$$\text{var}[X + Y] \neq \text{var}[X] + \text{var}[Y]$$

- Why not generally true?

$$\begin{aligned}\text{var}[X + Y] &= \mathbb{E}[(X + Y)^2] - (\mathbb{E}[X + Y])^2 \\ &= \mathbb{E}[X^2 + Y^2 + 2XY] - ((\mathbb{E}[X])^2 + (\mathbb{E}[Y])^2 + 2\mathbb{E}[X]\mathbb{E}[Y]) \\ &= \text{var}[X] + \text{var}[Y] + 2(\mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y])\end{aligned}$$

- $X \perp\!\!\!\perp Y$  is a sufficient condition for  $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$
- Also, a necessary condition? we will see later, when we study **covariance**.

## Example: The hat problem (1)

- $n$  people throw their hats in a box and then pick one at random
- $X$ : number of people with their own hat
- $\mathbb{E}[X]$ ?  $\text{var}[X]$ ?
- All permutations are equally likely as  $1/n!$ . Thus, this equals to picking one hat at a time.
- Key step 1.** Define a rv  $X_i = 1$  if  $i$  selects its own hat and 0 otherwise.

$$X = \sum_{i=1}^n X_i.$$

- $\{X_i\}, i = 1, 2, \dots, n$ : identically distributed (from symmetry)

- $\mathbb{E}[X] = n\mathbb{E}[X_1] = n\mathbb{P}(X_1 = 1) = n \times \frac{1}{n} = 1$ .
- **Key step 2.** Are  $X_i$ s independent? If yes, easy to get  $\text{var}(X)$ .
- Assume  $n = 2$ . Then,  $X_1 = 1 \rightarrow X_2 = 1$ , and  $X_1 = 0 \rightarrow X_2 = 0$ . Thus, **dependent**.

$$\text{var}(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2 = \mathbb{E}\left[\sum_i X_i^2 + \sum_{i,j:i \neq j} X_i X_j\right] - (\mathbb{E}[X])^2$$

$$\mathbb{E}[X_i^2] = \mathbb{E}[X_1^2] = 1 \times \frac{1}{n} + 0 \times \frac{n-1}{n} = \frac{1}{n}$$

$$\mathbb{E}[X_i X_j] = \mathbb{E}[X_1 X_2] = 1 \times \mathbb{P}(X_1 X_2 = 1) = \mathbb{P}(X_1 = 1)\mathbb{P}(X_2 = 1|X_1 = 1), \quad (i \neq j)$$

- $\mathbb{E}[X^2] = n\mathbb{E}[X_1^2] + n(n-1)\mathbb{E}[X_1 X_2] = n\frac{1}{n} + n(n-1)\frac{1}{n(n-1)} = 2$
- $\text{var}(X) = 2 - 1 = 1$

## Questions?

- 1) What is Random Variable? Why is it useful?
- 2) What is PMF (Probability Mass Function)?
- 3) Explain Bernoulli, Binomial, Poisson, Geometric rvs, when they are used and what their PMFs are.
- 4) What are joint and marginal PMFs?
- 5) Describe and explain the total probability/expectation theorem for random variables?
- 6) When is it useful to use total probability/expectation theorem?
- 7) What is conditional independence?