

Lecture 3: Random Variable, Part I

Yi, Yung (이웅)


EE210: Probability and Introductory Random Processes
KAIST EE

April 19, 2021

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables


- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

- In reality, many outcomes are , e.g., stock price.
- Even if not, very convenient if we map numerical values to random outcomes, e.g., '0' for male and '1' for female.



(b) Two rolls of tetrahedral dice

- In reality, many outcomes are **numerical**, e.g., stock price.
- Even if not, very convenient if we map numerical values to random outcomes, e.g., '0' for male and '1' for female.



(b) Two rolls of tetrahedral dice

¹Finite or countably infinite.

- Mathematically, a random variable X is a which maps from Ω to \mathbb{R} .

¹Finite or countably infinite.

- Mathematically, a random variable X is a function which maps from Ω to \mathbb{R} .

¹Finite or countably infinite.

- Mathematically, a random variable X is a function which maps from Ω to \mathbb{R} .
- Notation.** Random variable X , numerical value x .

¹Finite or countably infinite.

- Mathematically, a random variable X is a function which maps from Ω to \mathbb{R} .
- Notation.** Random variable X , numerical value x .
- Different random variables can be defined on the same sample space.

¹Finite or countably infinite.

- Mathematically, a random variable X is a function which maps from Ω to \mathbb{R} .
- Notation.** Random variable X , numerical value x .
- Different random variables can be defined on the same sample space.
- For a fixed value x , we can associate an that a random variable X has the value x , i.e.,

¹Finite or countably infinite.

- Mathematically, a random variable X is a **function** which maps from Ω to \mathbb{R} .
- Notation.** Random variable X , numerical value x .
- Different random variables can be defined on the same sample space.
- For a fixed value x , we can associate an **event** that a random variable X has the value x , i.e.,

¹Finite or countably infinite.

- Mathematically, a random variable X is a **function** which maps from Ω to \mathbb{R} .
- Notation.** Random variable X , numerical value x .
- Different random variables can be defined on the same sample space.
- For a fixed value x , we can associate an **event** that a random variable X has the value x , i.e., $\{\omega \in \Omega \mid X(\omega) = x\}$
- Assume that values x are discrete¹ such as $1, 2, 3, \dots$.
For notational convenience,

$$p_X(x) \triangleq \mathbb{P}(X = x) \triangleq \mathbb{P}(\{\omega \in \Omega \mid X(\omega) = x\})$$

¹Finite or countably infinite.

- Mathematically, a random variable X is a **function** which maps from Ω to \mathbb{R} .
- Notation.** Random variable X , numerical value x .
- Different random variables can be defined on the same sample space.
- For a fixed value x , we can associate an **event** that a random variable X has the value x , i.e., $\{\omega \in \Omega \mid X(\omega) = x\}$
- Assume that values x are discrete¹ such as $1, 2, 3, \dots$.
For notational convenience,

$$p_X(x) \triangleq \mathbb{P}(X = x) \triangleq \mathbb{P}(\{\omega \in \Omega \mid X(\omega) = x\})$$
- For a discrete random variable X , we call $p_X(x)$ (PMF).

¹Finite or countably infinite.

- Mathematically, a random variable X is a **function** which maps from Ω to \mathbb{R} .
- Notation.** Random variable X , numerical value x .
- Different random variables can be defined on the same sample space.
- For a fixed value x , we can associate an **event** that a random variable X has the value x , i.e., $\{\omega \in \Omega \mid X(\omega) = x\}$
- Assume that values x are discrete¹ such as $1, 2, 3, \dots$.
For notational convenience,
$$p_X(x) \triangleq \mathbb{P}(X = x) \triangleq \mathbb{P}(\{\omega \in \Omega \mid X(\omega) = x\})$$
- For a discrete random variable X , we call $p_X(x)$ **probability mass function** (PMF).

¹Finite or countably infinite.

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

- Only **binary** values

¹w.p.: with probability

- Only **binary** values

$$X = \begin{cases} 0, & \text{w.p. } 1 - p, \\ 1, & \text{w.p. } p \end{cases}$$

In other words, $p_X(0) = 1 - p$ and $p_X(1) = p$ from our PMF notation.

¹w.p.: with probability

- Only **binary** values

$$X = \begin{cases} 0, & \text{w.p. } 1 - p, \\ 1, & \text{w.p. } p \end{cases}$$

In other words, $p_X(0) = 1 - p$ and $p_X(1) = p$ from our PMF notation.

- Models a trial that results in binary results, e.g., success/failure, head/tail

¹w.p.: with probability

- Only **binary** values

$$X = \begin{cases} 0, & \text{w.p. } 1 - p, \\ 1, & \text{w.p. } p \end{cases}$$

In other words, $p_X(0) = 1 - p$ and $p_X(1) = p$ from our PMF notation.

- Models a trial that results in binary results, e.g., success/failure, head/tail
- Very useful for an of an event A .

¹w.p.: with probability

- Only **binary** values

$$X = \begin{cases} 0, & \text{w.p. } 1 - p, \\ 1, & \text{w.p. } p \end{cases}$$

In other words, $p_X(0) = 1 - p$ and $p_X(1) = p$ from our PMF notation.

- Models a trial that results in binary results, e.g., success/failure, head/tail
- Very useful for an **indicator rv** of an event A . Define a rv $\mathbf{1}_A$ as:

$$\mathbf{1}_A = \begin{cases} 1, & \text{if } A \text{ occurs,} \\ 0, & \text{otherwise} \end{cases}$$


¹w.p.: with probability

- integers a, b , where $a \leq b$


- integers a, b , where $a \leq b$
- Choose a number out of $\Omega = \{a, a + 1, \dots, b\}$ uniformly at random.

- integers a, b , where $a \leq b$
- Choose a number out of $\Omega = \{a, a + 1, \dots, b\}$ uniformly at random.
- $p_X(i) =$

- integers a, b , where $a \leq b$
- Choose a number out of $\Omega = \{a, a+1, \dots, b\}$ uniformly at random.
- $p_X(i) = \frac{1}{b-a+1}$, $i \in \Omega$



- integers a, b , where $a \leq b$
- Choose a number out of $\Omega = \{a, a + 1, \dots, b\}$ uniformly at random.
- $p_X(i) = \frac{1}{b-a+1}$, $i \in \Omega$



- Models complete **ignorance** (I don't know anything about X)

$$^1 \binom{n}{k} = \frac{n!}{k!(n-k)!}, \text{ which we read 'n choose k'.$$

L3(2)

- Models the number of **successes** in a given number of **independent** trials

$$^1 \binom{n}{k} = \frac{n!}{k!(n-k)!}, \text{ which we read 'n choose k'.$$

L3(2)

- Models the number of **successes** in a given number of **independent** trials
- n independent trials, where one trial has the success probability p .


$$p_X(k) =$$

$$^1 \binom{n}{k} = \frac{n!}{k!(n-k)!}, \text{ which we read 'n choose k'.$$

L3(2)

- Models the number of **successes** in a given number of **independent** trials
- n independent trials, where one trial has the success probability p .

$$p_X(k) =$$




$$p_X(k) = \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k}, \text{ which we read 'n choose k'.$$

L3(2)

- Models the number of **successes** in a given number of **independent** trials
- n independent trials, where one trial has the success probability p .

$$p_X(k) = \binom{n}{k} p^k (1-p)^{n-k}$$



$\binom{n}{k} = \frac{n!}{k!(n-k)!}$, which we read ' n choose k '.
L3(2)


- Infinitely many independent Bernoulli trials, where each trial has success probability p

- Infinitely many independent Bernoulli trials, where each trial has success probability p
- Random variable: number of trials until the **first success**.

$$p_X(k) =$$

- Infinitely many independent Bernoulli trials, where each trial has success probability p
- Random variable: number of trials until the **first success**.


$$p_X(k) = (1 - p)^{k-1} p$$



- Infinitely many independent Bernoulli trials, where each trial has success probability p
- Random variable: number of trials until the **first success**.

$$p_X(k) = (1 - p)^{k-1} p$$

- Models **waiting** times until something happens.



- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

- Average

Definition

$$\mathbb{E}[X] = \sum_x x p_X(x)$$

- $p_X(x)$: relative frequency of value x (trials with x /total trials)

- Average

Definition

$$\mathbb{E}[X] = \sum_x x p_X(x)$$

- $p_X(x)$: relative frequency of value x (trials with x /total trials)
- **Example.** Bernoulli rv with p

$$\mathbb{E}[X] = 1 \times p + 0 \times (1 - p) = p = p_X(1)$$


Not very surprising. Easy to prove using the definition.

- If $X \geq 0$, $\mathbb{E}[X] \geq 0$.
- If $a \leq X \leq b$, $a \leq \mathbb{E}[X] \leq b$.
- For a constant c , $\mathbb{E}[c] = c$.


- For a rv X , $Y = g(X)$ is also a r.v.

- For a rv X , $Y = g(X)$ is also a r.v.
- $\mathbb{E}[Y] = \mathbb{E}[g(X)] = \sum_x g(x)p_X(x)$

- For a rv X , $Y = g(X)$ is also a r.v.
- $\mathbb{E}[Y] = \mathbb{E}[g(X)] = \sum_x g(x)p_X(x)$
- Compute $\mathbb{E}[Y]$ for the following:




- For a rv X , $Y = g(X)$ is also a r.v.
- $\mathbb{E}[Y] = \mathbb{E}[g(X)] = \sum_x g(x)p_X(x)$
- Compute $\mathbb{E}[Y]$ for the following:



$$\begin{aligned} 4 \times (0.4 + 0.3) + 3 \times (0.1 + 0.2) \\ = 2.8 + 0.9 = 3.7 \end{aligned}$$

- For a rv X , $Y = g(X)$ is also a r.v.
- $\mathbb{E}[Y] = \mathbb{E}[g(X)] = \sum_x g(x)p_X(x)$
- Compute $\mathbb{E}[Y]$ for the following:



$$\begin{aligned} 4 \times (0.4 + 0.3) + 3 \times (0.1 + 0.2) \\ = 2.8 + 0.9 = 3.7 \end{aligned}$$

Linearity of Expectation

$$\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$$

- Measures how much the **spread** of a PMF is.

- Measures how much the spread of a PMF is.
- What about $\mathbb{E}[X - \mu]$, where $\mu = \mathbb{E}[X]$? Zero

- Measures how much the spread of a PMF is.
- What about $\mathbb{E}[X - \mu]$, where $\mu = \mathbb{E}[X]$? Zero
- Then, what about $\mathbb{E}[(X - \mu)^2]$?

- Measures how much the **spread** of a PMF is.
- What about $\mathbb{E}[X - \mu]$, where $\mu = \mathbb{E}[X]$? Zero
- Then, what about $\mathbb{E}[(X - \mu)^2]$?

Variance, Standard Deviation

$$\text{var}[X] = \mathbb{E}[(X - \mu)^2]$$

$$\sigma_X = \sqrt{\text{var}[X]}$$

- $\text{var}[X] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$
- $Y = X + b, \text{var}[Y] = \text{var}[X]$
- $Y = aX, \text{var}[Y] = a^2\text{var}[X]$

- $\text{var}[X] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$

$$\begin{aligned}\text{var}[X] &= \mathbb{E}[X^2 - 2\mu X + \mu^2] \\ &= \mathbb{E}[X^2] - 2\mu\mathbb{E}[X] + \mu^2 = \mathbb{E}[X^2] - \mu^2\end{aligned}$$

- $Y = X + b, \text{var}[Y] = \text{var}[X]$

- $Y = aX, \text{var}[Y] = a^2\text{var}[X]$

- $\text{var}[X] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$

$$\begin{aligned}\text{var}[X] &= \mathbb{E}[X^2 - 2\mu X + \mu^2] \\ &= \mathbb{E}[X^2] - 2\mu\mathbb{E}[X] + \mu^2 = \mathbb{E}[X^2] - \mu^2\end{aligned}$$

- $Y = X + b, \text{var}[Y] = \text{var}[X]$

$$\text{var}[Y] = \mathbb{E}[(X + b)^2] - (\mathbb{E}[X + b])^2$$

- $Y = aX, \text{var}[Y] = a^2\text{var}[X]$

- $\text{var}[X] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$

$$\begin{aligned}\text{var}[X] &= \mathbb{E}[X^2 - 2\mu X + \mu^2] \\ &= \mathbb{E}[X^2] - 2\mu\mathbb{E}[X] + \mu^2 = \mathbb{E}[X^2] - \mu^2\end{aligned}$$

- $Y = X + b, \text{var}[Y] = \text{var}[X]$

$$\text{var}[Y] = \mathbb{E}[(X + b)^2] - (\mathbb{E}[X + b])^2$$

- $Y = aX, \text{var}[Y] = a^2\text{var}[X]$

$$\text{var}[Y] = \mathbb{E}[a^2X^2] - (a\mathbb{E}[X])^2$$

- $\text{var}[X] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$
$$\text{var}[X] = \mathbb{E}[X^2 - 2\mu X + \mu^2]$$
$$= \mathbb{E}[X^2] - 2\mu\mathbb{E}[X] + \mu^2 = \mathbb{E}[X^2] - \mu^2$$
- $Y = X + b, \text{var}[Y] = \text{var}[X]$
$$\text{var}[Y] = \mathbb{E}[(X + b)^2] - (\mathbb{E}[X + b])^2$$
- $Y = aX, \text{var}[Y] = a^2\text{var}[X]$
$$\text{var}[Y] = \mathbb{E}[a^2X^2] - (a\mathbb{E}[X])^2$$

Example: Variance of a Bernoulli rv (p)

- $\text{var}[X] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$
$$\text{var}[X] = \mathbb{E}[X^2 - 2\mu X + \mu^2]$$
$$= \mathbb{E}[X^2] - 2\mu\mathbb{E}[X] + \mu^2 = \mathbb{E}[X^2] - \mu^2$$
- $Y = X + b, \text{var}[Y] = \text{var}[X]$
$$\text{var}[Y] = \mathbb{E}[(X + b)^2] - (\mathbb{E}[X + b])^2$$
- $Y = aX, \text{var}[Y] = a^2\text{var}[X]$
$$\text{var}[Y] = \mathbb{E}[a^2X^2] - (a\mathbb{E}[X])^2$$

Example: Variance of a Bernoulli rv (p)

$$\mu = \mathbb{E}[X] = 1 \times p + 0 \times (1 - p) = p$$

$$\mathbb{E}[X^2] = 1 \times p + 0 \times (1 - p) = p$$

$$\begin{aligned}\text{var}[X] &= \mathbb{E}[X^2] - \mu^2 = p - p^2 \\ &= p(1 - p)\end{aligned}$$

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

- For two random variables X, Y , consider two events $\{X = x\}$ and $\{Y = y\}$, and

$$\mathbb{P}(\{X = x\} \cap \{Y = y\})$$

- **Joint PMF.** For two random variables X, Y , consider two events $\{X = x\}$ and $\{Y = y\}$, and

$$p_{X,Y}(x,y) \triangleq \mathbb{P}(\{X = x\} \cap \{Y = y\})$$

- **Joint PMF.** For two random variables X, Y , consider two events $\{X = x\}$ and $\{Y = y\}$, and

$$p_{X,Y}(x,y) \triangleq \mathbb{P}(\{X = x\} \cap \{Y = y\})$$

- $\sum_x \sum_y p_{X,Y}(x,y) = 1$

- **Joint PMF.** For two random variables X, Y , consider two events $\{X = x\}$ and $\{Y = y\}$, and

$$p_{X,Y}(x,y) \triangleq \mathbb{P}(\{X = x\} \cap \{Y = y\})$$

- $\sum_x \sum_y p_{X,Y}(x,y) = 1$

-

$$p_X(x) = \sum_y p_{X,Y}(x,y),$$

$$p_Y(y) = \sum_x p_{X,Y}(x,y)$$

- Joint PMF.** For two random variables X, Y , consider two events $\{X = x\}$ and $\{Y = y\}$, and

$$p_{X,Y}(x,y) \triangleq \mathbb{P}(\{X = x\} \cap \{Y = y\})$$

- $\sum_x \sum_y p_{X,Y}(x,y) = 1$

- Marginal PMF.**

$$p_X(x) = \sum_y p_{X,Y}(x,y),$$

$$p_Y(y) = \sum_x p_{X,Y}(x,y)$$

Example.

VIDEO PAUSE

y \ x	1	2	3	4
4	1/20	2/20	2/20	0
3	2/20	4/20	1/20	2/20
2	0	1/20	3/20	1/20
1	0	1/20	0	0

$$p_{X,Y}(1,3) =$$

$$p_X(4) =$$

$$\mathbb{P}(X = Y) =$$

- Joint PMF.** For two random variables X, Y , consider two events $\{X = x\}$ and $\{Y = y\}$, and

$$p_{X,Y}(x,y) \triangleq \mathbb{P}(\{X = x\} \cap \{Y = y\})$$

- $\sum_x \sum_y p_{X,Y}(x,y) = 1$

- Marginal PMF.**

$$p_X(x) = \sum_y p_{X,Y}(x,y),$$

$$p_Y(y) = \sum_x p_{X,Y}(x,y)$$

Example.

VIDEO PAUSE

y \ x	1	2	3	4
4	1/20	2/20	2/20	0
3	2/20	4/20	1/20	2/20
2	0	1/20	3/20	1/20
1	0	1/20	0	0

$$p_{X,Y}(1,3) = 2/20$$

$$p_X(4) = 2/20 + 1/20 = 3/20$$

$$\mathbb{P}(X = Y) = 1/20 + 4/20 + 3/20 = 8/20$$

- Consider a rv $Z = g(X, Y)$. (Ex) $X + Y, X^2 + Y^2$. Then, PMF of Z is:

- Similarly,

$$\mathbb{E}[Z] = \mathbb{E}[g(X, Y)] =$$

- Consider a rv $Z = g(X, Y)$. (Ex) $X + Y, X^2 + Y^2$. Then, PMF of Z is:

$$p_Z(z) = \mathbb{P}(g(X, Y) = z) = \sum_{(x,y): g(x,y)=z} p_{X,Y}(x, y)$$

- Similarly,

$$\mathbb{E}[Z] = \mathbb{E}[g(X, Y)] = \sum_x \sum_y g(x, y) p_{X,Y}(x, y)$$

- Remember: $\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$

- Remember: $\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$

- Similarly,

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

(easy to prove, using the definition.)

- Remember: $\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$

- Similarly,

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

(easy to prove, using the definition.)

- $\mathbb{E}[X_1 + \dots + X_n] = \mathbb{E}[X_1] + \dots + \mathbb{E}[X_n]$
- $\mathbb{E}[2X + 3Y - Z] = 2\mathbb{E}[X] + 3\mathbb{E}[Y] - \mathbb{E}[Z]$

- Remember: $\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$

- Similarly,

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

(easy to prove, using the definition.)

- $\mathbb{E}[X_1 + \dots + X_n] = \mathbb{E}[X_1] + \dots + \mathbb{E}[X_n]$
- $\mathbb{E}[2X + 3Y - Z] = 2\mathbb{E}[X] + 3\mathbb{E}[Y] - \mathbb{E}[Z]$

- Example.** Mean of a binomial rv Y with (n, p)
- Y : number of successes in n Bernoulli trials with p

- Remember: $\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$

- Similarly,

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

(easy to prove, using the definition.)

- $\mathbb{E}[X_1 + \dots + X_n] = \mathbb{E}[X_1] + \dots + \mathbb{E}[X_n]$
- $\mathbb{E}[2X + 3Y - Z] = 2\mathbb{E}[X] + 3\mathbb{E}[Y] - \mathbb{E}[Z]$

- Example.** Mean of a binomial rv Y with (n, p)
- Y : number of successes in n Bernoulli trials with p
- $Y = X_1 + \dots + X_n$, where X_i is a Bernoulli rv.
- $\mathbb{E}[Y] = n\mathbb{E}[X_i] = n\mathbb{P}(X_i = 1) = np$

- Remember: $\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$

- Similarly,

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

(easy to prove, using the definition.)

- $\mathbb{E}[X_1 + \dots + X_n] = \mathbb{E}[X_1] + \dots + \mathbb{E}[X_n]$
- $\mathbb{E}[2X + 3Y - Z] = 2\mathbb{E}[X] + 3\mathbb{E}[Y] - \mathbb{E}[Z]$

- Example.** Mean of a binomial rv Y with (n, p)

- Y : number of successes in n Bernoulli trials with p

- $Y = X_1 + \dots + X_n$, where X_i is a Bernoulli rv.

- $\mathbb{E}[Y] = n\mathbb{E}[X_i] = n\mathbb{P}(X_i = 1) = np$

Message. When some rv X is write as a linear combination of other rvs, it is often easy to handle X .

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) **Conditioning for random variables**
- (6) Independence for random variables

Remember two probability laws: $\mathbb{P}(\cdot)$ and $\mathbb{P}(\cdot|A)$ for an event A .

Remember two probability laws: $\mathbb{P}(\cdot)$ and $\mathbb{P}(\cdot|A)$ for an event A .

- $p_X(x) \triangleq \mathbb{P}(X = x)$

- $p_{X|A}(x) \triangleq \mathbb{P}(X = x|A)$

Remember two probability laws: $\mathbb{P}(\cdot)$ and $\mathbb{P}(\cdot|A)$ for an event A .

- $p_X(x) \triangleq \mathbb{P}(X = x)$

- $\mathbb{E}[X] = \sum_x x p_X(x)$

- $p_{X|A}(x) \triangleq \mathbb{P}(X = x|A)$

- $\mathbb{E}[X|A] \triangleq \sum_x x p_{X|A}(x)$

Remember two probability laws: $\mathbb{P}(\cdot)$ and $\mathbb{P}(\cdot|A)$ for an event A .

- | | |
|--|--|
| <ul style="list-style-type: none">• $p_X(x) \triangleq \mathbb{P}(X = x)$• $\mathbb{E}[X] = \sum_x x p_X(x)$• $\mathbb{E}[g(X)] = \sum_x g(x) p_X(x)$ | <ul style="list-style-type: none">• $p_{X A}(x) \triangleq \mathbb{P}(X = x A)$• $\mathbb{E}[X A] \triangleq \sum_x x p_{X A}(x)$• $\mathbb{E}[g(X) A] \triangleq \sum_x g(x) p_{X A}(x)$ |
|--|--|


Remember two probability laws: $\mathbb{P}(\cdot)$ and $\mathbb{P}(\cdot|A)$ for an event A .

- $p_X(x) \triangleq \mathbb{P}(X = x)$
 - $\mathbb{E}[X] = \sum_x x p_X(x)$
 - $\mathbb{E}[g(X)] = \sum_x g(x) p_X(x)$
 - $\text{var}[X] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$
- $p_{X|A}(x) \triangleq \mathbb{P}(X = x|A)$
 - $\mathbb{E}[X|A] \triangleq \sum_x x p_{X|A}(x)$
 - $\mathbb{E}[g(X)|A] \triangleq \sum_x g(x) p_{X|A}(x)$
 - $\text{var}[X|A] \triangleq \mathbb{E}[X^2|A] - (\mathbb{E}[X|A])^2$

Remember two probability laws: $\mathbb{P}(\cdot)$ and $\mathbb{P}(\cdot|A)$ for an event A .


- $p_X(x) \triangleq \mathbb{P}(X = x)$
 - $\mathbb{E}[X] = \sum_x x p_X(x)$
 - $\mathbb{E}[g(X)] = \sum_x g(x) p_X(x)$
 - $\text{var}[X] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$
- $p_{X|A}(x) \triangleq \mathbb{P}(X = x|A)$
 - $\mathbb{E}[X|A] \triangleq \sum_x x p_{X|A}(x)$
 - $\mathbb{E}[g(X)|A] \triangleq \sum_x g(x) p_{X|A}(x)$
 - $\text{var}[X|A] \triangleq \mathbb{E}[X^2|A] - (\mathbb{E}[X|A])^2$
 - (Note) $p_{X|A}(x)$, $\mathbb{E}[X|A]$, $\mathbb{E}[g(X)|A]$, and $\text{var}[X|A]$ are all just notations!

$$A = \{X \geq 2\}$$



$$\mathbb{E}[X] =$$


$$\text{var}[X] =$$



$$\mathbb{E}[X|A] =$$


$$\text{var}[X|A] =$$

$$A = \{X \geq 2\}$$



$$\mathbb{E}[X] = \frac{1}{4} (1 + 2 + 3 + 4) = 2.5$$


$$\text{var}[X] =$$



$$\mathbb{E}[X|A] =$$


$$\text{var}[X|A] =$$

$$A = \{X \geq 2\}$$



$$\mathbb{E}[X] = \frac{1}{4} (1 + 2 + 3 + 4) = 2.5$$


$$\begin{aligned} \text{var}[X] &= \mathbb{E}[X^2] - (\mathbb{E}[X])^2 \\ &= \frac{1}{4} (1 + 2^2 + 3^2 + 4^2) - 2.5^2 \end{aligned}$$



$$\mathbb{E}[X|A] =$$


$$\text{var}[X|A] =$$

$$A = \{X \geq 2\}$$



$$\mathbb{E}[X] = \frac{1}{4}(1 + 2 + 3 + 4) = 2.5$$


$$\begin{aligned}\text{var}[X] &= \mathbb{E}[X^2] - (\mathbb{E}[X])^2 \\ &= \frac{1}{4}(1 + 2^2 + 3^2 + 4^2) - 2.5^2\end{aligned}$$



$$\mathbb{E}[X|A] = \frac{1}{3}(2 + 3 + 4) = 3$$


$$\text{var}[X|A] =$$

$$A = \{X \geq 2\}$$



$$\mathbb{E}[X] = \frac{1}{4}(1 + 2 + 3 + 4) = 2.5$$


$$\begin{aligned}\text{var}[X] &= \mathbb{E}[X^2] - (\mathbb{E}[X])^2 \\ &= \frac{1}{4}(1 + 2^2 + 3^2 + 4^2) - 2.5^2\end{aligned}$$



$$\mathbb{E}[X|A] = \frac{1}{3}(2 + 3 + 4) = 3$$


$$\text{var}[X|A] = \mathbb{E}[X^2|A] - (\mathbb{E}[X|A])^2$$

$$A = \{X \geq 2\}$$



$$\mathbb{E}[X] = \frac{1}{4}(1 + 2 + 3 + 4) = 2.5$$

$$\begin{aligned}\text{var}[X] &= \mathbb{E}[X^2] - (\mathbb{E}[X])^2 \\ &= \frac{1}{4}(1 + 2^2 + 3^2 + 4^2) - 2.5^2\end{aligned}$$



$$\mathbb{E}[X|A] = \frac{1}{3}(2 + 3 + 4) = 3$$

$$\begin{aligned}\text{var}[X|A] &= \mathbb{E}[X^2|A] - (\mathbb{E}[X|A])^2 \\ &= \frac{1}{3}(2^2 + 3^2 + 4^2) - 3^2 = 2/3\end{aligned}$$

What do we mean by “conditioning on a rv”? Consider $A = \{Y = y\}$ for a rv Y .



What do we mean by “conditioning on a rv”? Consider $A = \{Y = y\}$ for a rv Y .

- $p_{X|A}(x) \triangleq \mathbb{P}(X = x|A)$

- $p_{X|Y}(x|y) \triangleq \mathbb{P}(X = x|Y = y)$

What do we mean by “conditioning on a rv”? Consider $A = \{Y = y\}$ for a rv Y .

- $p_{X|A}(x) \triangleq \mathbb{P}(X = x|A)$

- $\mathbb{E}[X|A] \triangleq \sum_x x p_{X|A}(x)$

- $p_{X|Y}(x|y) \triangleq \mathbb{P}(X = x|Y = y)$

- $\mathbb{E}[X|Y = y] \triangleq \sum_x x p_{X|Y}(x|y)$

What do we mean by “conditioning on a rv”? Consider $A = \{Y = y\}$ for a rv Y .

- $p_{X|A}(x) \triangleq \mathbb{P}(X = x|A)$

- $\mathbb{E}[X|A] \triangleq \sum_x x p_{X|A}(x)$

- $\mathbb{E}[g(X)|A] \triangleq \sum_x g(x) p_{X|A}(x)$

- $p_{X|Y}(x|y) \triangleq \mathbb{P}(X = x|Y = y)$

- $\mathbb{E}[X|Y = y] \triangleq \sum_x x p_{X|Y}(x|y)$

- $\mathbb{E}[g(X)|Y = y] \triangleq \sum_x g(x) p_{X|Y}(x|y)$

What do we mean by “conditioning on a rv”? Consider $A = \{Y = y\}$ for a rv Y .

- $p_{X|A}(x) \triangleq \mathbb{P}(X = x|A)$
- $\mathbb{E}[X|A] \triangleq \sum_x x p_{X|A}(x)$
- $\mathbb{E}[g(X)|A] \triangleq \sum_x g(x) p_{X|A}(x)$
- $\text{var}[X|A] \triangleq \mathbb{E}[X^2|A] - (\mathbb{E}[X|A])^2$

- $p_{X|Y}(x|y) \triangleq \mathbb{P}(X = x|Y = y)$
- $\mathbb{E}[X|Y = y] \triangleq \sum_x x p_{X|Y}(x|y)$
- $\mathbb{E}[g(X)|Y = y] \triangleq \sum_x g(x) p_{X|Y}(x|y)$
- $\text{var}[X|Y = y] \triangleq \mathbb{E}[X^2|Y = y] - (\mathbb{E}[X|Y = y])^2$

- Conditional PMF

- Multiplication rule.

$$p_{X,Y}(x,y) =$$

- $p_{X,Y,Z}(x,y,z) =$

- Conditional PMF

$$p_{X|Y}(x|y) \triangleq \mathbb{P}(X = x|Y = y) = \frac{p_{X,Y}(x,y)}{p_Y(y)}$$

for y such that $p_Y(y) > 0$.

- Multiplication rule.

$$p_{X,Y}(x,y) =$$

- $p_{X,Y,Z}(x,y,z) =$

- Conditional PMF

$$p_{X|Y}(x|y) \triangleq \mathbb{P}(X = x|Y = y) = \frac{p_{X,Y}(x,y)}{p_Y(y)}$$

for y such that $p_Y(y) > 0$.

- $\sum_x p_{X|Y}(x|y) = 1$

- Multiplication rule.

$$p_{X,Y}(x,y) =$$

- $p_{X,Y,Z}(x,y,z) =$

- Conditional PMF

$$p_{X|Y}(x|y) \triangleq \mathbb{P}(X = x|Y = y) = \frac{p_{X,Y}(x,y)}{p_Y(y)}$$

for y such that $p_Y(y) > 0$.

- $\sum_x p_{X|Y}(x|y) = 1$

- Multiplication rule.

$$\begin{aligned} p_{X,Y}(x,y) &= p_Y(y)p_{X|Y}(x|y) \\ &= p_X(x)p_{Y|X}(y|x) \end{aligned}$$

- $p_{X,Y,Z}(x,y,z) =$

- Conditional PMF

$$p_{X|Y}(x|y) \triangleq \mathbb{P}(X = x|Y = y) = \frac{p_{X,Y}(x,y)}{p_Y(y)}$$

for y such that $p_Y(y) > 0$.

- $\sum_x p_{X|Y}(x|y) = 1$

- Multiplication rule.

$$\begin{aligned} p_{X,Y}(x,y) &= p_Y(y)p_{X|Y}(x|y) \\ &= p_X(x)p_{Y|X}(y|x) \end{aligned}$$

- $p_{X,Y,Z}(x,y,z) = p_X(x)p_{Y|X}(y|x)p_{Z|X,Y}(z|x,y)$

VIDEO PAUSE

4	1/20	2/20	2/20	
3	2/20	4/20	1/20	2/20
2		1/20	3/20	1/20
1		1/20		
	1	2	3	4

$$p_{X|Y}(2|2) =$$

$$p_{X|Y}(3|2) =$$

$$\mathbb{E}[X|Y = 3] =$$

- Conditional PMF

$$p_{X|Y}(x|y) \triangleq \mathbb{P}(X = x|Y = y) = \frac{p_{X,Y}(x,y)}{p_Y(y)}$$

for y such that $p_Y(y) > 0$.

- $\sum_x p_{X|Y}(x|y) = 1$

- Multiplication rule.

$$\begin{aligned} p_{X,Y}(x,y) &= p_Y(y)p_{X|Y}(x|y) \\ &= p_X(x)p_{Y|X}(y|x) \end{aligned}$$

- $p_{X,Y,Z}(x,y,z) = p_X(x)p_{Y|X}(y|x)p_{Z|X,Y}(z|x,y)$

VIDEO PAUSE

4	1/20	2/20	2/20	
3	2/20	4/20	1/20	2/20
2		1/20	3/20	1/20
1		1/20		
	1	2	3	4

$$p_{X|Y}(2|2) = \frac{1}{1+3+1}$$



$$p_{X|Y}(3|2) = \frac{3}{1+3+1}$$

$$\mathbb{E}[X|Y = 3] = 1(2/9) + 2(4/9) + 3(1/9) + 4(2/9)$$

- Partition of Ω into A_1, A_2, A_3
- Known: $\mathbb{P}(A_i)$ and $\mathbb{P}(B|A_i)$
- What is $\mathbb{P}(B)$? (probability of result)

Total Probability Theorem


$$\mathbb{P}(B) = \sum_i \mathbb{P}(A_i) \mathbb{P}(B|A_i)$$



- Partition of Ω into A_1, A_2, A_3

Total Probability Theorem

$$p_X(x) = \sum_i \mathbb{P}(A_i) \mathbb{P}(X = x | A_i) = \sum_i \mathbb{P}(A_i) p_{X|A_i}(x)$$




- Partition of Ω into A_1, A_2, A_3

Total Probability Theorem

$$p_X(x) = \sum_i \mathbb{P}(A_i) \mathbb{P}(X = x | A_i) = \sum_i \mathbb{P}(A_i) p_{X|A_i}(x)$$

Total Expectation Theorem


$$\mathbb{E}[X] = \sum_i \mathbb{P}(A_i) \mathbb{E}[X | A_i]$$



- Partition of Ω into A_1, A_2, A_3

Total Expectation Theorem

$$\mathbb{E}[X] = \sum_i \mathbb{P}(A_i) \mathbb{E}[X|A_i]$$





- Partition of Ω into A_1, A_2, A_3

Total Expectation Theorem

$$\mathbb{E}[X] = \sum_i \mathbb{P}(A_i) \mathbb{E}[X|A_i]$$


Total Expectation Theorem

$$\mathbb{E}[X] = \sum_y \mathbb{P}(Y = y) \mathbb{E}[X|Y = y] = \sum_y p_Y(y) \mathbb{E}[X|Y = y]$$




- Using the definition of expectation,

$$\mathbb{E}[X] =$$



- Using the definition of expectation,

$$\begin{aligned}\mathbb{E}[X] &= \frac{1}{9}(0 + 1 + 2) + \frac{2}{9}(6 + 7 + 8) \\ &= \frac{3 + 12 + 14 + 16}{9} = 5\end{aligned}$$




- Using the definition of expectation,

$$\begin{aligned}\mathbb{E}[X] &= \frac{1}{9}(0 + 1 + 2) + \frac{2}{9}(6 + 7 + 8) \\ &= \frac{3 + 12 + 14 + 16}{9} = 5\end{aligned}$$

- Let's use TET, for which consider

$$A_1 = \{X \in \{0, 1, 2\}\}, \quad A_2 = \{X \in \{6, 7, 8\}\}$$




- Using the definition of expectation,

$$\begin{aligned}\mathbb{E}[X] &= \frac{1}{9}(0 + 1 + 2) + \frac{2}{9}(6 + 7 + 8) \\ &= \frac{3 + 12 + 14 + 16}{9} = 5\end{aligned}$$

- Let's use TET, for which consider

$$A_1 = \{X \in \{0, 1, 2\}\}, \quad A_2 = \{X \in \{6, 7, 8\}\}$$

$$\begin{aligned}\mathbb{E}[X] &= \sum_{i=1,2} \mathbb{P}(A_i) \mathbb{E}[X|A_i] \\ &= 1/3 \cdot 1 + 2/3 \cdot 7 = 5\end{aligned}$$



- Some random variable often does not have **memory**.

- Some random variable often does not have **memory**.
- **Definition.** A random variable X is called **memoryless** if, for any $n, m \geq 0$,

$$\mathbb{P}(X > n + m | X > m) = \mathbb{P}(X > n)$$

- Some random variable often does not have **memory**.
- **Definition.** A random variable X is called **memoryless** if, for any $n, m \geq 0$,

$$\mathbb{P}(X > n + m | X > m) = \mathbb{P}(X > n)$$

- **Meaning.** Conditioned on $X > m$, $X - m$'s distribution is the same as the original X .

$$\mathbb{P}(X - m > n | X > m) = \mathbb{P}(X > n)$$

- Some random variable often does not have **memory**.
- **Definition.** A random variable X is called **memoryless** if, for any $n, m \geq 0$,

$$\mathbb{P}(X > n + m | X > m) = \mathbb{P}(X > n)$$

- **Meaning.** Conditioned on $X > m$, $X - m$'s distribution is the same as the original X .

$$\mathbb{P}(X - m > n | X > m) = \mathbb{P}(X > n)$$

- Suppose that X is the time of waiting for a bus and X is memoryless. At the bus stop, I have waited for the bus for 10 mins. Then, the time until the bus arrival does not depend on how much I have waited for a bus. **No memory.**

- **Theorem.** Any geometric random variable is **memoryless**.

- **Theorem.** Any geometric random variable is **memoryless**.
- **Remind.** Geometric rv X with parameter p

$$\mathbb{P}(X = k) = (1 - p)^{k-1}p, \quad \mathbb{P}(X > k) = 1 - \sum_{k'=1}^k (1 - p)^{k'-1}p = (1 - p)^k$$

- **Theorem.** Any geometric random variable is **memoryless**.
- **Remind.** Geometric rv X with parameter p

$$\mathbb{P}(X = k) = (1 - p)^{k-1}p, \quad \mathbb{P}(X > k) = 1 - \sum_{k'=1}^k (1 - p)^{k'-1}p = (1 - p)^k$$

- **Proof.**

$$\begin{aligned}\mathbb{P}(X > n + m | X > m) &= \frac{\mathbb{P}(X > n + m \text{ and } X > m)}{\mathbb{P}(X > m)} = \frac{\mathbb{P}(X > n + m)}{\mathbb{P}(X > m)} \\ &= \frac{(1 - p)^{n+m}}{(1 - p)^m} = (1 - p)^n = \mathbb{P}(X > n)\end{aligned}$$

- **Theorem.** Any geometric random variable is **memoryless**.
- **Remind.** Geometric rv X with parameter p

$$\mathbb{P}(X = k) = (1 - p)^{k-1}p, \quad \mathbb{P}(X > k) = 1 - \sum_{k'=1}^k (1 - p)^{k'-1}p = (1 - p)^k$$

- **Proof.**

$$\begin{aligned}\mathbb{P}(X > n + m | X > m) &= \frac{\mathbb{P}(X > n + m \text{ and } X > m)}{\mathbb{P}(X > m)} = \frac{\mathbb{P}(X > n + m)}{\mathbb{P}(X > m)} \\ &= \frac{(1 - p)^{n+m}}{(1 - p)^m} = (1 - p)^n = \mathbb{P}(X > n)\end{aligned}$$

- **Meaning.** Conditioned on $X > m$, $X - m$ is geometric with the same parameter.

- Write softwares over and over, and each time w.p. p of working correctly (independent from prev. programs).

- Write softwares over and over, and each time w.p. p of working correctly (independent from prev. programs).
- X : number of trials until the program works correctly.

- Write softwares over and over, and each time w.p. p of working correctly (independent from prev. programs).
- X : number of trials until the program works correctly.
- (Q) mean of X

- Write softwares over and over, and each time w.p. p of working correctly (independent from prev. programs).
- X : number of trials until the program works correctly.
- (Q) mean of X
- X is geometric

- Write softwares over and over, and each time w.p. p of working correctly (independent from prev. programs).
- X : number of trials until the program works correctly.
- (Q) mean of X
- X is geometric
- Direct computation is boring.

$$\mathbb{E}[X] = \sum_{k=1}^{\infty} k(1-p)^{k-1}p$$

- Write softwares over and over, and each time w.p. p of working correctly (independent from prev. programs).
- X : number of trials until the program works correctly.
- (Q) mean of X
- X is geometric
- Direct computation is boring.

$$\mathbb{E}[X] = \sum_{k=1}^{\infty} k(1-p)^{k-1}p$$

- Total expectation theorem and memorylessness helps a lot.

- Write softwares over and over, and each time w.p. p of working correctly (independent from prev. programs).
- X : number of trials until the program works correctly.
- (Q) mean of X
- X is geometric
- Direct computation is boring.

$$\mathbb{E}[X] = \sum_{k=1}^{\infty} k(1-p)^{k-1}p$$

- Total expectation theorem and memorylessness helps a lot.

- $A_1 = \{X = 1\}$ (first try is success),
 $A_2 = \{X > 1\}$ (first try is failure).

$$\mathbb{E}[X] = 1 + \mathbb{E}[X - 1]$$

=

=

- Write softwares over and over, and each time w.p. p of working correctly (independent from prev. programs).
- X : number of trials until the program works correctly.
- (Q) mean of X
- X is geometric
- Direct computation is boring.

$$\mathbb{E}[X] = \sum_{k=1}^{\infty} k(1-p)^{k-1}p$$

- Total expectation theorem and memorylessness helps a lot.

- $A_1 = \{X = 1\}$ (first try is success),
 $A_2 = \{X > 1\}$ (first try is failure).

$$\begin{aligned}\mathbb{E}[X] &= 1 + \mathbb{E}[X - 1] \\ &= 1 + \mathbb{P}(A_1)\mathbb{E}[X - 1|X = 1] \\ &\quad + \mathbb{P}(A_2)\mathbb{E}[X - 1|X > 1] \\ &= \end{aligned}$$

- Write softwares over and over, and each time w.p. p of working correctly (independent from prev. programs).
- X : number of trials until the program works correctly.
- (Q) mean of X
- X is geometric
- Direct computation is boring.

$$\mathbb{E}[X] = \sum_{k=1}^{\infty} k(1-p)^{k-1}p$$

- Total expectation theorem and memorylessness helps a lot.

- $A_1 = \{X = 1\}$ (first try is success),
 $A_2 = \{X > 1\}$ (first try is failure).

$$\begin{aligned}\mathbb{E}[X] &= 1 + \mathbb{E}[X - 1] \\ &= 1 + \mathbb{P}(A_1)\mathbb{E}[X - 1|X = 1] \\ &\quad + \mathbb{P}(A_2)\mathbb{E}[X - 1|X > 1] \\ &= 1 + (1 - p)\mathbb{E}[X]\end{aligned}$$

- Write softwares over and over, and each time w.p. p of working correctly (independent from prev. programs).
- X : number of trials until the program works correctly.
- (Q) mean of X
- X is geometric
- Direct computation is boring.

$$\mathbb{E}[X] = \sum_{k=1}^{\infty} k(1-p)^{k-1}p$$

- Total expectation theorem and memorylessness helps a lot.

- $A_1 = \{X = 1\}$ (first try is success),
 $A_2 = \{X > 1\}$ (first try is failure).

$$\begin{aligned}\mathbb{E}[X] &= 1 + \mathbb{E}[X - 1] \\ &= 1 + \mathbb{P}(A_1)\mathbb{E}[X - 1|X = 1] \\ &\quad + \mathbb{P}(A_2)\mathbb{E}[X - 1|X > 1] \\ &= 1 + (1 - p)\mathbb{E}[X]\end{aligned}$$

$$\mathbb{E}[X] = 1 + (1 - p)\frac{1}{p} = 1/p.$$

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

- Two events

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$$

$$\mathbb{P}(A \cap B | C) = \mathbb{P}(A | C) \cdot \mathbb{P}(B | C)$$

- Two events

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$$

$$\mathbb{P}(A \cap B | C) = \mathbb{P}(A | C) \cdot \mathbb{P}(B | C)$$

- A rv and an event

$$\mathbb{P}(\{X = x\} \cap B) = \mathbb{P}(X = x) \cdot \mathbb{P}(B), \quad \text{for all } x$$

$$\mathbb{P}(\{X = x\} \cap B | C) = \mathbb{P}(X = x | C) \cdot \mathbb{P}(B | C), \quad \text{for all } x$$

- Two events

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$$

$$\mathbb{P}(A \cap B | C) = \mathbb{P}(A | C) \cdot \mathbb{P}(B | C)$$

- A rv and an event

$$\mathbb{P}(\{X = x\} \cap B) = \mathbb{P}(X = x) \cdot \mathbb{P}(B), \quad \text{for all } x$$

$$\mathbb{P}(\{X = x\} \cap B | C) = \mathbb{P}(X = x | C) \cdot \mathbb{P}(B | C), \quad \text{for all } x$$

- Two rvs

$$\mathbb{P}(\{X = x\} \cap \{Y = y\}) = \mathbb{P}(X = x) \cdot \mathbb{P}(Y = y), \quad \text{for all } x, y$$

$$\mathbb{P}(\{X = x\} \cap \{Y = y\} | Z = z) = \mathbb{P}(X = x | Z = z) \cdot \mathbb{P}(Y = y | Z = z), \quad \text{for all } x, y$$

- Two events

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$$

$$\mathbb{P}(A \cap B | C) = \mathbb{P}(A | C) \cdot \mathbb{P}(B | C)$$

- A rv and an event

$$\mathbb{P}(\{X = x\} \cap B) = \mathbb{P}(X = x) \cdot \mathbb{P}(B), \quad \text{for all } x$$

$$\mathbb{P}(\{X = x\} \cap B | C) = \mathbb{P}(X = x | C) \cdot \mathbb{P}(B | C), \quad \text{for all } x$$

- Two rvs

$$\mathbb{P}(\{X = x\} \cap \{Y = y\}) = \mathbb{P}(X = x) \cdot \mathbb{P}(Y = y), \quad \text{for all } x, y$$


$$p_{X,Y}(x, y) = p_X(x) \cdot p_Y(y)$$

$$\mathbb{P}(\{X = x\} \cap \{Y = y\} | Z = z) = \mathbb{P}(X = x | Z = z) \cdot \mathbb{P}(Y = y | Z = z), \quad \text{for all } x, y$$

$$p_{X,Y|Z}(x, y) = p_{X|Z}(x) \cdot p_{Y|Z}(y)$$

- $X \perp\!\!\!\perp Y$?

- $X \perp\!\!\!\perp Y | \{X \leq 2 \text{ and } Y \geq 3\}$?




- $X \perp\!\!\!\perp Y$?

$$p_{X,Y}(1,1) = 0, \quad p_X(1) = 3/20$$

$$p_Y(1) = 1/20.$$

- $X \perp\!\!\!\perp Y | \{X \leq 2 \text{ and } Y \geq 3\}$?




- $X \perp\!\!\!\perp Y$?

$$p_{X,Y}(1,1) = 0, \quad p_X(1) = 3/20$$

$$p_Y(1) = 1/20.$$

- $X \perp\!\!\!\perp Y | \{X \leq 2 \text{ and } Y \geq 3\}$?
- Yes.



$Y = 4 \ (1/3)$	$1/9$	$2/9$
$Y = 3 \ (2/3)$	$2/9$	$4/9$
	$X = 1 \ (1/3)$	$X = 2 \ (2/3)$

- Always true.

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

- Always true.

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

- Generally, $\mathbb{E}[g(X, Y)] \neq g(\mathbb{E}[X], \mathbb{E}[Y])$

- Always true.

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

- Generally, $\mathbb{E}[g(X, Y)] \neq g(\mathbb{E}[X], \mathbb{E}[Y])$

- However, if $X \perp\!\!\!\perp Y$,

$$\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$$

$$\mathbb{E}[g(X)h(Y)] = \mathbb{E}[g(X)]\mathbb{E}[h(Y)]$$

- Always true.

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

- Generally, $\mathbb{E}[g(X, Y)] \neq g(\mathbb{E}[X], \mathbb{E}[Y])$

- However, if $X \perp\!\!\!\perp Y$,

$$\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$$

$$\mathbb{E}[g(X)h(Y)] = \mathbb{E}[g(X)]\mathbb{E}[h(Y)]$$

- Proof.

$$\begin{aligned}\mathbb{E}[g(X)h(Y)] &= \sum_x \sum_y g(x)h(y)p_{X,Y}(x,y) \\ &= \sum_x x p_X(x) \sum_y y p_Y(y)\end{aligned}$$

- Always true.

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

- Generally, $\mathbb{E}[g(X, Y)] \neq g(\mathbb{E}[X], \mathbb{E}[Y])$

- However, if $X \perp\!\!\!\perp Y$,

$$\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$$

$$\mathbb{E}[g(X)h(Y)] = \mathbb{E}[g(X)]\mathbb{E}[h(Y)]$$

- Proof.

$$\begin{aligned}\mathbb{E}[g(X)h(Y)] &= \sum_x \sum_y g(x)h(y)p_{X,Y}(x,y) \\ &= \sum_x xp_X(x) \sum_y yp_Y(y)\end{aligned}$$

- Always true.

$$\text{var}[aX] = a^2\text{var}[X], \text{var}[X + a] = \text{var}[X]$$

- Always true.

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

- Generally, $\mathbb{E}[g(X, Y)] \neq g(\mathbb{E}[X], \mathbb{E}[Y])$

- However, if $X \perp\!\!\!\perp Y$,

$$\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$$

$$\mathbb{E}[g(X)h(Y)] = \mathbb{E}[g(X)]\mathbb{E}[h(Y)]$$

- Proof.

$$\begin{aligned}\mathbb{E}[g(X)h(Y)] &= \sum_x \sum_y g(x)h(y)p_{X,Y}(x,y) \\ &= \sum_x xp_X(x) \sum_y yp_Y(y)\end{aligned}$$

- Always true.

$$\text{var}[aX] = a^2\text{var}[X], \text{var}[X + a] = \text{var}[X]$$

- Generally, $\text{var}[X + Y] \neq \text{var}[X] + \text{var}[Y]$

- Always true.

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

- Generally, $\mathbb{E}[g(X, Y)] \neq g(\mathbb{E}[X], \mathbb{E}[Y])$

- However, if $X \perp\!\!\!\perp Y$,

$$\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$$

$$\mathbb{E}[g(X)h(Y)] = \mathbb{E}[g(X)]\mathbb{E}[h(Y)]$$

- Proof.

$$\begin{aligned}\mathbb{E}[g(X)h(Y)] &= \sum_x \sum_y g(x)h(y)p_{X,Y}(x,y) \\ &= \sum_x x p_X(x) \sum_y y p_Y(y)\end{aligned}$$

- Always true.

$$\text{var}[aX] = a^2\text{var}[X], \text{var}[X + a] = \text{var}[X]$$

- Generally, $\text{var}[X + Y] \neq \text{var}[X] + \text{var}[Y]$

- However, if $X \perp\!\!\!\perp Y$,

$$\text{var}[X + Y] = \text{var}[X] + \text{var}[Y]$$

- Practice.

- Always true.

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

- Generally, $\mathbb{E}[g(X, Y)] \neq g(\mathbb{E}[X], \mathbb{E}[Y])$

- However, if $X \perp\!\!\!\perp Y$,

$$\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$$

$$\mathbb{E}[g(X)h(Y)] = \mathbb{E}[g(X)]\mathbb{E}[h(Y)]$$

- **Proof.**

$$\begin{aligned}\mathbb{E}[g(X)h(Y)] &= \sum_x \sum_y g(x)h(y)p_{X,Y}(x,y) \\ &= \sum_x xp_X(x) \sum_y yp_Y(y)\end{aligned}$$

- Always true.

$$\text{var}[aX] = a^2\text{var}[X], \text{var}[X + a] = \text{var}[X]$$

- Generally, $\text{var}[X + Y] \neq \text{var}[X] + \text{var}[Y]$

- However, if $X \perp\!\!\!\perp Y$,

$$\text{var}[X + Y] = \text{var}[X] + \text{var}[Y]$$

- Practice.

- $X = Y \implies \text{var}[X + Y] = 4\text{var}[X]$

- Always true.

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

- Generally, $\mathbb{E}[g(X, Y)] \neq g(\mathbb{E}[X], \mathbb{E}[Y])$

- However, if $X \perp\!\!\!\perp Y$,

$$\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$$

$$\mathbb{E}[g(X)h(Y)] = \mathbb{E}[g(X)]\mathbb{E}[h(Y)]$$

- **Proof.**

$$\begin{aligned}\mathbb{E}[g(X)h(Y)] &= \sum_x \sum_y g(x)h(y)p_{X,Y}(x,y) \\ &= \sum_x xp_X(x) \sum_y yp_Y(y)\end{aligned}$$

- Always true.

$$\text{var}[aX] = a^2\text{var}[X], \text{var}[X + a] = \text{var}[X]$$

- Generally, $\text{var}[X + Y] \neq \text{var}[X] + \text{var}[Y]$

- However, if $X \perp\!\!\!\perp Y$,

$$\text{var}[X + Y] = \text{var}[X] + \text{var}[Y]$$

- Practice.

- $X = Y \implies \text{var}[X + Y] = 4\text{var}[X]$
- $X = -Y \implies \text{var}[X + Y] = 0$

- Always true.

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

- Generally, $\mathbb{E}[g(X, Y)] \neq g(\mathbb{E}[X], \mathbb{E}[Y])$

- However, if $X \perp\!\!\!\perp Y$,

$$\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$$

$$\mathbb{E}[g(X)h(Y)] = \mathbb{E}[g(X)]\mathbb{E}[h(Y)]$$

- **Proof.**

$$\begin{aligned}\mathbb{E}[g(X)h(Y)] &= \sum_x \sum_y g(x)h(y)p_{X,Y}(x,y) \\ &= \sum_x xp_X(x) \sum_y yp_Y(y)\end{aligned}$$

- Always true.

$$\text{var}[aX] = a^2\text{var}[X], \text{var}[X + a] = \text{var}[X]$$

- Generally, $\text{var}[X + Y] \neq \text{var}[X] + \text{var}[Y]$

- However, if $X \perp\!\!\!\perp Y$,

$$\text{var}[X + Y] = \text{var}[X] + \text{var}[Y]$$

- Practice.

- $X = Y \implies \text{var}[X + Y] = 4\text{var}[X]$
- $X = -Y \implies \text{var}[X + Y] = 0$
- $X \perp\!\!\!\perp Y \implies \text{var}[X - 3Y] = \text{var}[X] + 9\text{var}[Y]$

$$\text{var}[X + Y] \neq \text{var}[X] + \text{var}[Y]$$

- Why not generally true?

$$\text{var}[X + Y] \neq \text{var}[X] + \text{var}[Y]$$

- Why not generally true?

$$\begin{aligned}\text{var}[X + Y] &= \mathbb{E}[(X + Y)^2] - (\mathbb{E}[X + Y])^2 \\ &= \mathbb{E}[X^2 + Y^2 + 2XY] - ((\mathbb{E}[X])^2 + (\mathbb{E}[Y])^2 + 2\mathbb{E}[X]\mathbb{E}[Y]) \\ &= \text{var}[X] + \text{var}[Y] + 2(\mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y])\end{aligned}$$

$$\text{var}[X + Y] \neq \text{var}[X] + \text{var}[Y]$$

- Why not generally true?

$$\begin{aligned}\text{var}[X + Y] &= \mathbb{E}[(X + Y)^2] - (\mathbb{E}[X + Y])^2 \\ &= \mathbb{E}[X^2 + Y^2 + 2XY] - ((\mathbb{E}[X])^2 + (\mathbb{E}[Y])^2 + 2\mathbb{E}[X]\mathbb{E}[Y]) \\ &= \text{var}[X] + \text{var}[Y] + 2(\mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y])\end{aligned}$$

- is a sufficient condition for $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$

$$\text{var}[X + Y] \neq \text{var}[X] + \text{var}[Y]$$

- Why not generally true?

$$\begin{aligned}\text{var}[X + Y] &= \mathbb{E}[(X + Y)^2] - (\mathbb{E}[X + Y])^2 \\ &= \mathbb{E}[X^2 + Y^2 + 2XY] - ((\mathbb{E}[X])^2 + (\mathbb{E}[Y])^2 + 2\mathbb{E}[X]\mathbb{E}[Y]) \\ &= \text{var}[X] + \text{var}[Y] + 2(\mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y])\end{aligned}$$

- $X \perp\!\!\!\perp Y$ is a sufficient condition for $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$

$$\text{var}[X + Y] \neq \text{var}[X] + \text{var}[Y]$$

- Why not generally true?

$$\begin{aligned}\text{var}[X + Y] &= \mathbb{E}[(X + Y)^2] - (\mathbb{E}[X + Y])^2 \\ &= \mathbb{E}[X^2 + Y^2 + 2XY] - ((\mathbb{E}[X])^2 + (\mathbb{E}[Y])^2 + 2\mathbb{E}[X]\mathbb{E}[Y]) \\ &= \text{var}[X] + \text{var}[Y] + 2(\mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y])\end{aligned}$$

- $X \perp\!\!\!\perp Y$ is a sufficient condition for $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$
- Also, a necessary condition? we will see later, when we study **covariance**.

- n people throw their hats in a box and then pick one at random
- X : number of people with their own hat

- n people throw their hats in a box and then pick one at random
- X : number of people with their own hat
- $\mathbb{E}[X]$? $\text{var}[X]$?

- n people throw their hats in a box and then pick one at random
- X : number of people with their own hat
- $\mathbb{E}[X]$? $\text{var}[X]$?
- All permutations are equally likely as $1/n!$. Thus, this equals to picking one hat at a time.

- n people throw their hats in a box and then pick one at random
- X : number of people with their own hat
- $\mathbb{E}[X]$? $\text{var}[X]$?
- All permutations are equally likely as $1/n!$. Thus, this equals to picking one hat at a time.
- **Key step 1.** Define a rv $X_i = 1$ if i selects own hat and 0 otherwise.

$$X = \sum_{i=1}^n X_i.$$

- n people throw their hats in a box and then pick one at random
- X : number of people with their own hat
- $\mathbb{E}[X]$? $\text{var}[X]$?
- All permutations are equally likely as $1/n!$. Thus, this equals to picking one hat at a time.
- **Key step 1.** Define a rv $X_i = 1$ if i selects own hat and 0 otherwise.

$$X = \sum_{i=1}^n X_i.$$

- $\{X_i\}, i = 1, 2, \dots, n$: identically distributed (symmetry)

- $\mathbb{E}[X] = n\mathbb{E}[X_1] = n\mathbb{P}(X_1 = 1) = n \times \frac{1}{n} = 1.$

- $\mathbb{E}[X] = n\mathbb{E}[X_1] = n\mathbb{P}(X_1 = 1) = n \times \frac{1}{n} = 1.$
- **Key step 2.** Are X_i s are independent? If yes, easy to get $\text{var}(X).$

- $\mathbb{E}[X] = n\mathbb{E}[X_1] = n\mathbb{P}(X_1 = 1) = n \times \frac{1}{n} = 1$.
- **Key step 2.** Are X_i s are independent? If yes, easy to get $\text{var}(X)$.
- Assume $n = 2$. Then, $X_1 = 1 \rightarrow X_2 = 1$, and $X_1 = 0 \rightarrow X_2 = 0$. Thus, **dependent**.

- $\mathbb{E}[X] = n\mathbb{E}[X_1] = n\mathbb{P}(X_1 = 1) = n \times \frac{1}{n} = 1$.
- **Key step 2.** Are X_i s are independent? If yes, easy to get $\text{var}(X)$.
- Assume $n = 2$. Then, $X_1 = 1 \rightarrow X_2 = 1$, and $X_1 = 0 \rightarrow X_2 = 0$. Thus, **dependent**.

$$\begin{aligned}\text{var}(X) &= \mathbb{E}[X^2] - (\mathbb{E}[X])^2 \\ &= \mathbb{E}\left[\sum_i X_i^2 + \sum_{i,j:i \neq j} X_i X_j\right] - (\mathbb{E}[X])^2\end{aligned}$$

- $\mathbb{E}[X] = n\mathbb{E}[X_1] = n\mathbb{P}(X_1 = 1) = n \times \frac{1}{n} = 1$.
- **Key step 2.** Are X_i s are independent? If yes, easy to get $\text{var}(X)$.
- Assume $n = 2$. Then, $X_1 = 1 \rightarrow X_2 = 1$, and $X_1 = 0 \rightarrow X_2 = 0$. Thus, **dependent**.

$$\begin{aligned}\text{var}(X) &= \mathbb{E}[X^2] - (\mathbb{E}[X])^2 \\ &= \mathbb{E}\left[\sum_i X_i^2 + \sum_{i,j:i \neq j} X_i X_j\right] - (\mathbb{E}[X])^2\end{aligned}$$

$$\mathbb{E}[X_i^2] = 1 \times \frac{1}{n} + 0 \times \frac{n-1}{n} = \frac{1}{n}$$

- $\mathbb{E}[X] = n\mathbb{E}[X_1] = n\mathbb{P}(X_1 = 1) = n \times \frac{1}{n} = 1$.
- **Key step 2.** Are X_i s are independent? If yes, easy to get $\text{var}(X)$.
- Assume $n = 2$. Then, $X_1 = 1 \rightarrow X_2 = 1$, and $X_1 = 0 \rightarrow X_2 = 0$. Thus, **dependent**.

$$\begin{aligned}\text{var}(X) &= \mathbb{E}[X^2] - (\mathbb{E}[X])^2 \\ &= \mathbb{E}\left[\sum_i X_i^2 + \sum_{i,j:i \neq j} X_i X_j\right] - (\mathbb{E}[X])^2\end{aligned}$$

$$\mathbb{E}[X_i^2] = 1 \times \frac{1}{n} + 0 \times \frac{n-1}{n} = \frac{1}{n}$$

$$\mathbb{E}[X_i X_j] = \mathbb{E}[X_1 X_2] = 1 \times \mathbb{P}(X_1 X_2 = 1) = \mathbb{P}(X_1 = 1)\mathbb{P}(X_2 = 1|X_1 = 1), \quad (i \neq j)$$

- $\mathbb{E}[X] = n\mathbb{E}[X_1] = n\mathbb{P}(X_1 = 1) = n \times \frac{1}{n} = 1$.
- **Key step 2.** Are X_i s are independent? If yes, easy to get $\text{var}(X)$.
- Assume $n = 2$. Then, $X_1 = 1 \rightarrow X_2 = 1$, and $X_1 = 0 \rightarrow X_2 = 0$. Thus, **dependent**.

$$\begin{aligned}\text{var}(X) &= \mathbb{E}[X^2] - (\mathbb{E}[X])^2 \\ &= \mathbb{E}\left[\sum_i X_i^2 + \sum_{i,j:i \neq j} X_i X_j\right] - (\mathbb{E}[X])^2\end{aligned}$$

$$\mathbb{E}[X_i^2] = 1 \times \frac{1}{n} + 0 \times \frac{n-1}{n} = \frac{1}{n}$$

$$\mathbb{E}[X_i X_j] = \mathbb{E}[X_1 X_2] = 1 \times \mathbb{P}(X_1 X_2 = 1) = \mathbb{P}(X_1 = 1)\mathbb{P}(X_2 = 1|X_1 = 1), \quad (i \neq j)$$

- $\mathbb{E}[X^2] = n\mathbb{E}[X_1^2] + n(n-1)\mathbb{E}[X_1 X_2] = n\frac{1}{n} + n(n-1)\frac{1}{n(n-1)} = 2$

- $\mathbb{E}[X] = n\mathbb{E}[X_1] = n\mathbb{P}(X_1 = 1) = n \times \frac{1}{n} = 1$.
- **Key step 2.** Are X_i s are independent? If yes, easy to get $\text{var}(X)$.
- Assume $n = 2$. Then, $X_1 = 1 \rightarrow X_2 = 1$, and $X_1 = 0 \rightarrow X_2 = 0$. Thus, **dependent**.

$$\begin{aligned}\text{var}(X) &= \mathbb{E}[X^2] - (\mathbb{E}[X])^2 \\ &= \mathbb{E}\left[\sum_i X_i^2 + \sum_{i,j:i \neq j} X_i X_j\right] - (\mathbb{E}[X])^2\end{aligned}$$

$$\mathbb{E}[X_i^2] = 1 \times \frac{1}{n} + 0 \times \frac{n-1}{n} = \frac{1}{n}$$

$$\mathbb{E}[X_i X_j] = \mathbb{E}[X_1 X_2] = 1 \times \mathbb{P}(X_1 X_2 = 1) = \mathbb{P}(X_1 = 1) \mathbb{P}(X_2 = 1 | X_1 = 1), \quad (i \neq j)$$

- $\mathbb{E}[X^2] = n\mathbb{E}[X_1^2] + n(n-1)\mathbb{E}[X_1 X_2] = n\frac{1}{n} + n(n-1)\frac{1}{n(n-1)} = 2$
- $\text{var}(X) = 2 - 1 = 1$

Questions?

- 1) What is Random Variable? Why is it useful?
- 2) What is PMF (Probability Mass Function)?
- 3) Explain Bernoulli, Binomial, Poisson, Geometric rvs, when they are used and what their PMFs are.
- 4) What are joint and marginal PMFS?
- 5) Describe and explain the total probability/expectation theorem for random variables?
- 6) When is it useful to use total probability/expectation theorem?
- 7) What is conditional independence?