

## Lecture 5: Random Variable, Part III

Yi, Yung (이웅)

EE210: Probability and Introductory Random Processes  
KAIST EE

June 2, 2021

- (1) Derived distribution of  $Y = g(X)$  or  $Z = g(X, Y)$
- (2) Derived distribution of  $Z = X + Y$
- (3) Covariance: Degree of dependence between two rvs.
- (4) Correlation coefficient
- (5) Conditional expectation and law of iterative expectations
- (6) Conditional variance and law of total variance
- (7) Random number of sum of random variables

June 2, 2021 1 / 46

June 2, 2021 2 / 46

- (1) Derived distribution of  $Y = g(X)$  or  $Z = g(X, Y)$
- (2) Derived distribution of  $Z = X + Y$
- (3) Covariance: Degree of dependence between two rvs.
- (4) Correlation coefficient
- (5) Conditional expectation and law of iterative expectations
- (6) Conditional variance and law of total variance
- (7) Random number of sum of random variables


- Given the PDF of  $X$ , What is the PDF of  $Y = g(X)$ ?
- Wait! Didn't we cover this topic? No. We covered just  $\mathbb{E}[g(X)]$ .
- Examples:  $Y = X$ ,  $Y = X + 1$ ,  $Y = X^2$ , etc.
- What are easy or difficult cases?
- Easy cases
  - Discrete
  - Linear:  $Y = aX + b$

- Take all values of  $x$  such that  $g(x) = y$ , i.e.,

$$p_Y(y) = \mathbb{P}(g(X) = y) \\ = \sum_{x: g(x)=y} p_X(x)$$

$$p_Y(3) = p_X(2) + p_X(3) = 0.1 + 0.2 = 0.3$$

$$p_Y(4) = p_X(4) + p_X(5) = 0.3 + 0.4 = 0.7$$



L5(1)

June 2, 2021 5 / 46

$$\text{If } a > 0, \quad F_Y(y) = \mathbb{P}(aX + b \leq y) = \mathbb{P}(X \leq \frac{y-b}{a}) = F_X(\frac{y-b}{a})$$

$$\rightarrow f_Y(y) = \frac{1}{a} f_X\left(\frac{y-b}{a}\right)$$

$$\text{If } a < 0, \quad F_Y(y) = \mathbb{P}(aX + b \leq y) = \mathbb{P}(X \geq \frac{y-b}{a}) = 1 - F_X(\frac{y-b}{a})$$

$$\rightarrow f_Y(y) = -\frac{1}{a} f_X\left(\frac{y-b}{a}\right)$$

Therefore,

$$f_Y(y) = \frac{1}{|a|} f_X\left(\frac{y-b}{a}\right)$$

L5(1)

June 2, 2021 6 / 46

$$f_X(x) = \begin{cases} \lambda e^{-\lambda x}, & \text{if } x \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

$$f_Y(y) = \begin{cases} \frac{\lambda}{|a|} e^{-\lambda(y-b)/a}, & \text{if } (y-b)/a \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

- If  $b = 0$  and  $a > 0$ ,  $Y$  is exponential with parameter  $\frac{\lambda}{a}$ , but generally not.

L5(1)

June 2, 2021 7 / 46

- Remember? Linear transformation preserves normality. Time to prove.

$$\text{If } X \sim \mathcal{N}(\mu, \sigma^2), \text{ then for } a \neq 0 \text{ and } b, Y = aX + b \sim \mathcal{N}(a\mu + b, a^2\sigma^2).$$

- Proof.**

$$f_X(x) = \frac{1}{\sqrt{2\pi}} e^{-(x-\mu)^2/2\sigma^2}$$

$$f_Y(y) = \frac{1}{|a|} f_X\left(\frac{y-b}{a}\right) = \frac{1}{|a|} \frac{1}{\sqrt{2\pi}} \exp\left\{-\left(\frac{y-b}{a} - \mu\right)^2/2\sigma^2\right\} \\ = \frac{1}{\sqrt{2\pi}|a|\sigma} \exp\left\{-\frac{(y-b-a\mu)^2}{2a^2\sigma^2}\right\}$$

L5(1)

June 2, 2021 8 / 46

Step 1. Find the CDF of  $Y$ :

$$F_Y(y) = \mathbb{P}(Y \leq y) = \mathbb{P}(g(X) \leq y)$$

Step 2. Differentiate:  $f_Y(y) = \frac{dF_Y}{dy}(y)$

Ex1.  $Y = X^2$ .

$$F_Y(y) = \mathbb{P}(X^2 \leq y) = \mathbb{P}(-\sqrt{y} \leq X \leq \sqrt{y}) \\ = F_X(\sqrt{y}) - F_X(-\sqrt{y})$$

$$f_Y(y) = \frac{1}{2\sqrt{y}} f_X(\sqrt{y}) + \\ \frac{1}{2\sqrt{y}} f_X(-\sqrt{y}), \quad y \geq 0$$

Ex2.  $X \sim \mathcal{U}[0, 1]$ .  $Y = \sqrt{X}$ .

$$F_Y(y) = \mathbb{P}(\sqrt{X} \leq y) = \mathbb{P}(X \leq y^2) = y^2$$

$$f_Y(y) = 2y, \quad 0 \leq y \leq 1$$

Ex3.  $X \sim \mathcal{U}[0, 2]$ .  $Y = X^3$ .

$$F_Y(y) = \mathbb{P}(X^3 \leq y) = \mathbb{P}(X \leq \sqrt[3]{y}) = \frac{1}{2} y^{1/3}$$

$$f_Y(y) = \frac{1}{6} y^{-2/3}, \quad 0 \leq y \leq 8$$

When  $Y = g(X)$  is monotonic, a **general formula** can be drawn (see the textbook at pp 207)

Basically, follow two-step approach: (i) CDF and (ii) differentiate.

Ex1.  $X, Y \sim \mathcal{U}[0, 1]$ , and  $X \perp\!\!\!\perp Y$ .  $Z = \max(X, Y)$ .

\*  $\mathbb{P}(X \leq z) = \mathbb{P}(Y \leq z) = z$ ,  $z \in [0, 1]$ .

$$F_Z(z) = \mathbb{P}(\max(X, Y) \leq z) = \mathbb{P}(X \leq z, Y \leq z) \\ = \mathbb{P}(X \leq z) \mathbb{P}(Y \leq z) = z^2 \quad (\text{from } X \perp\!\!\!\perp Y)$$

$$f_Z(z) = \begin{cases} 2z, & \text{if } 0 \leq z \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

Basically, follow two step approach: (i) CDF and (ii) differentiate.


Ex2.  $X, Y \sim \mathcal{U}[0, 1]$ , and  $X \perp\!\!\!\perp Y$ .  $Z = Y/X$ .

VIDEO PAUSE

$$F_Z(z) = \mathbb{P}(Y/X \leq z) \\ = \begin{cases} z/2, & 0 \leq z \leq 1 \\ 1 - 1/2z, & z > 1 \\ 0, & \text{otherwise} \end{cases}$$

$$f_Z(z) = \begin{cases} 1/2, & 0 \leq z \leq 1 \\ 1/(2z^2), & z > 1 \\ 0, & \text{otherwise} \end{cases}$$

- Depending on the value of  $z$ , two cases need to be considered separately.



(Note) Sometimes, the problem is tricky, which requires careful case-by-case handling. :-)

(1) Derived distribution of  $Y = g(X)$  or  $Z = g(X, Y)$

(2) **Derived distribution of  $Z = X + Y$**

(3) Covariance: Degree of dependence between two rvs.

(4) Correlation coefficient

(5) Conditional expectation and law of iterative expectations

(6) Conditional variance and law of total variance

(7) Random number of sum of random variables

- Sum of two independent rvs
- A very basic case with many applications
- Assume that  $X, Y \in \mathbb{Z}$

$$p_Z(z) = \mathbb{P}(X + Y = z) = \sum_{\{(x,y): x+y=z\}} \mathbb{P}(X=x, Y=y) = \sum_x \mathbb{P}(X=x, Y=z-x)$$

$$= \sum_x \mathbb{P}(X=x) \mathbb{P}(Y=z-x) = \sum_x p_X(x) p_Y(z-x)$$

- $p_Z(z)$  is called **convolution** of the PMFs of  $X$  and  $Y$ .

L5(2)


June 2, 2021 13 / 46

- Convolution:  $p_Z(z) = \sum_x p_X(x) p_Y(z-x)$

- Interpretation for a given  $z$ :

- Flip (horizontally) the PMF of  $Y$  ( $p_Y(-x)$ )
- Put it underneath the PMF of  $X$
- Right-shift the flipped PMF by  $z$  ( $p_Y(-x+z)$ )

Example.  $z = 3$



L5(2)


June 2, 2021 14 / 46

- Same logic as the discrete case


$$f_Z(z) = \int_{-\infty}^{\infty} f_X(x) f_Y(z-x) dx$$

- Youtube animation for convolution: <https://www.youtube.com/watch?v=C1N55M1VD2o>

For a fixed  $z$ ,



- Example.  $X, Y \sim \mathcal{U}[0, 1]$  and  $X \perp\!\!\!\perp Y$ . What is the PDF of  $Z = X + Y$ ? Draw the PDF of  $Z$ .



L5(2)

June 2, 2021 15 / 46

L5(2)

June 2, 2021 16 / 46

<https://www.youtube.com/watch?v=MQm6ZP1F6ms>

- Very special, but useful case
  - $X$  and  $Y$  are **normal**.

#### Sum of two independent normal rvs

$X \sim \mathcal{N}(\mu_x, \sigma_x^2)$  and  $Y \sim \mathcal{N}(\mu_y, \sigma_y^2)$  Then,  $X + Y \sim \mathcal{N}(\mu_x + \mu_y, \sigma_x^2 + \sigma_y^2)$

- Why normal rvs are used to model the **sum of random noises**.
- **Extension**. The sum of **finitely many** independent normals is also normal.

$$\begin{aligned} f_Z(z) &= \int_{-\infty}^{\infty} f_X(x) f_Y(z-x) dx \\ &= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}\sigma_x} \exp\left\{-\frac{(x-\mu_x)^2}{2\sigma_x^2}\right\} \frac{1}{\sqrt{2\pi}\sigma_y} \exp\left\{-\frac{(z-x-\mu_y)^2}{2\sigma_y^2}\right\} dx \end{aligned}$$

- The details of integration is a little bit tedious. :-)

$$f_Z(z) = \frac{1}{\sqrt{2\pi(\sigma_x^2 + \sigma_y^2)}} \exp\left\{-\frac{(z - \mu_x - \mu_y)^2}{2(\sigma_x^2 + \sigma_y^2)}\right\}$$


- (1) Derived distribution of  $Y = g(X)$  or  $Z = g(X, Y)$
- (2) Derived distribution of  $Z = X + Y$
- (3) **Covariance: Degree of dependence between two rvs**
- (4) Correlation coefficient
- (5) Conditional expectation and law of iterative expectations
- (6) Conditional variance and law of total variance
- (7) Random number of sum of random variables

- Goal: Given two rvs  $X$  and  $Y$ , assign some number that quantifies the degree of their dependence.
  - feeling/weather, university ranking/annual salary,
- Requirements
  - R1.** Increases (resp. decreases) as they become more (resp. less) dependent. 0 when they are independent.
  - R2.** Shows the 'direction' of dependence by + and -
  - R3.** Always bounded by some numbers (i.e., dimensionless metric). For example,  $[-1, 1]$
- Good engineers: Good at making good metrics
  - Metric of how our society is economically polarized
  - Cybermetrics in MLB (Major League Baseball):  
<http://m.mlb.com/glossary/advanced-stats>

L5(3)

June 2, 2021 21 / 46

- Simple case:  $\mathbb{E}[X] = \mu_X = 0$  and  $\mathbb{E}[Y] = \mu_Y = 0$
- Dependent: Positive (If  $X \uparrow$ ,  $Y \uparrow$ ) or Negative (If  $X \uparrow$ ,  $Y \downarrow$ )
- What about  $\mathbb{E}[XY]$ ? Seems good.
  - $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y] = 0$  when  $X \perp\!\!\!\perp Y$
  - More data points (thus increases) when  $xy > 0$  (both positive or negative)
  - $|\mathbb{E}[XY]|$  also quantifies the **amount of spread**.

(Q) What about  $\mathbb{E}[X + Y]$ ?

- When they are positively dependent, but have negative values?

L5(3)

June 2, 2021 22 / 46

- **Solution:** Centering.  $X \rightarrow X - \mu_X$  and  $Y \rightarrow Y - \mu_Y$

## Covariance


$$\text{cov}(X, Y) = \mathbb{E}[(X - \mathbb{E}[X]) \cdot (Y - \mathbb{E}[Y])]$$

- After some algebra,  $\text{cov}(X, Y) = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y]$
- $X \perp\!\!\!\perp Y \implies \text{cov}(X, Y) = 0$
- $\text{cov}(X, Y) = 0 \implies X \perp\!\!\!\perp Y$ ? NO.
- When  $\text{cov}(X, Y) = 0$ , we say that  $X$  and  $Y$  are **uncorrelated**.

L5(3)

June 2, 2021 23 / 46

- $p_{X,Y}(1, 0) = p_{X,Y}(0, 1) = p_{X,Y}(-1, 0) = p_{X,Y}(0, -1) = 1/4$ .
- $\mathbb{E}[X] = \mathbb{E}[Y] = 0$ , and  $\mathbb{E}[XY] = 0$ . So,  $\text{cov}(X, Y) = 0$
- Are they independent? No, because if  $X = 1$ , then we should have  $Y = 0$ .



L5(3)

June 2, 2021 24 / 46

$$\text{cov}(X, X) = \text{var}(X)$$

$$\text{cov}(aX + b, Y) = \mathbb{E}[(aX + b)Y] - \mathbb{E}[aX + b]\mathbb{E}[Y] = a \cdot \text{cov}(X, Y)$$

$$\text{cov}(X, Y + Z) = \mathbb{E}[X(Y + Z)] - \mathbb{E}[X]\mathbb{E}[Y + Z] = \text{cov}(X, Y) + \text{cov}(X, Z)$$

$$\text{var}[X + Y] = \mathbb{E}[(X + Y)^2] - (\mathbb{E}[X + Y])^2 = \text{var}[X] + \text{var}[Y] + 2\text{cov}(X, Y)$$

$$\text{var}\left[\sum X_i\right] = \sum \text{var}[X_i] + \sum_{i \neq j} \text{cov}(X_i, X_j)$$

L5(3)

June 2, 2021 25 / 46

- $n$  people throw their hats in a box and then pick one at random
- $X$ : number of people with their own hat
- (Q)  $\text{var}[X]$
- Key step 1. Define a rv  $X_i = 1$  if  $i$  selects own hat and 0 otherwise. Then,  $X = \sum_{i=1}^n X_i$ .
- Key step 2. Are  $X_i$ s are independent?
- $X_i \sim \text{Bern}(1/n)$ . Thus,  $\mathbb{E}[X_i] = 1/n$  and  $\text{var}[X_i] = \frac{1}{n}(1 - \frac{1}{n})$

- For  $i \neq j$ ,

$$\begin{aligned} \text{cov}(X_i, X_j) &= \mathbb{E}[X_i X_j] - \mathbb{E}[X_i]\mathbb{E}[X_j] \\ &= \mathbb{P}(X_i = 1 \text{ and } X_j = 1) - \frac{1}{n^2} \\ &= \mathbb{P}(X_i = 1)\mathbb{P}(X_j = 1 | X_i = 1) - \frac{1}{n^2} \\ &= \frac{1}{n} \frac{1}{n-1} - \frac{1}{n^2} = \frac{1}{n^2(n-1)} \end{aligned}$$

$$\begin{aligned} \text{var}[X] &= \text{var}\left[\sum X_i\right] \\ &= \sum \text{var}[X_i] + \sum_{i \neq j} \text{cov}(X_i, X_j) \\ &= n \frac{1}{n} \left(1 - \frac{1}{n}\right) + n(n-1) \frac{1}{n^2(n-1)} = 1 \end{aligned}$$

L5(3)

June 2, 2021 26 / 46

- (1) Derived distribution of  $Y = g(X)$  or  $Z = g(X, Y)$
- (2) Derived distribution of  $Z = X + Y$
- (3) Covariance: Degree of dependence between two rvs
- (4) Correlation coefficient
- (5) Conditional expectation and law of iterative expectations
- (6) Conditional variance and law of total variance
- (7) Random number of sum of random variables

L5(4)

June 2, 2021 27 / 46

- Reqs. **R1** and **R2** are satisfied.
- **R3.** Always bounded by some numbers (dimensionless metric)
- How? Normalization, but by what?

## Correlation Coefficient

$$\rho(X, Y) = \mathbb{E}\left[\frac{(X - \mu_X)}{\sigma_X} \cdot \frac{(Y - \mu_Y)}{\sigma_Y}\right] = \frac{\text{cov}(X, Y)}{\sqrt{\text{var}[X]\text{var}[Y]}}$$

- Theorem.
  1.  $-1 \leq \rho \leq 1$  (proof at the next slide)
  2.  $|\rho| = 1 \Leftrightarrow X - \mu_X = c(Y - \mu_Y)$  for some constant  $c$  ( $c > 0$  when  $\rho = 1$  and  $c < 0$  when  $\rho = -1$ ). In other words, linear relation, meaning VERY related.

L5(4)

June 2, 2021 28 / 46

- **Cauchy-Schwarz inequality.** For any rvs  $X$  and  $Y$ ,  $(\mathbb{E}(XY))^2 \leq \mathbb{E}(X^2)\mathbb{E}(Y^2)$

- **Proof of  $-1 \leq \rho \leq 1$ :**

Let  $\tilde{X} = X - \mathbb{E}(X)$  and  $\tilde{Y} = Y - \mathbb{E}(Y)$ . Then,  $(\rho(X, Y))^2 = \frac{(\mathbb{E}[\tilde{X}\tilde{Y}])^2}{\mathbb{E}(\tilde{X}^2)\mathbb{E}(\tilde{Y}^2)} \leq 1$

- **Proof of CSI:** For any constant  $a$ ,

$$0 \leq \mathbb{E}[(X - aY)^2] = \mathbb{E}[X^2 - 2aXY + a^2Y^2] = \mathbb{E}(X^2) - 2a\mathbb{E}(XY) + a^2\mathbb{E}(Y^2)$$

Now, choose  $a = \frac{\mathbb{E}(XY)}{\mathbb{E}(Y^2)}$ . Then,

$$\mathbb{E}(X^2) - 2\frac{\mathbb{E}(XY)}{\mathbb{E}(Y^2)}\mathbb{E}(XY) + \frac{(\mathbb{E}[XY])^2}{(\mathbb{E}[Y^2])^2}\mathbb{E}(Y^2) = \mathbb{E}(X^2) - \frac{(\mathbb{E}[XY])^2}{\mathbb{E}(Y^2)} \geq 0$$

L5(4)

June 2, 2021 29 / 46

( $\Rightarrow$ ) Suppose that  $|\rho| = 1$ . In the proof of CSI,

$$\mathbb{E} \left[ \left( \tilde{X} - \frac{\mathbb{E}(\tilde{X}\tilde{Y})}{\mathbb{E}(\tilde{Y}^2)} \tilde{Y} \right)^2 \right] = \mathbb{E}(\tilde{X}^2) - \frac{(\mathbb{E}[\tilde{X}\tilde{Y}])^2}{\mathbb{E}(\tilde{Y}^2)} = \mathbb{E}(\tilde{X}^2)(1 - \rho^2) = 0$$

$$\tilde{X} - \frac{\mathbb{E}(\tilde{X}\tilde{Y})}{\mathbb{E}(\tilde{Y}^2)} \tilde{Y} = 0 \Leftrightarrow \tilde{X} = \frac{\mathbb{E}(\tilde{X}\tilde{Y})}{\mathbb{E}(\tilde{Y}^2)} \tilde{Y} = \rho \sqrt{\frac{\mathbb{E}(\tilde{X}^2)}{\mathbb{E}(\tilde{Y}^2)}} \tilde{Y}$$

( $\Leftarrow$ ) If  $\tilde{Y} = c\tilde{X}$ , then

$$\rho(X, Y) = \frac{\mathbb{E}(\tilde{X}c\tilde{X})}{\sqrt{\mathbb{E}[\tilde{X}^2]\mathbb{E}[(c\tilde{X})^2]}} = \frac{c}{|c|}$$

L5(4)

June 2, 2021 30 / 46

- (1) Derived distribution of  $Y = g(X)$  or  $Z = g(X, Y)$
- (2) Derived distribution of  $Z = X + Y$
- (3) Covariance: Degree of dependence between two rvs
- (4) Correlation coefficient
- (5) **Conditional expectation and law of iterative expectations**
- (6) Conditional variance and law of total variance
- (7) Random number of sum of random variables

L5(5)

June 2, 2021 31 / 46

- Consider a rv  $Y$ , such that

$$Y = \begin{cases} 0, & \text{w.p. } 1/4 \\ 1, & \text{w.p. } 1/4 \\ 2, & \text{w.p. } 1/2 \end{cases}$$

- If  $h(y) = y^2$ , then a new rv  $h(Y)$  is:

$$h(Y) = \begin{cases} 0, & \text{w.p. } 1/4 \\ 1, & \text{w.p. } 1/4 \\ 4, & \text{w.p. } 1/2 \end{cases}$$

- Consider other rv  $X$ , which, we assume, has:

$$g(y) = \mathbb{E}[X|Y=y] = \begin{cases} 3, & \text{if } y = 0 \\ 8, & \text{if } y = 1 \\ 9, & \text{if } y = 2 \end{cases}$$

- Then, a rv  $g(Y)$  is:

$$g(Y) = \begin{cases} 3, & \text{w.p. } 1/4 \\ 8, & \text{w.p. } 1/4 \\ 9, & \text{w.p. } 1/2 \end{cases}$$

- The rv  $g(Y)$  looks special, so let's give a fancy notation to it.

- What about?  $X_{\text{exp}}(Y)$ ,  $\mathbb{E}[X_Y]$ ,  $\mathbb{E}_X[Y]$ ?

L5(5)

June 2, 2021 32 / 46



## Conditional Expectation

A random variable  $g(Y) = \mathbb{E}[X|Y]$ , called **conditional expectation of  $X$  given  $Y$** , takes the value  $g(y) = \mathbb{E}[X|Y=y]$ , if  $Y$  happens to take the value  $y$ .

- A function of  $Y$
- A random variable
- Thus, having a distribution, expectation, variance, all the things that a random variable has.
- Often confusing because of the notation.

L5(5)

June 2, 2021 33 / 46

## Expectation of Conditional Expectation

$$\mathbb{E}[\mathbb{E}[X|Y]] = \mathbb{E}[X], \quad \text{Law of iterated expectations}$$

**Proof.**

$$\begin{aligned} \mathbb{E}[\mathbb{E}[X|Y]] &= \sum_y \mathbb{E}[X|Y=y] p_Y(y) \\ &= \mathbb{E}[X] \end{aligned}$$

L5(5)

June 2, 2021 34 / 46

- Stick of length  $l$
- Uniformly break at point  $Y$ , and break what is left uniformly at point  $X$ .
- $\mathbb{E}[X|Y=y] = y/2$
- $\mathbb{E}[X|Y] = Y/2$
- $\mathbb{E}[X] = \mathbb{E}[\mathbb{E}[X|Y]] = \mathbb{E}[Y/2] = \frac{1}{2} \frac{l}{2} = l/4$

- Forecasts on sales: calculating expected value, given any available information
- $X$  : February sales
- Forecast in the beg. of the year:  $\mathbb{E}[X]$
- End of Jan. new information  $Y = y$  (Jan. sales)  
Revised forecast:  $\mathbb{E}[X|Y=y]$   
Revised forecast  $\neq \mathbb{E}[X]$
- Law of iterated expectations  
 $\mathbb{E}[\text{revised forecast}] = \text{original one}$

L5(5)

June 2, 2021 35 / 46

- A class:  $n$  students, student  $i$ 's quiz score:  $x_i$
- Average quiz score:  $m = \frac{1}{n} \sum_{i=1}^n x_i$
- Students: partitioned into sections  $A_1, \dots, A_k$  and  $n_s$ : number of students in section  $s$
- average score in section  $s =$   
 $m_s = \frac{1}{n_s} \sum_{i \in A_s} x_i$
- whole average: (i) taking the average  $m_s$  of each section and (ii) forming a weighted average

$$\sum_{s=1}^k \frac{n_s}{n} m_s = \sum_{s=1}^k \frac{n_s}{n} \frac{1}{n_s} \sum_{i \in A_s} x_i = \frac{1}{n} \sum_{i=1}^n x_i = m$$

- Understanding from  $\mathbb{E}[\mathbb{E}[X|Y]] = \mathbb{E}[X]$
- $X$ : score of a randomly chosen student,  $Y$ : section of a student ( $\in \{1, \dots, k\}$ )

$$\begin{aligned} m &= \mathbb{E}(X) = \mathbb{E}[\mathbb{E}[X|Y]] \\ &= \sum_{s=1}^k \mathbb{E}(X|Y=s) \mathbb{P}(Y=s) \\ &= \sum_{s=1}^k \left( \frac{1}{n_s} \sum_{i \in A_s} x_i \right) \frac{n_s}{n} = \sum_{s=1}^k m_s \frac{n_s}{n} \end{aligned}$$

L5(5)

June 2, 2021 36 / 46

- (1) Derived distribution of  $Y = g(X)$  or  $Z = g(X, Y)$
- (2) Derived distribution of  $Z = X + Y$
- (3) Covariance: Degree of dependence between two rvs
- (4) Correlation coefficient
- (5) Conditional expectation and law of iterative expectations
- (6) **Conditional variance and law of total variance**
- (7) Random number of sum of random variables

L5(6)

June 2, 2021 37 / 46

$$\text{var}[X] = \mathbb{E}[(X - \mathbb{E}[X])^2]$$

$$g(y) = \text{var}[X|Y=y] = \mathbb{E}[(X - \mathbb{E}[X|Y=y])^2|Y=y]$$

$$g(Y) = \text{var}[X|Y] = \mathbb{E}[(X - \mathbb{E}[X|Y])^2|Y]$$

### Conditional Variance

A random variable  $g(Y) = \boxed{\text{var}[X|Y]}$  and called **conditional variance of  $X$  given  $Y$** , takes the value  $g(y) = \text{var}[X|Y=y]$ , if  $Y$  happens to take the value  $y$ .

- A function of  $Y$
- A random variable
- Thus, having a distribution, expectation, variance, all the things that a random variable has

L5(6)

June 2, 2021 38 / 46

	$\mathbb{E}[X Y]$	$\text{var}[X Y]$
Expectation	$\mathbb{E}[\mathbb{E}(X Y)]$	$\mathbb{E}[\text{var}(X Y)]$
Variance	$\text{var}[\mathbb{E}(X Y)]$	$\text{var}[\text{var}(X Y)]$

L5(6)

June 2, 2021 39 / 46

### Law of total variance (LTV)

$$\text{var}[X] = \mathbb{E}[\text{var}(X|Y)] + \text{var}[\mathbb{E}(X|Y)]$$

### Proof.

$$\text{var}(X|Y) = \mathbb{E}[X^2|Y] - (\mathbb{E}[X|Y])^2$$

$$\mathbb{E}[\text{var}(X|Y)] = \mathbb{E}[X^2] - \mathbb{E}[(\mathbb{E}[X|Y])^2] \quad (1)$$

$$\text{var}[\mathbb{E}(X|Y)] = \mathbb{E}[(\mathbb{E}[X|Y])^2] - (\mathbb{E}[\mathbb{E}(X|Y)])^2 = \mathbb{E}[(\mathbb{E}[X|Y])^2] - (\mathbb{E}[X])^2 \quad (2)$$

$$(1) + (2) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2 = \text{var}[X]$$

L5(6)

June 2, 2021 40 / 46

- Same setting as that in page 36
- $X$ : score of a randomly chosen student,  $Y$ : section of a student ( $\in \{1, \dots, k\}$ )
- Let's intuitively understand:  $\text{var}[X] = \mathbb{E}[\text{var}(X|Y)] + \text{var}[\mathbb{E}(X|Y)]$
- $\mathbb{E}[\text{var}(X|Y)] = \sum_{k=1}^s \mathbb{P}(Y = s) \text{var}(X|Y = s) = \sum_{k=1}^s \frac{n_s}{n} \text{var}(X|Y = s)$ 
  - Weighted average of the section variances
  - **average score variability within individual sections**
- $\text{var}[\mathbb{E}(X|Y)]$ : variability of the average of the different sections
  - $\mathbb{E}(X|Y = s)$ : average score in section  $s$
  - **variability between sections**

- Stick of length  $l$
- Uniformly break at point  $Y$ , and break what is left uniformly at point  $X$ .
- **Question.**  $\text{var}(X)$ ?
- LTV:  $\text{var}[X] = \mathbb{E}[\text{var}(X|Y)] + \text{var}[\mathbb{E}(X|Y)]$
- **Fact.** If a rv  $X \sim \mathcal{U}[0, \theta]$ , then  $\text{var}(X) = \frac{\theta^2}{12}$
- Since  $X \sim \mathcal{U}[0, Y]$ ,  $\text{var}(X|Y) = \frac{Y^2}{12} \rightarrow \mathbb{E}[\text{var}[X|Y]] = \frac{1}{12} \int_0^l \frac{1}{l} y^2 dy = \frac{l^2}{36}$
- $\mathbb{E}(X|Y) = Y/2 \rightarrow \text{var}[\mathbb{E}(X|Y)] = \frac{1}{4} \text{var}[Y] = \frac{1}{4} \frac{l^2}{12} = \frac{l^2}{48}$
- $\text{var}(X) = \frac{l^2}{36} + \frac{l^2}{48} = \frac{7l^2}{144}$

- (1) Derived distribution of  $Y = g(X)$  or  $Z = g(X, Y)$
- (2) Derived distribution of  $Z = X + Y$
- (3) Covariance: Degree of dependence between two rvs
- (4) Correlation coefficient
- (5) Conditional expectation and law of iterative expectations
- (6) Conditional variance and law of total variance
- (7) **Random number of sum of random variables**

- $N$ : number of stores visited (**random**)
- $X_i$ : money spent in store  $i$ , independent of other  $X_j$  and  $N$ ,  $X_i$ s are identically distributed with  $\mathbb{E}[X_i] = \mu$
- $Y = X_1 + X_2 + \dots + X_N$ . What are  $\mathbb{E}[Y]$  and  $\text{var}[Y]$ ?
- $\mathbb{E}[Y] = \mathbb{E}[\mathbb{E}[Y|N]] = \mathbb{E}[N\mathbb{E}[X_i]] = \mathbb{E}[N]\mathbb{E}[X_i] = \mu\mathbb{E}[N]$
- $\text{var}[Y] = \mathbb{E}[\text{var}(Y|N)] + \text{var}[\mathbb{E}(Y|N)] = \mathbb{E}[N\text{var}[X_i]] + \mu^2\text{var}[N]$   
 $\text{var}(\mathbb{E}[Y|N]) = \text{var}(N\mu) = \mu^2\text{var}[N]$   
 $\text{var}[Y|N] = N\text{var}[X_i]$   
 $\mathbb{E}[\text{var}(Y|N)] = \mathbb{E}[N\text{var}[X_i]] = \mathbb{E}[N]\text{var}[X_i]$

## Questions?

- 1) What are the key steps to get the derived distributions of  $Y = g(X)$  or  $Z = g(X, Y)$ ?
- 2) How can we compute the distribution of  $Z = X + Y$  when  $X$  and  $Y$  are independent?
- 3) What are covariance and correlation coefficient? Why do we need them?
- 4) Please explain the concepts of conditional expectation and conditional variance.