

Lecture 7: Random Processes, Part I

Yi, Yung (이웅)

EE210: Probability and Introductory Random Processes
KAIST EE

June 12, 2021

- (1) Introduction of Random Processes
- (2) Bernoulli Processes
- (3) Poisson Processes: Poisson RV and Basic Idea
- (4) Poisson Processes: Definition and Properties
- (5) Playing with Bernoulli and Poisson Processes

June 12, 2021 1 / 1


June 12, 2021 2 / 1

Roadmap


Things that evolve in time

- (1) Introduction of Random Processes
- (2) Bernoulli Processes
- (3) Poisson Processes: Poisson RV and Basic Idea
- (4) Poisson Processes: Definition and Properties
- (5) Playing with Bernoulli and Poisson Processes

- Many probabilistic experiments that **evolve in time**
 - Sequence of daily prices of a stock
 - Sequence of scores in football
 - Sequence of failure times of a machine
 - Sequence of traffic loads in Internet
- Random process is a mathematical model for it.



(a) Prices of a cryptocurrency



(b) Internet traffic traces

- A random process is a **sequence** of random variables indexed by **time**.
- Time: discrete or continuous (a modeling choice in most cases)
- Notation
 - $(X_t)_{t \in \mathcal{T}}$ or $(X(t))_{t \in \mathcal{T}}$, where $\mathcal{T} = \mathbb{R}$ (continuous) or $\mathcal{T} = \{0, 1, 2, \dots\}$ (discrete)
 - For the discrete case, we also often use $(X_n)_{n \in \mathbb{Z}_+}$.
 - We will use all of them, unless confusion arises.
- For a **fixed** time t , X_t (or $X(t)$) is a random variable.
- The values that X_t (or $X(t)$) can take: discrete or continuous

L8(1)

June 12, 2021 5 / 1

- **Example.** Discrete time RP. $\{X_1, X_2, X_3, \dots\}$.
 - X_i : # of Covid-19 infections at day i in South Korea, which is a random variable.
 - Then, $X_i : \Omega \mapsto \mathbb{R}$, where the sample space Ω is the set of all outcomes.
 - An outcome $\omega \in \Omega$ is a infinite sequence of infections
 - For example, $\omega_1 = (100, 150, 130, \dots)$, $\omega_2 = (200, 300, 400, \dots)$
 - $X_3(\omega_1) = 130$, $X_2(\omega_2) = 300$, $X_1(\omega_2) = 200$, etc.
- **Example.** Continuous time RP. $(X(t))_{t \in \mathbb{R}^+}$
 - $X(t)$: bitcoin price at time t , which is a random variable.
 - Then, $X(t) : \Omega \mapsto \mathbb{R}$.
 - An outcome $\omega \in \Omega$ is a trajectory of prices over $[0, \infty)$
 - $X(3.7, \omega_1) = 3409$, $X(2, \omega_2) = 5000$, $X(7.8, \omega_3) = 2800$, etc.

L8(1)

June 12, 2021 6 / 1


- **Question.** Already studied a sequence (or a collection) of rvs X_1, X_2, \dots, X_n . What's the difference?
 - Assume a discrete time random process for our discussion.
- **Physical difference:** **infinite** sequence of X_1, X_2, \dots .
 - Sample space? set of all outcomes?
 - an outcome: an infinite sequence of sample values x_1, x_2, \dots ,
- **Semantic difference:** Understand i in X_i as time. Also, interesting questions are asked from the random process point of view.
 - **Dependence:** How X_1, X_2, \dots are related to each other as a time series. Prediction of values in the future.
 - **Long-term behavior:** What is the fraction of times that a stock price is above 3000?
 - Other interesting questions, depending on the target random process

L8(1)

June 12, 2021 7 / 1

- Types of time and value

- (a) continuous time, continuous value
- (b) continuous time, discrete value
- (c) discrete time, continuous value
- (d) discrete time, discrete value



L8(1)

June 12, 2021 8 / 1

- The simplest RP
- discrete time
- $X_i \perp\!\!\!\perp \{X_{i-1}, X_{i-2}, \dots, X_1\}$
- **Bernoulli Process (BP)**
- “today” independent of “past”

Jacob Bernoulli (1654 - 1705),
Swiss



L8(1)

- The simplest RP
- Continuous time version of BP
- $[X(s)]_{s=0}^t \perp\!\!\!\perp [X(s)]_{s=t}^{t+a}$
- **Poisson Process (PP)**
- “today” independent of “past”


Simeon Denis Poisson (1781 -
1840), France



June 12, 2021 9 / 1

- One-step more general than BP/PP
- discrete time
- X_i depends on X_{i-1} , but $\perp\!\!\!\perp \{X_{i-2}, X_{i-3}, \dots, X_1\}$
- **Markov Chain (MC)**
- “today” depends only on “yesterday”

Andrey Markov (1856 - 1922),
Russia



- (1) Introduction of Random Processes
- (2) **Bernoulli Processes**
- (3) Poisson Processes: Poisson RV and Basic Idea
- (4) Poisson Processes: Definition and Properties
- (5) Playing with Bernoulli and Poisson Processes

L8(2)

June 12, 2021 10 / 1

- At each “minute”, we toss a coin with probability of head $0 < p < 1$.
 - Sequence of lottery wins/loses
 - Customers (each second) to a bank
 - Clicks (at each time slot) to server
- A sequence of **independent Bernoulli trials** X_1, X_2, \dots ,
 - We call index 1, 2, ... **time slots** (or simply slots)

0 0 1 0 0 0 0 1 0 1 1 0 0 →

- Discrete time, discrete value
- One of the simplest random processes
- A type of “arrival” process

Please pause the video and write down the questions that you want to ask about Bernoulli process.

VIDEO PAUSE

Q1.

Q2.

Q3.

Q4.

Q5.

L8(2)

June 12, 2021 11 / 1

L8(2)

June 12, 2021 12 / 1

(Q1) # of arrivals in the first n slots?

- $S_n = X_1 + X_2 + \dots + X_n$
- $S_n \sim \text{Binomial}(n, p)$
- $\mathbb{E}(S_n) = np$, $\text{var}(S_n) = np(1-p)$
- This will hold for any n consecutive slots.

(Q2) # of slots T_1 until the first arrival?

- $T_1 \sim \text{Geom}(p)$
- $\mathbb{E}(T_1) = 1/p$, $\text{var}(T_1) = \frac{1-p}{p^2}$

- T_1 is geometric? **Memoryless**
- Conditioned no first arrival at some time, what's the distribution of the remaining time until the first arrival?
- Still, geometric.
- But, more than that, as we will see. Independence across time slots leads to many useful properties, allowing the quick solution of many problems.

Independence across slots \implies the fresh-start anytime when I look at the process?

(Q3) $U = X_1 + X_2 \perp\!\!\!\perp V = X_5 + X_6$?

- Yes
- Because X_i s are independent

(Q4) After time $n = 6$, I start to look at the process $(X_n)_{n=6}^\infty$?

- $(X_1, \dots, X_5) \perp\!\!\!\perp (X_n)_{n=6}^\infty$
- **Fresh-start** after a deterministic time n (doesn't matter what happened until $n = 5$).
- If you watch the on-going Bernoulli process(p) from some time n , you still see the same Bernoulli process(p).


(Q5) The process $(X_N, X_{N+1}, X_{N+2}, \dots)$? Fresh-start even after **random** N ?

- This means that the time I start to look at the process is a random variable.
- Examples of N

E1. Time of 3rd arrival

E2. First time when 3 consecutive arrivals have been observed

E3. Time just before 3 consecutive arrivals




- Difference of N from n
 - The time when I watch the on-going Bernoulli process is **random**.
 - N is a random variable, i.e., $N : \Omega \mapsto \mathbb{R}$. What is Ω ?
- Do we experience the fresh-start for any N ? **E1, E2, and E3?**

(Q5) The process $(X_N, X_{N+1}, X_{N+2}, \dots)$? Fresh-start even after random N ?

E1. Time of 3rd arrival

E2. First time when 3 consecutive arrivals have been observed

E3. Time just before 3 consecutive arrivals



- E1.** When I watch the process, N has been already determined. **Yes**
- E2.** Same as **E1**. **Yes**
- E3.** Need the future knowledge. '111' does not become random. **No**

- The question of $N = n$? can be answered just from the knowledge about X_1, X_2, \dots, X_n ? Then, Yes! (see pp. 301 for more formal description)


- In probability theory, a random time N is said to be a **stopping time**, if the question of " $N = n$?" can be answered only from the present and the past knowledge of X_1, X_2, \dots, X_n .
- https://en.wikipedia.org/wiki/Stopping_time
- Fresh-start after N in Bernoulli process? **Yes, if N is a stopping time.**
- Please think about two examples of stopping time and not.
 - Yes.** Time when 10 consecutive arrivals have been observed
 - No.** Time of 2nd arrival in 10 consecutive arrivals

VIDEO PAUSE

L8(2)

June 12, 2021 17 / 1


- Regard an arrival as a server being busy (just for our easy understanding)
- First busy period B_1** : starts with the first busy slot and ends just before the first subsequent idle slot



- (Q6)** B_1 is a random variable. Distribution of B_1 ?
- N : time of the first busy slot. N is a stopping time?
 - Yes.** Thus, **fresh-start after N** . Because we can answer the question of $N = n$?, just using X_1, X_2, \dots, X_n .
- B_1 is geometric with parameter $(1 - p)$

L8(2)

June 12, 2021 18 / 1



- Question.** What about the second busy period B_2 ?
- N : time of the first busy slot of the second busy period. N is a stopping time?
- Yes.** Thus, fresh-start after N .
- Then, B_1 and B_2 are **identically distributed** as $\text{Geom}(1 - p)$.
- B_3, B_4, \dots ?


L8(2)

June 12, 2021 19 / 1

- Time of the first arrival $Y_1 \sim \text{Geom}(p)$
- (Q7)** Time of the k -th arrival Y_k ?

VIDEO PAUSE

- $T_k = Y_k - Y_{k-1}$: k -th inter-arrival ($k \geq 2, T_1 = Y_1$)
- $Y_k = T_1 + T_2 + \dots + T_k$.



- After each T_k , the fresh-start occurs.
- $\{T_i\}$ are i.i.d. and $\sim \text{Geom}(p)$
- We know Y_k 's expectation and variance: $\mathbb{E}[Y_k] = \frac{k}{p}$, $\text{var}[Y_k] = \frac{k(1-p)}{p^2}$, but **its distribution?**

L8(2)

June 12, 2021 20 / 1

- $Y_k = T_1 + T_2 + \dots + T_k$.
- $\{T_i\}$ are i.i.d. and $\sim \text{Geom}(p)$

$$\begin{aligned}\mathbb{P}(Y_k = t) &= \mathbb{P}(X_k = 1 \text{ and } k-1 \text{ arrivals during the first } t-1 \text{ slots}) \\ &= \mathbb{P}(X_k = 1) \cdot \mathbb{P}(k-1 \text{ arrivals during the first } t-1 \text{ slots}) \\ &= p \times \binom{t-1}{k-1} p^{k-1} (1-p)^{t-k} = \binom{t-1}{k-1} p^k (1-p)^{t-k}, \quad t = k, k+1, \dots\end{aligned}$$

L8(2)

June 12, 2021 21 / 1

- In the sequence of Bernoulli trials, the time Y_k of k -th success
- PMF of Y_k

$$\mathbb{P}(Y_k = t) = \begin{cases} \binom{t-1}{k-1} p^k (1-p)^{t-k}, & \text{if } t = k, k+1, \dots \\ 0, & \text{if } t = 1, 2, \dots, k-1 \end{cases}$$

- $\text{Pascal}(1, p) = \text{Geom}(p)$

L8(2)

June 12, 2021 22 / 1

- (1) Introduction of Random Processes
- (2) Bernoulli Processes
- (3) Poisson Processes: Poisson RV and Basic Idea
- (4) Poisson Processes: Definition and Properties
- (5) Playing with Bernoulli and Poisson Processes

L8(3)

June 12, 2021 23 / 1

- A random variable $S \sim \text{Bin}(n, p)$: Models the number of successes in a given number n of independent trials with success probability p .

$$p_S(k) = \frac{n!}{(n-k)!k!} p^k (1-p)^{n-k}$$

- Our interest: very large n and very small p , such that $np = \lambda$, i.e., $\lim_{n \rightarrow \infty} p_S(k)$?

$$\begin{aligned}p_S(k) &= \frac{n(n-1) \cdots (n-k+1)}{k!} \cdot \frac{\lambda^k}{n^k} \cdot \left(1 - \frac{\lambda}{n}\right)^{n-k} \\ &= \frac{n}{n} \cdot \frac{(n-1)}{n} \cdots \frac{(n-k+1)}{n} \cdot \frac{\lambda^k}{k!} \cdot \left(1 - \frac{\lambda}{n}\right)^n \cdot \left(1 - \frac{\lambda}{n}\right)^{-k} \xrightarrow{n \rightarrow \infty} e^{-\lambda} \frac{\lambda^k}{k!}\end{aligned}$$

L8(3)

June 12, 2021 24 / 1

- A Poisson random variable Z with parameter λ takes nonnegative integer values, whose PMF is:

$$p_Z(k) = e^{-\lambda} \frac{\lambda^k}{k!}, \quad k = 0, 1, \dots$$

- Infinitely many slots (n) with the infinitely small slot duration (thus infinitely small success probability $p = \lambda/n$)
- $\mathbb{E}(Z) = \lambda$ (because $\lambda = np$ is the mean of binomial rv)
- $\text{var}(Z) = \lambda$ (because $np(1-p)$ is the variance of binomial rv)

L8(3)

June 12, 2021 25 / 1

- A packet consisting of a string of n symbols is transmitted over a noisy channel.
- Each symbol: erroneous transmission with probability of 0.0001, independent of other symbols. Incorrect transmission is when at least one symbol is in error.
- Question.** How small should n be in order for the probability of incorrect transmission to be less than 0.001?
- p is very small, and n is reasonably large \rightarrow Poisson approximation
- Prob. of incorrect transmission =
 $1 - \mathbb{P}(\text{no symbol error}) = 1 - e^{-\lambda} = 1 - e^{-0.0001n} < 0.0001$
- $n < \frac{-\ln 0.999}{0.0001} = 10.005$

L8(3)


June 12, 2021 26 / 1

- Remind.** Geometric vs. Exponential L4(3)
 - Two rvs with memoryless property
 - continuous system = discrete system with infinitely many slots whose duration is infinitely small.
- Independence between what happens in a different time region
- Memoryless and fresh-start property
- Choose one of discrete or continuous versions for our modeling convenience.
- Question.** How do we design the continuous analog of Bernoulli process?
 - Key idea: Making it as a limit of a sequence of Bernoulli processes
- Need a “modeling sense” to make this possible. It’s a good practice for engineers!
- VIDEO PAUSE

L8(3)

June 12, 2021 27 / 1


- Continuous twin
 - Key point: Understand the number of arrivals over a given interval $[0, \tau]$.
 - Assume that it has some arrival rate λ (# of arrivals/unit time).
 - We know how to handle Bernoulli process with discrete time slots.
- Divide $[0, \tau]$ into slots whose length = δ . Then, $n = \#$ of slots = $\frac{\tau}{\delta}$.



- What’s the limit as $\delta \rightarrow 0$ (equivalently, $n \rightarrow \infty$)

L8(3)

June 12, 2021 28 / 1




- Now, our design idea: during one time slot of length δ ,

$\mathbb{P}(1 \text{ arrival}) \propto \text{arrival rate and slot length}$
 $\mathbb{P}(\geq 2 \text{ arrivals}) \propto \text{something, but very small}$
 for small slot length
 $\mathbb{P}(0 \text{ arrival}) = 1 - \mathbb{P}(1 \text{ arrival or } \geq 2 \text{ arrivals})$

$$\begin{aligned}
 \mathbb{P}(1 \text{ arrival}) &= \lambda\delta + o(\delta) \\
 \mathbb{P}(\geq 2 \text{ arrivals}) &= o(\delta) \\
 \mathbb{P}(0 \text{ arrival}) &= 1 - \lambda\delta + o(\delta)
 \end{aligned}$$

- $o(\delta)$: some function that goes to zero faster than δ .
 - Thus, for very small δ , $o(\delta)$ becomes negligible, compared to δ .
 - Example: $o(\delta) = \delta^\alpha$, where any $\alpha > 1$



- Our interest: probability of k arrivals over $[0, \tau]$
- Given “small” δ , # of arrivals $\sim \text{Bin}(n, p)$, where $n = \tau/\delta$ and $p = \lambda\delta$
- As $\delta \rightarrow 0$, $np = \tau/\delta \times \lambda\delta = \lambda\tau$.
- # of arrivals over $[0, \tau]$, $\sim \text{Poisson}(\lambda\tau)$
- This is a continuous twin process of Bernous process, which we call **Poisson process**.

- (1) Introduction of Random Processes
- (2) Bernoulli Processes
- (3) Poisson Processes: Poisson RV and Basic Idea
- (4) **Poisson Processes: Definition and Properties**
- (5) Playing with Bernoulli and Poisson Processes

An arrival process is called a **Poisson process** with rate λ , if the following are satisfied:

- (Independence)** Let N_τ be the number of arrivals over the interval $[0, \tau]$. For any $\tau_1, \tau_2 > 0$, $N_{s+\tau_1} - N_s$ is independent of $N_{t+\tau_2} - N_t$, if $t > s + \tau_1$.
 - The number of arrivals over two disjoint intervals are independent.
- (Time homogeneity)** For any s , the distribution of $N_{s+\tau} - N_s$ is equal to that of N_τ .
 - N_τ becomes the number of arrivals over any interval of length τ .
- (Small interval probability)** Let $\mathbb{P}(k, \tau) = \mathbb{P}(N_\tau = k)$, which satisfy:

$$\begin{aligned}
 \mathbb{P}(0, \tau) &= 1 - \lambda\tau + o(\tau) \\
 \mathbb{P}(1, \tau) &= \lambda\tau + o_1(\tau) \\
 \mathbb{P}(k, \tau) &= o_k(\tau) \quad \text{for } k = 2, 3, \dots, \quad \text{where } \lim_{\tau \rightarrow 0} \frac{o(\tau)}{\tau} = 0, \quad \lim_{\tau \rightarrow 0} \frac{o_k(\tau)}{\tau} = 0
 \end{aligned}$$

An arrival process is called a **Poisson process** with rate λ , if the following are satisfied:

- **(Independence)** Let N_τ be the number of arrivals over the interval $[0, \tau]$. For any $\tau_1, \tau_2 > 0$, $N_{s+\tau_1} - N_s$ is independent of $N_{t+\tau_2} - N_t$, if $t > s + \tau_1$.
 - The number of arrivals over two disjoint intervals are independent.
- **(Time homogeneity)** For any s , the distribution of $N_{s+\tau} - N_s$ is equal to that of N_τ .
 - N_τ becomes the number of arrivals over any interval of length τ .
- **(Distribution of N_τ)** N_τ is the Poisson rv with parameter $\lambda\tau$, i.e., if we let $\mathbb{P}(k, \tau) = \mathbb{P}(N_\tau = k)$, we have:

$$\mathbb{P}(k, \tau) = e^{-\lambda\tau} \frac{(\lambda\tau)^k}{k!}, \quad k = 0, 1, 2, \dots$$

L8(4)

June 12, 2021 33 / 1

(Q1) Number of arrivals of **any interval** with length $\tau \sim \text{Poisson}(\lambda\tau)$, i.e.,

$$\mathbb{P}(k, \tau) = e^{-\lambda\tau} \frac{(\lambda\tau)^k}{k!}, \quad k = 0, 1, 2, \dots$$

$$\mathbb{E}(N_\tau) = \lambda\tau \text{ and } \text{var}(N_\tau) = \lambda\tau$$

(Q2) Time of first arrival T

$$F_T(t) = \mathbb{P}(T \leq t) = 1 - \mathbb{P}(T > t) = 1 - P(0, t) = 1 - e^{-\lambda t}$$

$$f_T(t) = \frac{dF_T(t)}{dt} = \lambda e^{-\lambda t}, \quad t \geq 0.$$

- $T \sim \text{Exp}(\lambda)$. Thus, $\mathbb{E}(T) = 1/\lambda$ and $\text{var}(T) = 1/\lambda^2$
 - Continuous twin of geometric rv in Bernoulli process
 - Memoryless

L8(4)

June 12, 2021 34 / 1

- Receive emails according to a Poisson process at rate $\lambda = 5$ messages/hour
- Mean and variance of mails received during a day
 - $5 \cdot 24 = 120$
- $\mathbb{P}[\text{one new message in the next hour}]$
 - $\mathbb{P}(1, 1) = \frac{(5 \cdot 1)^1 e^{-5 \cdot 1}}{1!} = 5e^{-5}$
- $\mathbb{P}[\text{exactly two msgs during each of the next three hours}]$
 - $\left(\frac{5^2 e^{-5}}{2!}\right)^3$

L8(4)

June 12, 2021 35 / 1

- **Remind.** Similar property for Bernoulli processes, but here no time slots.
- **Fresh-start at deterministic time t :** Start watching at time t , then you see the Poisson process, independent of what has happened in the past.
- **Fresh-start at random time T :** Similarly, it holds.
 - For example, when you start watching at random time T_1 (time of first arrival).
 - Generally, it holds when T is a **stopping time**.

(Q3) The k -th arrival time Y_k ?

- k -th inter-arrival time $T_k = Y_k - Y_{k-1}$, $k \geq 2$, and $T_1 = Y_1$.
- $Y_k = T_1 + T_2 + \dots + T_k$ is sum of i.i.d. exponential rvs.
- $\mathbb{E}[Y_k] = k/\lambda$ and $\text{var}[Y_k] = k/\lambda^2$, but what is the distribution of Y_k ?

L8(4)

June 12, 2021 36 / 1

- For a given δ , $\delta \cdot f_{Y_k}(y)$: prob. of k -th arrival over $[y, y + \delta]$.
- When δ is small, only one arrival occurs. Thus,

$$\delta \cdot f_{Y_k}(y) = \mathbb{P}(\text{an arrival over } [y, y + \delta]) \times \mathbb{P}(k-1 \text{ arrivals before } y)$$

$$\approx \lambda \delta \times \mathbb{P}(k-1, y) = \lambda \delta \times \frac{\lambda^{k-1} y^{k-1} e^{-\lambda y}}{(k-1)!}$$


$$f_{Y_k}(y) = \frac{\lambda^k y^{k-1} e^{-\lambda y}}{(k-1)!}, \quad y \geq 0.$$

- This is called **Erlang** rv.

An Erlang random variable Z with parameter (k, λ) has the following pdf:

$$f_Z(z) = \frac{\lambda^k z^{k-1} e^{-\lambda z}}{(k-1)!}, \quad z \geq 0$$

$$- n = \tau/\delta, \quad p = \lambda\delta, \quad np = \lambda\tau$$



	Bernoulli process	Poisson process
time of arrival	Discrete	Continuous
PMF of # of arrivals	Binomial	Poisson
Interarrival time	Geometric	Exponential
Time of k -th arrival	Pascal	Erlang
Arrival rate	p /per slot	λ /unit time

Example: Poisson Fishing (Problem 10, page 329)

Roadmap

- Catching fish: Poisson process $\lambda = 0.6$ /hour.
- Fish for 2 hours. Within 2 hours, if he catches at least one fish, then he stops at 2 hours. Otherwise, he fishes until one fish is caught.

(Q1) $\mathbb{P}(\text{fishing time} > 2 \text{ hours})$

Method 1: $\mathbb{P}(0, 2)$

Method 2: $\mathbb{P}(T_1 > 2)$

(Q2) $\mathbb{P}(2 < \text{fishing time} < 5)$

Method 1: $\mathbb{P}(0, 2)(1 - \mathbb{P}(0, 3))$

Method 2: $\mathbb{P}(2 < T_1 < 5)$

(Q3) $\mathbb{P}(\text{Catch at least two fish})$

Method 1: $\sum_{k=2}^{\infty} = 1 - \mathbb{P}(0, 2) - \mathbb{P}(1, 2)$

Method 2: $\mathbb{P}(Y_2 \leq 2)$

(Q4) $\mathbb{E}[\text{future fi. time} | \text{already fished for 3h}]$

Fresh-start. So,

$$\mathbb{E}[\exp(\lambda)] = 1/\lambda = 1/0.6$$

(Q5) $\mathbb{E}[F = \text{total fishing time}]$

$$2 + \mathbb{E}[F - 2] = 2 + \mathbb{P}(F = 2) \cdot 0 +$$

$$\mathbb{P}(F > 2) \cdot \mathbb{E}[F - 2 | F > 2]$$

$$= 2 + \mathbb{P}(0, 2) \cdot \frac{1}{\lambda}$$

(Q6) $\mathbb{E}[\text{number of fish}] = \lambda \cdot 2 + \mathbb{P}(0, 2) \cdot 1$

(1) Introduction of Random Processes

(2) Bernoulli Processes

(3) Poisson Processes: Poisson RV and Basic Idea

(4) Poisson Processes: Definition and Properties

(5) **Playing with Bernoulli and Poisson Processes**

Alternative Description of the Bernoulli Process

1. Start with a sequence of independent geometric random variables T_1, T_2, \dots , with common parameter p , and let these stand for the interarrival times.
2. Record a success (or arrival) at times $T_1, T_1 + T_2, T_1 + T_2 + T_3$, etc.

Alternative Description of the Poisson Process


1. Start with a sequence of independent exponential random variables T_1, T_2, \dots , with common parameter λ , and let these stand for the interarrival times.
2. Record an arrival at times $T_1, T_1 + T_2, T_1 + T_2 + T_3$, etc.

- It has been observed that after a rainy day, the number of days until the next rain $\sim \text{Geom}(p)$, independent of the past.
- **Question.** What is the probability that it rains on both the 5th and the 8th day of the month?
- **Approach 1:** Handling this problem directly with the geometric PMFs is very tedious and complex.
- **Approach 2:** Rainy days is a Bernoulli process with arrival probability p .
- Thus, the answer is p^2 .

- **Question.** How to make software codes of Bernoulli process with p and Poisson process with λ
- Inter-arrival times are very handy.
- **Bernoulli process with p :** Obtain a sequence of random values following the **geometric distribution** with parameter p .
- **Poisson process with λ :** Obtain a sequence of random values following the **exponential distribution** with parameter λ .

- Bernoulli random variable: **Bern(p)**
- Bernoulli process: **BP(p)**
- Poisson random variable: **Poisson(λ)**
- Poisson process: **PP(λ)**


- Split $BP(p)$ into two processes. Whenever there is an arrival, keep it w.p. q and discard it w.p. $(1 - q)$.
- Split decisions are independent of arrivals.
- Question.** What are the two split processes?
- $BP(pq)$ and $BP(p(1 - q))$. Why?
- Are they independent? **No.**



L8(5)

June 12, 2021 45 / 1

- Merge $BP(p)$ and $BP(q)$ into one process.
- Collided arrival is regarded just one arrival in the merged process
- Probability of having at least one arrival: $1 - (1 - p)(1 - q) = p + q - pq$
- Merged process: $BP(p + q - pq)$
- $\mathbb{P}(\text{arrival from proc. 1} \mid \text{arrival in the merged proc.}) = \frac{p}{p+q-pq}$



L8(5)

June 12, 2021 46 / 1

- Split a Poisson process $PP(\lambda)$ into two processes by keeping each arrival w.p. p and discarding it w.p. $(1 - p)$
- Question.** What are the split processes?
- Let's focus on the process that we keep
- Independence and time-homogeneity? **Yes**
- Small interval probability over δ -interval
 - $\mathbb{P}(1 \text{ arrival}) = p\lambda\delta + p \cdot o(\delta) = p\lambda\delta + o(\delta)$
 - $\mathbb{P}(2 \text{ arrivals}) = p \cdot o(\delta) = o(\delta)$
 - $\mathbb{P}(0 \text{ arrival}) = 1 - p\lambda\delta - p \cdot o(\delta) = 1 - p\lambda\delta + o(\delta)$
- $PP(\lambda p)$ and $PP(\lambda(1 - p))$

L8(5)

June 12, 2021 47 / 1

- Merge from $PP(\lambda_1)$ and $PP(\lambda_2)$
- Independence and time-homogeneity? **Yes**
- Small interval probability over δ -interval (ignoring $o(\delta)$ for small δ)
 - $\mathbb{P}(0 \text{ arrival}) \approx (1 - \lambda_1\delta)(1 - \lambda_2\delta) \approx 1 - (\lambda_1 + \lambda_2)\delta$
 - $\mathbb{P}(1 \text{ arrival}) \approx (\lambda_1\delta)(1 - \lambda_2\delta) + \lambda_2\delta(1 - \lambda_1\delta) \approx (\lambda_1 + \lambda_2)\delta$
- Merged process: $PP(\lambda_1 + \lambda_2)$

L8(5)

June 12, 2021 48 / 1

- **Red**: $PP(\lambda_1)$ and **Blue**: $PP(\lambda_2)$
- $\mathbb{P}(\text{from red} \mid \text{arrival at time } t \text{ in the merged proc.})? \frac{\lambda_1 \delta}{\lambda_1 \delta + \lambda_2 \delta} = \frac{\lambda_1}{\lambda_1 + \lambda_2}$
- Consider an event $A_k = \{k\text{-th arrival in the merged proc. is red}\}$
 - $\mathbb{P}(A_k)? \frac{\lambda_1}{\lambda_1 + \lambda_2}$
 - A_1, A_2, \dots are independent: origins (red or blue) of arrivals in the merged proc. are independent
- $\mathbb{P}(k \text{ out of first 10 arrivals are red})? \binom{10}{k} \left(\frac{\lambda_1}{\lambda_1 + \lambda_2} \right)^k \left(\frac{\lambda_2}{\lambda_1 + \lambda_2} \right)^{10-k}$

L8(5)

June 12, 2021 49 / 1

1. Competing exponentials
2. Sum of independent Poisson rvs
3. Poisson arrivals during and Exponential interval

L8(5)

June 12, 2021 50 / 1

- Two independent light bulbs have life times $T_a \sim \text{Exp}(\lambda_a)$ and $T_b \sim \text{Exp}(\lambda_b)$.
- (Q) Distribution of $Z = \min\{T_a, T_b\}$, the first time when a bulb burns out?
 - **Approach 1**
 - $\mathbb{P}(Z \geq z) = \mathbb{P}(T_a \geq z)\mathbb{P}(T_b \geq z) = e^{-\lambda_a z} e^{-\lambda_b z} = e^{-(\lambda_a + \lambda_b)z}$
 - Thus, $Z \sim \text{Exp}(\lambda_a + \lambda_b)$
 - **Approach 2**
 - T_a and T_b are the first arrival times of two Poisson processes of λ_a and λ_b , respectively.
 - Z is the first arrival time of merged Poisson process $(\lambda_a + \lambda_b)$.
 - Thus, $Z \sim \text{Exp}(\lambda_a + \lambda_b)$

L8(5)

June 12, 2021 51 / 1

- Three independent light bulbs have life time $T \sim \text{Exp}(\lambda)$.

(Q) $\mathbb{E}[\text{time until the last bulb burns out}]?$

- Understanding from the merged Poisson process
 - T_1 : time until the first burn-out, T_2 : time until the second burn-out, T_3 : time until the third burn-out
 - $\mathbb{E}(T_1 + T_2 + T_3)?$
 - $PP(3\lambda) \xrightarrow{\text{1st burn out}} PP(2\lambda) \xrightarrow{\text{2nd burn out}} PP(\lambda)$
 - $T_1 \sim \text{Exp}(3\lambda)$, $T_2 \sim \text{Exp}(2\lambda)$, $T_3 \sim \text{Exp}(\lambda)$

$$\mathbb{E}[T_1 + T_2 + T_3] = \frac{1}{3\lambda} + \frac{1}{2\lambda} + \frac{1}{\lambda}$$

L8(5)

June 12, 2021 52 / 1

- Two independent rvs X and Y , where $X \sim \text{Poisson}(\mu)$ and $Y \sim \text{Poisson}(\nu)$.
- **Question.** Distribution of $X + Y$? Complex convolution, but any other easy way?
- Poisson process perspective
 - X : number of arrivals of PP(1) over a time interval of length μ
 - Y : number of arrivals of PP(1) over a time interval of length ν
 - Two intervals do not overlap and located in a consecutive manner $\implies X \perp\!\!\!\perp Y$
- Distribution of $X + Y$: the number of arrivals of PP(1) over a time interval of length $\mu + \nu$
- Thus, $X + Y \sim \text{Poisson}(\mu + \nu)$

L8(5)

June 12, 2021 53 / 1

- Problem 24, pp. 335
- Consider PP(λ) and an independent rv $T \sim \text{Exp}(\nu)$
- **Question.** Distribution of N_T ?

- **Approach 1:** Total probability theorem

$$\mathbb{P}(N_T = k) = \int_0^\infty \mathbb{P}(N_T = k | T = \tau) f_T(\tau) d\tau = \int_0^\infty \mathbb{P}(N_\tau = k) f_T(\tau) d\tau$$

- Very tedious and not very intuitive.

L8(5)

June 12, 2021 54 / 1

- Consider PP(λ) and an independent rv $T \sim \text{Exp}(\nu)$
- Consider another PP(ν), and let's view T as the first arrival time in PP(ν).
- Now, consider the merged process of PP(λ) and PP(ν).
 - $\mathbb{P}[\text{from PP}(\lambda) | \text{arrival}] = \frac{\lambda}{\lambda + \nu}$ and $\mathbb{P}[\text{from PP}(\nu) | \text{arrival}] = \frac{\nu}{\lambda + \nu}$
- K : number of total arrivals until we get the first arrival from PP(ν).
 - Then, $K \sim \text{Geom}(\frac{\nu}{\lambda + \nu})$.
- Let L be the number of arrivals from PP(λ) until we get the first arrival from PP(ν).

$$p_L(l) = p_K(l+1) = \left(\frac{\nu}{\lambda + \nu}\right) \left(\frac{\lambda}{\lambda + \nu}\right)^l, \quad l = 0, 1, \dots$$

L8(5)

June 12, 2021 55 / 1

- Consider $S_n \sim \text{Binomial}(n, p)$. Then, $S_n = \sum_{i=1}^n X_i$, where $X_i \sim \text{Bern}(p)$.
- **Poisson approximation.** Poisson(np) is a good approximation of S_n
 - Holds when $np = \lambda$ ($n \rightarrow \infty, p \rightarrow 0$), i.e., X_i 's behavior changes over n .
- **Normal approximation.** $\sigma\sqrt{n}Z + n\mu$ is a good approximation of S_n
 - Holds for a fixed p ($n \rightarrow \infty$), i.e., X_i 's behavior does not change over n .
- In practice, an actual numbers of n and p are given, so which approximation is good under what situation?
 - $p = 1/100, n = 100$: $np = 1$, very asymmetric X_i , small $p \implies$ **Poisson**
 - $p = 1/3, n = 100$: large, reasonably symmetric p , at least moderate $n \implies$ **Normal**
 - $p = 1/100, n = 10,000$: small p , but large $n \implies$ **Both Poisson and Normal**

L8(5)

June 12, 2021 56 / 1

Questions?

1)