

Lecture 4: Random Variable, Part II

Yi, Yung (이윤)

EE210: Probability and Introductory Random Processes
KAIST EE

MONTH DAY, 2021

- Continuous Random Variable
- PDF (Probability Density Function)
- CDF (Cumulative Distribution Function)
- Exponential and Normal Distribution
- Joint PDF, Conditional PDF
- Bayes' rule for continuous and even mixed cases

- Famous discrete random variables used in the community
 - Bernoulli, Uniform, Binomial, Geometric, Poisson, etc.
- Summarizing a random variable: Expectation and Variance
- Functions of a single random variable, Functions of multiple random variables
- Conditioning for random variables, Independence for random variables
- Continuous random variables
 - Normal, Uniform, Exponential, etc.
- Bayes' rule for random variables



Continuous RV and Probability Density Function

- Many cases when random variable have “continuous values”, e.g., velocity of a car



Continuous RV and Probability Density Function

- Many cases when random variable have “continuous values”, e.g., velocity of a car

Continuous Random Variable

A rv X is **continuous** if \exists a function f_X , called **probability density function (PDF)**, s.t.

$$\mathbb{P}(X \in B) = \int_B f_X(x) dx$$

Continuous RV and Probability Density Function

- Many cases when random variable have “continuous values”, e.g., velocity of a car

Continuous Random Variable

A rv X is **continuous** if \exists a function f_X , called **probability density function (PDF)**, s.t.

$$\mathbb{P}(X \in B) = \int_B f_X(x) dx$$

- All of the concepts and methods (expectation, PMFs, and conditioning) for discrete rvs have continuous counterparts

Continuous RV and Probability Density Function


- Many cases when random variable have “continuous values”, e.g., velocity of a car

Continuous Random Variable

A rv X is **continuous** if \exists a function f_X , called **probability density function (PDF)**, s.t.

$$\mathbb{P}(X \in B) = \int_B f_X(x) dx$$

- All of the concepts and methods (expectation, PMFs, and conditioning) for discrete rvs have continuous counterparts



- $\mathbb{P}(a \leq X \leq b) = \sum_{x:a \leq x \leq b} p_X(x)$
- $p_X(x) \geq 0, \sum_x p_X(x) = 1$

Continuous RV and Probability Density Function


- Many cases when random variable have “continuous values”, e.g., velocity of a car

Continuous Random Variable


A rv X is **continuous** if \exists a function f_X , called **probability density function (PDF)**, s.t.

$$\mathbb{P}(X \in B) = \int_B f_X(x) dx$$


- All of the concepts and methods (expectation, PMFs, and conditioning) for discrete rvs have continuous counterparts



- $\mathbb{P}(a \leq X \leq b) = \sum_{x: a \leq x \leq b} p_X(x)$
- $p_X(x) \geq 0, \sum_x p_X(x) = 1$




- $\mathbb{P}(a \leq X \leq b) = \int_a^b f_X(x) dx$
- $f_X(x) \geq 0, \int_{-\infty}^{\infty} f_X(x) dx = 1$




- $\mathbb{P}(a \leq X \leq a + \delta) \approx$

Examples




- $\mathbb{P}(a \leq X \leq a + \delta) \approx f_X(a) \cdot \delta$

Examples





- $\mathbb{P}(a \leq X \leq a + \delta) \approx f_X(a) \cdot \delta$
- $\mathbb{P}(X = a) = 0$


Examples




- $\mathbb{P}(a \leq X \leq a + \delta) \approx f_X(a) \cdot \delta$
- $\mathbb{P}(X = a) = 0$

Examples







- $\mathbb{E}[X] = \int_{-\infty}^{\infty} x f_X(x) dx =$
- $\mathbb{E}[X^2] = \int_{-\infty}^{\infty} x^2 f_X(x) dx =$
- $\text{var}[X] =$



- $\mathbb{E}[X] = \int_{-\infty}^{\infty} x f_X(x) dx = \int_a^b \frac{x}{b-a} dx = \frac{1}{b-a} \frac{b^2 - a^2}{2} = \frac{b+a}{2}$
- $\mathbb{E}[X^2] = \int_{-\infty}^{\infty} x^2 f_X(x) dx =$
- $\text{var}[X] =$



- $\mathbb{E}[X] = \int_{-\infty}^{\infty} x f_X(x) dx = \int_a^b \frac{x}{b-a} dx = \frac{1}{b-a} \frac{b^2 - a^2}{2} = \frac{b+a}{2}$
- $\mathbb{E}[X^2] = \int_{-\infty}^{\infty} x^2 f_X(x) dx = \int_a^b \frac{x^2}{b-a} dx = \frac{1}{b-a} \frac{b^3 - a^3}{3} = \frac{a^2 + ab + b^2}{3}$
- $\text{var}[X] =$



- $\mathbb{E}[X] = \int_{-\infty}^{\infty} x f_X(x) dx = \int_a^b \frac{x}{b-a} dx = \frac{1}{b-a} \frac{b^2 - a^2}{2} = \frac{b+a}{2}$
- $\mathbb{E}[X^2] = \int_{-\infty}^{\infty} x^2 f_X(x) dx = \int_a^b \frac{x^2}{b-a} dx = \frac{1}{b-a} \frac{b^3 - a^3}{3} = \frac{a^2 + ab + b^2}{3}$
- $\text{var}[X] = \frac{a^2 + ab + b^2}{3} - \frac{a^2 + 2ab + b^2}{4}$

Cumulative Distribution Function (CDF)

- Discrete: PMF, Continuous: PDF

Cumulative Distribution Function (CDF)

- Discrete: PMF, Continuous: PDF
- Can we describe all rvs with a single mathematical concept?

Cumulative Distribution Function (CDF)

- Discrete: PMF, Continuous: PDF
- Can we describe all rvs with a single mathematical concept?

$$F_X(x) = \mathbb{P}(X \leq x) =$$

$$\begin{cases} \sum_{k \leq x} p_X(k), & \text{discrete} \\ \int_{-\infty}^x f_X(t) dt, & \text{continuous} \end{cases}$$

Cumulative Distribution Function (CDF)

- Discrete: PMF, Continuous: PDF
- Can we describe all rvs with a single mathematical concept?

$$F_X(x) = \mathbb{P}(X \leq x) =$$

$$\begin{cases} \sum_{k \leq x} p_X(k), & \text{discrete} \\ \int_{-\infty}^x f_X(t) dt, & \text{continuous} \end{cases}$$

- always well defined, because we can always compute the probability for the event $\{X \leq x\}$

Cumulative Distribution Function (CDF)

- Discrete: PMF, Continuous: PDF
- Can we describe all rvs with a single mathematical concept?

$$F_X(x) = \mathbb{P}(X \leq x) =$$

$$\begin{cases} \sum_{k \leq x} p_X(k), & \text{discrete} \\ \int_{-\infty}^x f_X(t) dt, & \text{continuous} \end{cases}$$

- always well defined, because we can always compute the probability for the event $\{X \leq x\}$
- CCDF (Complementary CDF): $\mathbb{P}(X > x)$


Cumulative Distribution Function (CDF)

- Discrete: PMF, Continuous: PDF
- Can we describe all rvs with a single mathematical concept?

$$F_X(x) = \mathbb{P}(X \leq x) =$$

$$\begin{cases} \sum_{k \leq x} p_X(k), & \text{discrete} \\ \int_{-\infty}^x f_X(t) dt, & \text{continuous} \end{cases}$$

- always well defined, because we can always compute the probability for the event $\{X \leq x\}$
- CCDF (Complementary CDF): $\mathbb{P}(X > x)$




Cumulative Distribution Function (CDF)

- Discrete: PMF, Continuous: PDF
- Can we describe all rvs with a single mathematical concept?

$$F_X(x) = \mathbb{P}(X \leq x) =$$

$$\begin{cases} \sum_{k \leq x} p_X(k), & \text{discrete} \\ \int_{-\infty}^x f_X(t) dt, & \text{continuous} \end{cases}$$

- always well defined, because we can always compute the probability for the event $\{X \leq x\}$
- CCDF (Complementary CDF): $\mathbb{P}(X > x)$



- Non-decreasing

- Non-decreasing
- $F_X(x)$ tends to 1, as $x \rightarrow \infty$
- $F_X(x)$ tends to 0, as $x \rightarrow -\infty$

- Non-decreasing
- $F_X(x)$ tends to 1, as $x \rightarrow \infty$
- $F_X(x)$ tends to 0, as $x \rightarrow -\infty$


Now, let's look at famous continuous random variables popularly used in our life.

Exponential RV with parameter $\lambda > 0$: $\exp(\lambda)$

Exponential RV with parameter $\lambda > 0$: $\exp(\lambda)$

- A rv X is called **exponential with λ** , if

$$f_X(x) = \begin{cases} \lambda e^{-\lambda x}, & x \geq 0 \\ 0, & x < 0 \end{cases} \quad \text{or} \quad F_X(x) = 1 - e^{-\lambda x}$$




Exponential RV with parameter $\lambda > 0$: $\exp(\lambda)$

- A rv X is called **exponential with λ** , if

$$f_X(x) = \begin{cases} \lambda e^{-\lambda x}, & x \geq 0 \\ 0, & x < 0 \end{cases} \quad \text{or} \quad F_X(x) = 1 - e^{-\lambda x}$$

- Models a waiting time




Exponential RV with parameter $\lambda > 0$: $\exp(\lambda)$

- A rv X is called **exponential with λ** , if

$$f_X(x) = \begin{cases} \lambda e^{-\lambda x}, & x \geq 0 \\ 0, & x < 0 \end{cases} \quad \text{or} \quad F_X(x) = 1 - e^{-\lambda x}$$

- Models a waiting time
- CCDF $\mathbb{P}(X \geq x) = e^{-\lambda x}$ (waiting time decays exponentially)




Exponential RV with parameter $\lambda > 0$: $\exp(\lambda)$

- A rv X is called **exponential with λ** , if

$$f_X(x) = \begin{cases} \lambda e^{-\lambda x}, & x \geq 0 \\ 0, & x < 0 \end{cases} \quad \text{or} \quad F_X(x) = 1 - e^{-\lambda x}$$

- Models a waiting time
- CCDF $\mathbb{P}(X \geq x) = e^{-\lambda x}$ (waiting time decays exponentially)
- $\mathbb{E}[X] = 1/\lambda$, $\mathbb{E}[X^2] = 2/\lambda^2$, $\text{var}[X] = 1/\lambda^2$




Exponential RV with parameter $\lambda > 0$: $\exp(\lambda)$

- A rv X is called **exponential with λ** , if

$$f_X(x) = \begin{cases} \lambda e^{-\lambda x}, & x \geq 0 \\ 0, & x < 0 \end{cases} \quad \text{or} \quad F_X(x) = 1 - e^{-\lambda x}$$

- Models a waiting time
- CCDF $\mathbb{P}(X \geq x) = e^{-\lambda x}$ (waiting time decays exponentially)
- $\mathbb{E}[X] = 1/\lambda$, $\mathbb{E}[X^2] = 2/\lambda^2$, $\text{var}[X] = 1/\lambda^2$
- (Q) What is the discrete rv which models a waiting time?



Modeling Waiting Time? A Discrete Twin (1)

- A discrete twin for modeling waiting times is **geometric** rvs.

- A discrete twin for modeling waiting times is **geometric** rvs.
- Models a system evolution over time: Continuous time vs. Discrete time. In many cases, continuous case is the some type of of its corresponding discrete case.

- A discrete twin for modeling waiting times is **geometric** rvs.
- Models a system evolution over time: Continuous time vs. Discrete time. In many cases, continuous case is the some type of **limit** of its corresponding discrete case.
- Can you make mathematical description, where geometric and exponential rvs meet each other in the limit?

- A discrete twin for modeling waiting times is **geometric** rvs.
- Models a system evolution over time: Continuous time vs. Discrete time. In many cases, continuous case is the some type of **limit** of its corresponding discrete case.
- Can you make mathematical description, where geometric and exponential rvs meet each other in the limit?
- **Key idea.**
 - Continuous system: Discrete system with


- A discrete twin for modeling waiting times is **geometric** rvs.
- Models a system evolution over time: Continuous time vs. Discrete time. In many cases, continuous case is the some type of **limit** of its corresponding discrete case.
- Can you make mathematical description, where geometric and exponential rvs meet each other in the limit?
- **Key idea.**
 - Continuous system: Discrete system with **infinitely many slots whose duration is infinitely small.**

- A discrete twin for modeling waiting times is **geometric** rvs.
- Models a system evolution over time: Continuous time vs. Discrete time. In many cases, continuous case is the some type of **limit** of its corresponding discrete case.
- Can you make mathematical description, where geometric and exponential rvs meet each other in the limit?
- **Key idea.**
 - Continuous system: Discrete system with **infinitely many slots whose duration is infinitely small.**
- limiting system: $X_{exp}(\lambda)$ with CDF $F_{exp}(\cdot)$

- A discrete twin for modeling waiting times is **geometric** rvs.
- Models a system evolution over time: Continuous time vs. Discrete time. In many cases, continuous case is the some type of **limit** of its corresponding discrete case.
- Can you make mathematical description, where geometric and exponential rvs meet each other in the limit?
- **Key idea.**
 - Continuous system: Discrete system with **infinitely many slots whose duration is infinitely small.**
- limiting system: $X_{exp}(\lambda)$ with CDF $F_{exp}(\cdot)$
- n -th system: $X_{geo}^n(p_n)$ with CDF $F_{geo}^n(\cdot)$

Modeling Waiting Time? A Discrete Twin (2)


For a given $x > 0$,



Modeling Waiting Time? A Discrete Twin (2)

For a given $x > 0$,

- Define $\delta = \frac{x}{n}$ (a slot length in the n -th system)




Modeling Waiting Time? A Discrete Twin (2)

For a given $x > 0$,

- Define $\delta = \frac{x}{n}$ (a slot length in the n -th system)
- Remember

$$F_{exp}(x) = 1 - e^{-\lambda x}$$

$$F_{geo}^n(n) = 1 - (1 - p_n)^n$$



Modeling Waiting Time? A Discrete Twin (2)


For a given $x > 0$,

- Define $\delta = \frac{x}{n}$ (a slot length in the n -th system)
- Remember

$$F_{exp}(x) = 1 - e^{-\lambda x}$$

$$F_{geo}^n(n) = 1 - (1 - p_n)^n$$

- Choose $p_n = 1 - e^{-\lambda \delta} = 1 - e^{-\lambda \frac{x}{n}}$.



Modeling Waiting Time? A Discrete Twin (2)

For a given $x > 0$,


- Define $\delta = \frac{x}{n}$ (a slot length in the n -th system)
- Remember

$$F_{\text{exp}}(x) = 1 - e^{-\lambda x}$$

$$F_{\text{geo}}^n(n) = 1 - (1 - p_n)^n$$

- Choose $p_n = 1 - e^{-\lambda \delta} = 1 - e^{-\lambda \frac{x}{n}}$.
- As $n \rightarrow \infty$, the slot length $\delta \rightarrow 0$ thus $p_n \rightarrow 0$
- The CDF values of exponential and n -th geometric rvs become equal whenever $x = \delta, 2\delta, 3\delta, \dots$, i.e.,

$$F_{\text{exp}}(n\delta) = F_{\text{geo}}^n(n), \quad n = 1, 2, \dots$$



Modeling Waiting Time? A Discrete Twin (2)

For a given $x > 0$,


- Define $\delta = \frac{x}{n}$ (a slot length in the n -th system)
- Remember

$$F_{\text{exp}}(x) = 1 - e^{-\lambda x}$$

$$F_{\text{geo}}^n(n) = 1 - (1 - p_n)^n$$

- Choose $p_n = 1 - e^{-\lambda \delta} = 1 - e^{-\lambda \frac{x}{n}}$.
- As $n \rightarrow \infty$, the slot length $\delta \rightarrow 0$ thus $p_n \rightarrow 0$
- The CDF values of exponential and n -th geometric rvs become equal whenever $x = \delta, 2\delta, 3\delta, \dots$, i.e.,

$$F_{\text{exp}}(n\delta) = F_{\text{geo}}^n(n), \quad n = 1, 2, \dots$$



- As n grows, the number of slots grows, but the success probability over one slot decreases, so that everything is balanced up.

Modeling Waiting Time? A Discrete Twin (2)

For a given $x > 0$,


- Define $\delta = \frac{x}{n}$ (a slot length in the n -th system)
- Remember

$$F_{\text{exp}}(x) = 1 - e^{-\lambda x}$$

$$F_{\text{geo}}^n(n) = 1 - (1 - p_n)^n$$

- Choose $p_n = 1 - e^{-\lambda \delta} = 1 - e^{-\lambda \frac{x}{n}}$.
- As $n \rightarrow \infty$, the slot length $\delta \rightarrow 0$ thus $p_n \rightarrow 0$
- The CDF values of exponential and n -th geometric rvs become equal whenever $x = \delta, 2\delta, 3\delta, \dots$, i.e.,

$$F_{\text{exp}}(n\delta) = F_{\text{geo}}^n(n), \quad n = 1, 2, \dots$$



- As n grows, the number of slots grows, but the success probability over one slot decreases, so that everything is balanced up.
- As n grows, $F_{\text{geo}}^n(n)$ approaches $F_{\text{exp}}(n\delta)$.

Why important?

- Central limit theorem (중심극한정리)
 - One of the most remarkable findings in the probability theory
- Convenient analytical properties
- Modeling aggregate noise with many small, independent noise terms

- Standard Normal $N(0, 1)$

$$f_X(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$$

- $\mathbb{E}[X] = 0$
- $\text{var}[X] = 1$

- Standard Normal $N(0, 1)$

$$f_X(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$$

- $\mathbb{E}[X] = 0$
- $\text{var}[X] = 1$

- General Normal $N(\mu, \sigma^2)$

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-(x-\mu)^2/2\sigma^2}$$

- $\mathbb{E}[X] = \mu$
- $\text{var}[X] = \sigma^2$

- Standard Normal $N(0, 1)$

$$f_X(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$$

- $\mathbb{E}[X] = 0$
- $\text{var}[X] = 1$

- General Normal $N(\mu, \sigma^2)$

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-(x-\mu)^2/2\sigma^2}$$

- $\mathbb{E}[X] = \mu$
- $\text{var}[X] = \sigma^2$

Need to check:

- a legitimate PDF or not
- expectation/variance

- Linear transformation preserves normality

Linear transformation of Normal

If $X \sim \text{Norm}(\mu, \sigma^2)$, then for $a \neq 0$ and b $Y = aX + b \sim \text{Norm}(a\mu + b, a^2\sigma^2)$.

- Linear transformation preserves normality

Linear transformation of Normal

If $X \sim \text{Norm}(\mu, \sigma^2)$, then for $a \neq 0$ and b $Y = aX + b \sim \text{Norm}(a\mu + b, a^2\sigma^2)$.

- Thus, every normal rv can be :

If $X \sim \text{Norm}(\mu, \sigma^2)$, then $\sim \text{Norm}(0, 1)$

- Linear transformation preserves normality

Linear transformation of Normal

If $X \sim \text{Norm}(\mu, \sigma^2)$, then for $a \neq 0$ and b $Y = aX + b \sim \text{Norm}(a\mu + b, a^2\sigma^2)$.

- Thus, every normal rv can be **standardized** :

If $X \sim \text{Norm}(\mu, \sigma^2)$, then $Y = \frac{X - \mu}{\sigma} \sim \text{Norm}(0, 1)$

- Linear transformation preserves normality

Linear transformation of Normal

If $X \sim \text{Norm}(\mu, \sigma^2)$, then for $a \neq 0$ and b $Y = aX + b \sim \text{Norm}(a\mu + b, a^2\sigma^2)$.

- Thus, every normal rv can be **standardized** :

If $X \sim \text{Norm}(\mu, \sigma^2)$, then $Y = \frac{X - \mu}{\sigma} \sim \text{Norm}(0, 1)$

- Thus, we can make the **table** which records the following CDF values:

$$\Phi(y) = \mathbb{P}(Y \leq y) = \mathbb{P}(Y < y) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^y e^{-t^2/2} dt$$

Example

- Annual snowfall X is modeled as $Norm(60, 20^2)$. What is the probability that this year's snowfall is at least 80 inches?

| | .00 | .01 | .02 | .03 | .04 | .05 | .06 | .07 | .08 | .09 |
|-----|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 0.0 | .5000 | .5040 | .5080 | .5120 | .5160 | .5199 | .5239 | .5279 | .5319 | .5359 |
| 0.1 | .5398 | .5438 | .5478 | .5517 | .5557 | .5596 | .5636 | .5675 | .5714 | .5753 |
| 0.2 | .5793 | .5832 | .5871 | .5910 | .5948 | .5987 | .6026 | .6064 | .6103 | .6141 |
| 0.3 | .6179 | .6217 | .6255 | .6293 | .6331 | .6368 | .6406 | .6443 | .6480 | .6517 |
| 0.4 | .6554 | .6591 | .6628 | .6664 | .6700 | .6736 | .6772 | .6808 | .6844 | .6879 |
| 0.5 | .6915 | .6950 | .6985 | .7019 | .7054 | .7088 | .7123 | .7157 | .7190 | .7224 |
| 0.6 | .7257 | .7291 | .7324 | .7357 | .7389 | .7422 | .7454 | .7486 | .7517 | .7549 |
| 0.7 | .7580 | .7611 | .7642 | .7673 | .7704 | .7734 | .7764 | .7794 | .7823 | .7852 |
| 0.8 | .7881 | .7910 | .7939 | .7967 | .7995 | .8023 | .8051 | .8078 | .8106 | .8133 |
| 0.9 | .8159 | .8186 | .8212 | .8238 | .8264 | .8289 | .8315 | .8340 | .8365 | .8389 |
| 1.0 | .8413 | .8438 | .8461 | .8485 | .8508 | .8531 | .8554 | .8577 | .8599 | .8621 |
| 1.1 | .8643 | .8665 | .8686 | .8708 | .8729 | .8749 | .8770 | .8790 | .8810 | .8830 |
| 1.2 | .8849 | .8869 | .8888 | .8907 | .8925 | .8944 | .8962 | .8980 | .8997 | .9015 |
| 1.3 | .9032 | .9049 | .9066 | .9082 | .9099 | .9115 | .9131 | .9147 | .9162 | .9177 |
| 1.4 | .9192 | .9207 | .9222 | .9236 | .9251 | .9265 | .9279 | .9292 | .9306 | .9319 |
| 1.5 | .9332 | .9345 | .9357 | .9370 | .9382 | .9394 | .9406 | .9418 | .9429 | .9441 |
| 1.6 | .9452 | .9463 | .9474 | .9484 | .9495 | .9505 | .9515 | .9525 | .9535 | .9545 |
| 1.7 | .9554 | .9564 | .9573 | .9582 | .9591 | .9599 | .9608 | .9616 | .9625 | .9633 |
| 1.8 | .9641 | .9649 | .9656 | .9664 | .9671 | .9678 | .9686 | .9693 | .9699 | .9706 |
| 1.9 | .9713 | .9719 | .9726 | .9732 | .9738 | .9744 | .9750 | .9756 | .9761 | .9767 |
| 2.0 | .9772 | .9778 | .9783 | .9788 | .9793 | .9798 | .9803 | .9808 | .9812 | .9817 |
| 2.1 | .9821 | .9826 | .9830 | .9834 | .9838 | .9842 | .9846 | .9850 | .9854 | .9857 |
| 2.2 | .9861 | .9864 | .9868 | .9871 | .9875 | .9878 | .9881 | .9884 | .9887 | .9890 |
| 2.3 | .9893 | .9896 | .9898 | .9901 | .9904 | .9906 | .9909 | .9911 | .9913 | .9916 |
| 2.4 | .9918 | .9920 | .9922 | .9925 | .9927 | .9929 | .9931 | .9932 | .9934 | .9936 |
| 2.5 | .9938 | .9940 | .9941 | .9943 | .9945 | .9946 | .9948 | .9949 | .9951 | .9952 |
| 2.6 | .9953 | .9955 | .9956 | .9957 | .9959 | .9960 | .9961 | .9962 | .9963 | .9964 |
| 2.7 | .9965 | .9966 | .9967 | .9968 | .9969 | .9970 | .9971 | .9972 | .9973 | .9974 |
| 2.8 | .9974 | .9975 | .9976 | .9977 | .9977 | .9978 | .9979 | .9979 | .9980 | .9981 |
| 2.9 | .9981 | .9982 | .9982 | .9983 | .9984 | .9984 | .9985 | .9985 | .9986 | .9986 |

Example

- Annual snowfall X is modeled as $Norm(60, 20^2)$. What is the probability that this year's snowfall is at least 80 inches?
- $Y = \frac{X-60}{20}$.

| | .00 | .01 | .02 | .03 | .04 | .05 | .06 | .07 | .08 | .09 |
|-----|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 0.0 | .5000 | .5040 | .5080 | .5120 | .5160 | .5199 | .5239 | .5279 | .5319 | .5359 |
| 0.1 | .5398 | .5438 | .5478 | .5517 | .5557 | .5596 | .5636 | .5675 | .5714 | .5753 |
| 0.2 | .5793 | .5832 | .5871 | .5910 | .5948 | .5987 | .6026 | .6064 | .6103 | .6141 |
| 0.3 | .6179 | .6217 | .6255 | .6293 | .6331 | .6368 | .6406 | .6443 | .6480 | .6517 |
| 0.4 | .6554 | .6591 | .6628 | .6664 | .6700 | .6736 | .6772 | .6808 | .6844 | .6879 |
| 0.5 | .6915 | .6950 | .6985 | .7019 | .7054 | .7088 | .7123 | .7157 | .7190 | .7224 |
| 0.6 | .7257 | .7291 | .7324 | .7357 | .7389 | .7422 | .7454 | .7486 | .7517 | .7549 |
| 0.7 | .7580 | .7611 | .7642 | .7673 | .7704 | .7734 | .7764 | .7794 | .7823 | .7852 |
| 0.8 | .7881 | .7910 | .7939 | .7967 | .7995 | .8023 | .8051 | .8078 | .8106 | .8133 |
| 0.9 | .8159 | .8186 | .8212 | .8238 | .8264 | .8289 | .8315 | .8340 | .8365 | .8389 |
| 1.0 | .8413 | .8438 | .8461 | .8485 | .8508 | .8531 | .8554 | .8577 | .8599 | .8621 |
| 1.1 | .8643 | .8665 | .8686 | .8708 | .8729 | .8749 | .8770 | .8790 | .8810 | .8830 |
| 1.2 | .8849 | .8869 | .8888 | .8907 | .8925 | .8944 | .8962 | .8980 | .8997 | .9015 |
| 1.3 | .9032 | .9049 | .9066 | .9082 | .9099 | .9115 | .9131 | .9147 | .9162 | .9177 |
| 1.4 | .9192 | .9207 | .9222 | .9236 | .9251 | .9265 | .9279 | .9292 | .9306 | .9319 |
| 1.5 | .9332 | .9345 | .9357 | .9370 | .9382 | .9394 | .9406 | .9418 | .9429 | .9441 |
| 1.6 | .9452 | .9463 | .9474 | .9484 | .9495 | .9505 | .9515 | .9525 | .9535 | .9545 |
| 1.7 | .9554 | .9564 | .9573 | .9582 | .9591 | .9599 | .9608 | .9616 | .9625 | .9633 |
| 1.8 | .9641 | .9649 | .9656 | .9664 | .9671 | .9678 | .9686 | .9693 | .9699 | .9706 |
| 1.9 | .9713 | .9719 | .9726 | .9732 | .9738 | .9744 | .9750 | .9756 | .9761 | .9767 |
| 2.0 | .9772 | .9778 | .9783 | .9788 | .9793 | .9798 | .9803 | .9808 | .9812 | .9817 |
| 2.1 | .9821 | .9826 | .9830 | .9834 | .9838 | .9842 | .9846 | .9850 | .9854 | .9857 |
| 2.2 | .9861 | .9864 | .9868 | .9871 | .9875 | .9878 | .9881 | .9884 | .9887 | .9890 |
| 2.3 | .9893 | .9896 | .9898 | .9901 | .9904 | .9906 | .9909 | .9911 | .9913 | .9916 |
| 2.4 | .9918 | .9920 | .9922 | .9925 | .9927 | .9929 | .9931 | .9932 | .9934 | .9936 |
| 2.5 | .9938 | .9940 | .9941 | .9943 | .9945 | .9946 | .9948 | .9949 | .9951 | .9952 |
| 2.6 | .9953 | .9955 | .9956 | .9957 | .9959 | .9960 | .9961 | .9962 | .9963 | .9964 |
| 2.7 | .9965 | .9966 | .9967 | .9968 | .9969 | .9970 | .9971 | .9972 | .9973 | .9974 |
| 2.8 | .9974 | .9975 | .9976 | .9977 | .9977 | .9978 | .9979 | .9979 | .9980 | .9981 |
| 2.9 | .9981 | .9982 | .9982 | .9983 | .9984 | .9984 | .9985 | .9985 | .9986 | .9986 |

Example

- Annual snowfall X is modeled as $Norm(60, 20^2)$. What is the probability that this year's snowfall is at least 80 inches?
- $Y = \frac{X-60}{20}$.

$$\begin{aligned}\mathbb{P}(X \geq 80) &= \mathbb{P}\left(Y \geq \frac{80 - 60}{20}\right) \\ &= \mathbb{P}(Y \geq 1) = 1 - \Phi(1) \\ &= 1 - 0.8413 = 0.1587\end{aligned}$$

| | .00 | .01 | .02 | .03 | .04 | .05 | .06 | .07 | .08 | .09 |
|-----|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 0.0 | .5000 | .5040 | .5080 | .5120 | .5160 | .5199 | .5239 | .5279 | .5319 | .5359 |
| 0.1 | .5398 | .5438 | .5478 | .5517 | .5557 | .5596 | .5636 | .5675 | .5714 | .5753 |
| 0.2 | .5793 | .5832 | .5871 | .5910 | .5948 | .5987 | .6026 | .6064 | .6103 | .6141 |
| 0.3 | .6179 | .6217 | .6255 | .6293 | .6331 | .6368 | .6406 | .6443 | .6480 | .6517 |
| 0.4 | .6554 | .6591 | .6628 | .6664 | .6700 | .6736 | .6772 | .6808 | .6844 | .6879 |
| 0.5 | .6915 | .6950 | .6985 | .7019 | .7054 | .7088 | .7123 | .7157 | .7190 | .7224 |
| 0.6 | .7257 | .7291 | .7324 | .7357 | .7389 | .7422 | .7454 | .7486 | .7517 | .7549 |
| 0.7 | .7580 | .7611 | .7642 | .7673 | .7704 | .7734 | .7764 | .7794 | .7823 | .7852 |
| 0.8 | .7881 | .7910 | .7939 | .7967 | .7995 | .8023 | .8051 | .8078 | .8106 | .8133 |
| 0.9 | .8159 | .8186 | .8212 | .8238 | .8264 | .8289 | .8315 | .8340 | .8365 | .8389 |
| 1.0 | .8413 | .8438 | .8461 | .8485 | .8508 | .8531 | .8554 | .8577 | .8599 | .8621 |
| 1.1 | .8643 | .8665 | .8686 | .8708 | .8729 | .8749 | .8770 | .8790 | .8810 | .8830 |
| 1.2 | .8849 | .8869 | .8888 | .8907 | .8925 | .8944 | .8962 | .8980 | .8997 | .9015 |
| 1.3 | .9032 | .9049 | .9066 | .9082 | .9099 | .9115 | .9131 | .9147 | .9162 | .9177 |
| 1.4 | .9192 | .9207 | .9222 | .9236 | .9251 | .9265 | .9279 | .9292 | .9306 | .9319 |
| 1.5 | .9332 | .9345 | .9357 | .9370 | .9382 | .9394 | .9406 | .9418 | .9429 | .9441 |
| 1.6 | .9452 | .9463 | .9474 | .9484 | .9495 | .9505 | .9515 | .9525 | .9535 | .9545 |
| 1.7 | .9554 | .9564 | .9573 | .9582 | .9591 | .9599 | .9608 | .9616 | .9625 | .9633 |
| 1.8 | .9641 | .9649 | .9656 | .9664 | .9671 | .9678 | .9686 | .9693 | .9699 | .9706 |
| 1.9 | .9713 | .9719 | .9726 | .9732 | .9738 | .9744 | .9750 | .9756 | .9761 | .9767 |
| 2.0 | .9772 | .9778 | .9783 | .9788 | .9793 | .9798 | .9803 | .9808 | .9812 | .9817 |
| 2.1 | .9821 | .9826 | .9830 | .9834 | .9838 | .9842 | .9846 | .9850 | .9854 | .9857 |
| 2.2 | .9861 | .9864 | .9868 | .9871 | .9875 | .9878 | .9881 | .9884 | .9887 | .9890 |
| 2.3 | .9893 | .9896 | .9898 | .9901 | .9904 | .9906 | .9909 | .9911 | .9913 | .9916 |
| 2.4 | .9918 | .9920 | .9922 | .9925 | .9927 | .9929 | .9931 | .9932 | .9934 | .9936 |
| 2.5 | .9938 | .9940 | .9941 | .9943 | .9945 | .9946 | .9948 | .9949 | .9951 | .9952 |
| 2.6 | .9953 | .9955 | .9956 | .9957 | .9959 | .9960 | .9961 | .9962 | .9963 | .9964 |
| 2.7 | .9965 | .9966 | .9967 | .9968 | .9969 | .9970 | .9971 | .9972 | .9973 | .9974 |
| 2.8 | .9974 | .9975 | .9976 | .9977 | .9977 | .9978 | .9979 | .9979 | .9980 | .9981 |
| 2.9 | .9981 | .9982 | .9982 | .9983 | .9984 | .9984 | .9985 | .9985 | .9986 | .9986 |

- Famous discrete random variables used in the community
 - Bernoulli, Uniform, Binomial, Geometric, Poisson, etc.
- Summarizing a random variable: Expectation and Variance
- Functions of a single random variable, Functions of multiple random variables
- Conditioning for random variables, Independence for random variables
- Continuous random variables
 - Normal, Uniform, Exponential, etc.
- Bayes' rule for random variables

** Continuous counterparts are intuitively understandable. So, we will be quick at reviewing them.

Jointly Continuous

Two continuous rvs are if a non-negative function $f_{X,Y}(x,y)$ (called joint PDF) satisfies: for every subset B of the two dimensional plane,

$$\mathbb{P}((X, Y) \in B) = \iint_{(x,y) \in B} f_{X,Y}(x,y) dx dy$$

Jointly Continuous

Two continuous rvs are **jointly continuous** if a non-negative function $f_{X,Y}(x,y)$ (called joint PDF) satisfies: for **every** subset B of the two dimensional plane,

$$\mathbb{P}((X, Y) \in B) = \iint_{(x,y) \in B} f_{X,Y}(x,y) dx dy$$

Jointly Continuous

Two continuous rvs are **jointly continuous** if a non-negative function $f_{X,Y}(x, y)$ (called joint PDF) satisfies: for **every** subset B of the two dimensional plane,

$$\mathbb{P}((X, Y) \in B) = \iint_{(x,y) \in B} f_{X,Y}(x, y) dx dy$$

1. The joint PDF is used to calculate probabilities

$$\mathbb{P}((X, Y) \in B) = \iint_{(x,y) \in B} f_{X,Y}(x, y) dx dy$$

Our particular interest: $B = \{(x, y) \mid a \leq x \leq b, c \leq y \leq d\}$

2. The marginal PDFs of X and Y are from the joint PDF as:

$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x,y)dy, \quad f_Y(y) = \int_{-\infty}^{\infty} f_{X,Y}(x,y)dx$$

2. The marginal PDFs of X and Y are from the joint PDF as:

$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dy, \quad f_Y(y) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx$$

3. The joint CDF is defined by $F_{X,Y}(x, y) = \mathbb{P}(X \leq x, Y \leq y)$, and determines the joint PDF as:

$$f_{X,Y}(x, y) = \frac{\partial^2 F_{x,y}}{\partial x \partial y}(x, y)$$

2. The marginal PDFs of X and Y are from the joint PDF as:

$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dy, \quad f_Y(y) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx$$

3. The joint CDF is defined by $F_{X,Y}(x, y) = \mathbb{P}(X \leq x, Y \leq y)$, and determines the joint PDF as:

$$f_{X,Y}(x, y) = \frac{\partial^2 F_{x,y}}{\partial x \partial y}(x, y)$$

4. A function $g(X, Y)$ of X and Y defines a new random variable, and

$$\mathbb{E}[g(X, Y)] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x, y) f_{X,Y}(x, y) dx dy$$

* Conditional PDF, given an event

* Conditional PDF, given $X \in B$

* Conditional PDF, given an event

- $f_X(x) \cdot \delta \approx \mathbb{P}(x \leq X \leq x + \delta)$
 $f_{X|A}(x) \cdot \delta \approx \mathbb{P}(x \leq X \leq x + \delta | A)$

* Conditional PDF, given $X \in B$

* Conditional PDF, given an event

- $f_X(x) \cdot \delta \approx \mathbb{P}(x \leq X \leq x + \delta)$
 $f_{X|A}(x) \cdot \delta \approx \mathbb{P}(x \leq X \leq x + \delta | A)$

- $\mathbb{P}(X \in B) = \int_B f_X(x) dx$
 $\mathbb{P}(X \in B | A) = \int_B f_{X|A}(x) dx$

Note: A is an event, but B is a subset that includes the possible values which can be taken by the rv X .

* Conditional PDF, given $X \in B$

* Conditional PDF, given an event

- $f_X(x) \cdot \delta \approx \mathbb{P}(x \leq X \leq x + \delta)$
 $f_{X|A}(x) \cdot \delta \approx \mathbb{P}(x \leq X \leq x + \delta | A)$

- $\mathbb{P}(X \in B) = \int_B f_X(x) dx$
 $\mathbb{P}(X \in B | A) = \int_B f_{X|A}(x) dx$

Note: A is an event, but B is a subset that includes the possible values which can be taken by the rv X .

- $\int f_{X|A}(x) = 1$

* Conditional PDF, given $X \in B$

* Conditional PDF, given an event

- $f_X(x) \cdot \delta \approx \mathbb{P}(x \leq X \leq x + \delta)$
 $f_{X|A}(x) \cdot \delta \approx \mathbb{P}(x \leq X \leq x + \delta | A)$

- $\mathbb{P}(X \in B) = \int_B f_X(x) dx$
 $\mathbb{P}(X \in B | A) = \int_B f_{X|A}(x) dx$

Note: A is an event, but B is a subset that includes the possible values which can be taken by the rv X .

- $\int f_{X|A}(x) = 1$

* Conditional PDF, given $X \in B$

$$\mathbb{P}(x \leq X \leq x + \delta | X \in B) \approx f_{X|X \in B}(x) \cdot \delta$$

$$f_{X|X \in B}(x) = \begin{cases} 0, & \text{if } x \notin B \\ \frac{f_X(x)}{\mathbb{P}(B)}, & \text{if } x \in B \end{cases}$$

* Conditional PDF, given an event

- $f_X(x) \cdot \delta \approx \mathbb{P}(x \leq X \leq x + \delta)$
 $f_{X|A}(x) \cdot \delta \approx \mathbb{P}(x \leq X \leq x + \delta | A)$

- $\mathbb{P}(X \in B) = \int_B f_X(x) dx$
 $\mathbb{P}(X \in B | A) = \int_B f_{X|A}(x) dx$

Note: A is an event, but B is a subset that includes the possible values which can be taken by the rv X .

- $\int f_{X|A}(x) = 1$

* Conditional PDF, given $X \in B$


$$\mathbb{P}(x \leq X \leq x + \delta | X \in B) \approx f_{X|X \in B}(x) \cdot \delta$$

$$f_{X|X \in B}(x) = \begin{cases} 0, & \text{if } x \notin B \\ \frac{f_X(x)}{\mathbb{P}(B)}, & \text{if } x \in B \end{cases}$$

(Q) In the discrete, we consider the event $\{X = x\}$, not $\{X \in B\}$. Why?



Continuous: Conditional Expectation

$$A = \left\{ \frac{a+b}{2} \leq X \leq b \right\}$$





Continuous: Conditional Expectation

$$A = \left\{ \frac{a+b}{2} \leq X \leq b \right\}$$



Continuous: Conditional Expectation



$$A = \left\{ \frac{a+b}{2} \leq X \leq b \right\}$$



- $\mathbb{E}[X] = \int x f_X(x) dx$
 $\mathbb{E}[X|A] = \int x f_{X|A}(x) dx$

Continuous: Conditional Expectation



$$A = \left\{ \frac{a+b}{2} \leq X \leq b \right\}$$



- $\mathbb{E}[X] = \int x f_X(x) dx$
 $\mathbb{E}[X|A] = \int x f_{X|A}(x) dx$
- $\mathbb{E}[g(X)] = \int g(x) f_X(x) dx$
 $\mathbb{E}[g(X)|A] = \int g(x) f_{X|A}(x) dx$

Continuous: Conditional Expectation

$$A = \left\{ \frac{a+b}{2} \leq X \leq b \right\}$$





- $\mathbb{E}[X] = \int x f_X(x) dx$
 $\mathbb{E}[X|A] = \int x f_{X|A}(x) dx$
- $\mathbb{E}[g(X)] = \int g(x) f_X(x) dx$
 $\mathbb{E}[g(X)|A] = \int g(x) f_{X|A}(x) dx$

$$\mathbb{E}[X|A] =$$

$$\mathbb{E}[X^2|A] =$$

Continuous: Conditional Expectation

$$A = \left\{ \frac{a+b}{2} \leq X \leq b \right\}$$



- $\mathbb{E}[X] = \int x f_X(x) dx$
 $\mathbb{E}[X|A] = \int x f_{X|A}(x) dx$
- $\mathbb{E}[g(X)] = \int g(x) f_X(x) dx$
 $\mathbb{E}[g(X)|A] = \int g(x) f_{X|A}(x) dx$

$$\mathbb{E}[X|A] = \int_{(a+b)/2}^b x \frac{2}{b-a} dx = \frac{a}{4} + \frac{3b}{4}$$

$$\mathbb{E}[X^2|A] = \int_{(a+b)/2}^b x^2 \frac{2}{b-a} dx =$$

- Exponential rv is a continuous counterpart of geometric rv.

- Exponential rv is a continuous counterpart of geometric rv.
- Thus, expected to be memoryless.

- Exponential rv is a continuous counterpart of geometric rv.
- Thus, expected to be memoryless.

Definition. A random variable X is called **memoryless** if, for any $n, m \geq 0$,

$$\mathbb{P}(X > n + m | X > m) = \mathbb{P}(X > n)$$

- Exponential rv is a continuous counterpart of geometric rv.
- Thus, expected to be memoryless.

Definition. A random variable X is called **memoryless** if, for any $n, m \geq 0$,

$$\mathbb{P}(X > n + m | X > m) = \mathbb{P}(X > n)$$

- **Proof.** Note that $\mathbb{P}(X > x) = e^{-\lambda x}$.

- Exponential rv is a continuous counterpart of geometric rv.
- Thus, expected to be memoryless.

Definition. A random variable X is called **memoryless** if, for any $n, m \geq 0$,

$$\mathbb{P}(X > n + m | X > m) = \mathbb{P}(X > n)$$

- **Proof.** Note that $\mathbb{P}(X > x) = e^{-\lambda x}$. Then,

$$\mathbb{P}(X > n + m | X > m) = \frac{\mathbb{P}(X > n + m)}{\mathbb{P}(X > m)} = \frac{e^{-\lambda(n+m)}}{e^{-\lambda m}} = e^{-\lambda n} = \mathbb{P}(X > n)$$

Total Probability/Expectation Theorem

Partition of Ω into A_1, A_2, A_3, \dots

* Discrete case

* Continuous case

Total Probability/Expectation Theorem

Partition of Ω into A_1, A_2, A_3, \dots

* Discrete case

Total Probability Theorem

$$\begin{aligned} p_X(x) &= \sum_i \mathbb{P}(A_i) \mathbb{P}(X = x | A_i) \\ &= \sum_i \mathbb{P}(A_i) p_{X|A_i}(x) \end{aligned}$$

Total Expectation Theorem

$$\mathbb{E}[X] = \sum_i \mathbb{P}(A_i) \mathbb{E}[X | A_i]$$

* Continuous case

Total Probability/Expectation Theorem

Partition of Ω into A_1, A_2, A_3, \dots

* Discrete case

Total Probability Theorem

$$\begin{aligned} p_X(x) &= \sum_i \mathbb{P}(A_i) \mathbb{P}(X = x | A_i) \\ &= \sum_i \mathbb{P}(A_i) p_{X|A_i}(x) \end{aligned}$$

Total Expectation Theorem

$$\mathbb{E}[X] = \sum_i \mathbb{P}(A_i) \mathbb{E}[X | A_i]$$

* Continuous case

Total Probability Theorem

$$f_X(x) = \sum_i \mathbb{P}(A_i) f_{X|A_i}(x)$$

Total Probability/Expectation Theorem

Partition of Ω into A_1, A_2, A_3, \dots

* Discrete case

Total Probability Theorem

$$\begin{aligned} p_X(x) &= \sum_i \mathbb{P}(A_i) \mathbb{P}(X = x | A_i) \\ &= \sum_i \mathbb{P}(A_i) p_{X|A_i}(x) \end{aligned}$$

Total Expectation Theorem

$$\mathbb{E}[X] = \sum_i \mathbb{P}(A_i) \mathbb{E}[X | A_i]$$

* Continuous case

Total Probability Theorem


$$f_X(x) = \sum_i \mathbb{P}(A_i) f_{X|A_i}(x)$$

Total Expectation Theorem

$$\mathbb{E}[X] = \sum_i \mathbb{P}(A_i) \mathbb{E}[X | A_i]$$


Example

- Your train's arrival every quarter hour (0, 15min, 30min, 45min).
- Your arrival $\sim \text{uniform}(7:10, 7:30)$ am.




Example

- Your train's arrival every quarter hour (0, 15min, 30min, 45min).
- Your arrival $\sim \text{uniform}(7:10, 7:30)$ am.
- What is the PDF of waiting time for the first train?



Example

- Your train's arrival every quarter hour (0, 15min, 30min, 45min).
- Your arrival $\sim \text{uniform}(7:10, 7:30)$ am.
- What is the PDF of waiting time for the first train?
- X : your arrival time, Y : waiting time.




Example

- Your train's arrival every quarter hour (0, 15min, 30min, 45min).
- Your arrival $\sim \text{uniform}(7:10, 7:30)$ am.
- What is the PDF of waiting time for the first train?
- X : your arrival time, Y : waiting time.
- The value of X makes a different waiting time. So, consider two events:

$$A = \{7:10 \leq X \leq 7:15\}$$

$$B = \{7:15 \leq X \leq 7:30\}$$




Example

- Your train's arrival every quarter hour (0, 15min, 30min, 45min).
- Your arrival $\sim \text{uniform}(7:10, 7:30)$ am.
- What is the PDF of waiting time for the first train?
- X : your arrival time, Y : waiting time.
- The value of X makes a different waiting time. So, consider two events:

$$A = \{7:10 \leq X \leq 7:15\}$$

$$B = \{7:15 \leq X \leq 7:30\}$$



$$f_Y(y) = \mathbb{P}(A)f_{Y|A}(y) + \mathbb{P}(B)f_{Y|B}(y)$$

$$\text{for } 0 \leq y \leq 5$$


$$\text{for } 5 < y \leq 15$$

Example

- Your train's arrival every quarter hour (0, 15min, 30min, 45min).
- Your arrival $\sim \text{uniform}(7:10, 7:30)$ am.
- What is the PDF of waiting time for the first train?
- X : your arrival time, Y : waiting time.
- The value of X makes a different waiting time. So, consider two events:

$$A = \{7:10 \leq X \leq 7:15\}$$

$$B = \{7:15 \leq X \leq 7:30\}$$



$$f_Y(y) = \mathbb{P}(A)f_{Y|A}(y) + \mathbb{P}(B)f_{Y|B}(y)$$

$$f_Y(y) = \frac{1}{4} \frac{1}{5} + \frac{3}{4} \frac{1}{15} = \frac{1}{10}, \quad \text{for } 0 \leq y \leq 5$$


$$\text{for } 5 < y \leq 15$$

Example

- Your train's arrival every quarter hour (0, 15min, 30min, 45min).
- Your arrival $\sim \text{uniform}(7:10, 7:30)$ am.
- What is the PDF of waiting time for the first train?
- X : your arrival time, Y : waiting time.
- The value of X makes a different waiting time. So, consider two events:

$$A = \{7:10 \leq X \leq 7:15\}$$

$$B = \{7:15 \leq X \leq 7:30\}$$



$$f_Y(y) = \mathbb{P}(A)f_{Y|A}(y) + \mathbb{P}(B)f_{Y|B}(y)$$

$$f_Y(y) = \frac{1}{4} \frac{1}{5} + \frac{3}{4} \frac{1}{15} = \frac{1}{10}, \quad \text{for } 0 \leq y \leq 5$$

$$f_Y(y) = \frac{1}{4} 0 + \frac{3}{4} \frac{1}{15} = \frac{1}{20}, \quad \text{for } 5 < y \leq 15$$

Continuous: Conditional PDF given a RV

- $p_{X|Y}(x|y) = \frac{p_{X,Y}(x,y)}{p_Y(y)}$

Continuous: Conditional PDF given a RV

- $p_{X|Y}(x|y) = \frac{p_{X,Y}(x,y)}{p_Y(y)}$
- Similarly, for $f_Y(y) > 0$,

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)}$$

Continuous: Conditional PDF given a RV

- $p_{X|Y}(x|y) = \frac{p_{X,Y}(x,y)}{p_Y(y)}$

- Similarly, for $f_Y(y) > 0$,

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)}$$

- Remember: For a fixed event A , $\mathbb{P}(\cdot|A)$ is a legitimate probability law.

Continuous: Conditional PDF given a RV

- $p_{X|Y}(x|y) = \frac{p_{X,Y}(x,y)}{p_Y(y)}$

- Similarly, for $f_Y(y) > 0$,

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)}$$

- Remember: For a fixed event A , $\mathbb{P}(\cdot|A)$ is a legitimate probability law.
- Similarly, For a fixed y , $f_{X|Y}(x|y)$ is a legitimate PDF, since

$$\int_{-\infty}^{\infty} f_{X|Y}(x|y) dx = \frac{\int_{-\infty}^{\infty} f_{X,Y}(x,y) dx}{f_Y(y)} = 1$$

Continuous: Conditional PDF given a RV

- $p_{X|Y}(x|y) = \frac{p_{X,Y}(x,y)}{p_Y(y)}$

- Similarly, for $f_Y(y) > 0$,

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)}$$

- Remember: For a fixed event A , $\mathbb{P}(\cdot|A)$ is a legitimate probability law.
- Similarly, For a fixed y , $f_{X|Y}(x|y)$ is a legitimate PDF, since

$$\int_{-\infty}^{\infty} f_{X|Y}(x|y) dx = \frac{\int_{-\infty}^{\infty} f_{X,Y}(x,y) dx}{f_Y(y)} = 1$$

- Multiplication rule.

$$\begin{aligned} f_{X,Y}(x,y) &= f_Y(y) \cdot f_{X|Y}(x|y) \\ &= f_X(x) f_{Y|X}(y|x) \end{aligned}$$

Continuous: Conditional PDF given a RV

- $p_{X|Y}(x|y) = \frac{p_{X,Y}(x,y)}{p_Y(y)}$

- Similarly, for $f_Y(y) > 0$,

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)}$$

- Remember: For a fixed event A , $\mathbb{P}(\cdot|A)$ is a legitimate probability law.
- Similarly, For a fixed y , $f_{X|Y}(x|y)$ is a legitimate PDF, since

$$\int_{-\infty}^{\infty} f_{X|Y}(x|y) dx = \frac{\int_{-\infty}^{\infty} f_{X,Y}(x,y) dx}{f_Y(y)} = 1$$

- Multiplication rule.

$$\begin{aligned} f_{X,Y}(x,y) &= f_Y(y) \cdot f_{X|Y}(x|y) \\ &= f_X(x) f_{Y|X}(y|x) \end{aligned}$$

- Total prob./exp. theorem.

$$f_X(x) = \int_{-\infty}^{\infty} f_Y(y) f_{X|Y}(x|y) dy$$

$$\mathbb{E}[X|Y = y] = \int_{-\infty}^{\infty} x f_{X|Y}(x|y) dx$$

$$\mathbb{E}[X] = \int_{-\infty}^{\infty} f_Y(y) \mathbb{E}[X|Y = y] dy$$

Continuous: Conditional PDF given a RV

- $p_{X|Y}(x|y) = \frac{p_{X,Y}(x,y)}{p_Y(y)}$

- Similarly, for $f_Y(y) > 0$,

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)}$$

- Remember: For a fixed event A , $\mathbb{P}(\cdot|A)$ is a legitimate probability law.
- Similarly, For a fixed y , $f_{X|Y}(x|y)$ is a legitimate PDF, since

$$\int_{-\infty}^{\infty} f_{X|Y}(x|y) dx = \frac{\int_{-\infty}^{\infty} f_{X,Y}(x,y) dx}{f_Y(y)} = 1$$

- Multiplication rule.

$$\begin{aligned} f_{X,Y}(x,y) &= f_Y(y) \cdot f_{X|Y}(x|y) \\ &= f_X(x) f_{Y|X}(y|x) \end{aligned}$$

- Total prob./exp. theorem.

$$f_X(x) = \int_{-\infty}^{\infty} f_Y(y) f_{X|Y}(x|y) dy$$

$$\mathbb{E}[X|Y = y] = \int_{-\infty}^{\infty} x f_{X|Y}(x|y) dx$$

$$\mathbb{E}[X] = \int_{-\infty}^{\infty} f_Y(y) \mathbb{E}[X|Y = y] dy$$

- Independence.

$$f_{X,Y}(x,y) = f_X(x) f_Y(y), \quad \text{for all } x \text{ and } y$$

Example: Stick-breaking (Ch 3. Prob 21)

- Break a stick of length l twice
 - first break at $X \sim \text{uniform}[0, l]$
 - second break at $Y \sim \text{uniform}[0, X]$

Example: Stick-breaking (Ch 3. Prob 21)

- Break a stick of length l twice
 - first break at $X \sim \text{uniform}[0, l]$
 - second break at $Y \sim \text{uniform}[0, X]$
- (Q) What is $\mathbb{E}[Y]$?

Example: Stick-breaking (Ch 3. Prob 21)

- Break a stick of length l twice
 - first break at $X \sim \text{uniform}[0, l]$
 - second break at $Y \sim \text{uniform}[0, X]$
- (Q) What is $\mathbb{E}[Y]$?
- Since Y depends on X , the total expectation theorem seems useful.

$$\mathbb{E}[Y] = \int_{-\infty}^{\infty} f_X(x) \mathbb{E}[Y|X = x] dx$$

Example: Stick-breaking (Ch 3. Prob 21)

- Break a stick of length l twice
 - first break at $X \sim \text{uniform}[0, l]$
 - second break at $Y \sim \text{uniform}[0, X]$
- (Q) What is $\mathbb{E}[Y]$?
- Since Y depends on X , the total expectation theorem seems useful.

$$\mathbb{E}[Y] = \int_{-\infty}^{\infty} f_X(x) \mathbb{E}[Y|X = x] dx$$

- Using the TET,

$$\begin{aligned} \mathbb{E}[Y] &= \int_0^l \frac{1}{l} \mathbb{E}[Y|X = x] dx \\ &= \int_0^l \frac{1}{l} \frac{x}{2} dx = \frac{l}{4} \end{aligned}$$

Example: Stick-breaking (Ch 3. Prob 21)

- Break a stick of length l twice
 - first break at $X \sim \text{uniform}[0, l]$
 - second break at $Y \sim \text{uniform}[0, X]$
- (Q) What is $\mathbb{E}[Y]$?
- Since Y depends on X , the total expectation theorem seems useful.

$$\mathbb{E}[Y] = \int_{-\infty}^{\infty} f_X(x) \mathbb{E}[Y|X = x] dx$$

- Using the TET,

$$\begin{aligned} \mathbb{E}[Y] &= \int_0^l \frac{1}{l} \mathbb{E}[Y|X = x] dx \\ &= \int_0^l \frac{1}{l} \frac{x}{2} dx = \frac{l}{4} \end{aligned}$$

- $f_X(x)$ and $f_{Y|X}(y|x)$ seems easy to compute. Thus,

$$f_{X,Y}(x, y) = f_X(x) f_{Y|X}(y|x) = \frac{1}{l} \cdot \frac{1}{x}$$

You can do many other things with the joint PDF.

- Famous discrete random variables used in the community
 - Bernoulli, Uniform, Binomial, Geometric, Poisson, etc.
- Summarizing a random variable: Expectation and Variance
- Functions of a single random variable, Functions of multiple random variables
- Conditioning for random variables, Independence for random variables
- Continuous random variables
 - Normal, Uniform, Exponential, etc.
- Bayes' rule for random variables

- X : state/cause/original value \rightarrow Y : result/resulting action/noisy measurement
- Model: $\mathbb{P}(X)$ (prior) and $\mathbb{P}(Y|X)$ (cause \rightarrow result)
- Inference: $\mathbb{P}(X|Y)$?

- X : state/cause/original value \rightarrow Y : result/resulting action/noisy measurement
- Model: $\mathbb{P}(X)$ (prior) and $\mathbb{P}(Y|X)$ (cause \rightarrow result)
- Inference: $\mathbb{P}(X|Y)$?

$$\begin{aligned}p_{X,Y}(x,y) &= p_X(x)p_{Y|X}(y|x) \\ &= p_Y(y)p_{X|Y}(x|y) \\ \textcolor{red}{p_{X|Y}(x|y)} &= \frac{p_X(x)p_{Y|X}(y|x)}{p_Y(y)} \\ p_Y(y) &= \sum_{x'} p_X(x')p_{Y|X}(y|x')\end{aligned}$$

$$\begin{aligned}f_{X,Y}(x,y) &= f_X(x)f_{Y|X}(y|x) \\ &= f_Y(y)f_{X|Y}(x|y) \\ \textcolor{red}{f_{X|Y}(x|y)} &= \frac{f_X(x)f_{Y|X}(y|x)}{f_Y(y)} \\ f_Y(y) &= \int f_X(x')f_{Y|X}(y|x')dx'\end{aligned}$$

K : discrete, Y : continuous

K : discrete, Y : continuous

- Inference of K given Y

- Inference of Y given K

K : discrete, Y : continuous

- Inference of K given Y

$$p_{K|Y}(k|y) = \frac{p_K(k)f_{Y|K}(y|k)}{f_Y(y)}$$

$$f_Y(y) = \sum_{k'} p_K(k')f_{Y|K}(y|k')$$

- Inference of Y given K

K : discrete, Y : continuous

- Inference of K given Y

$$p_{K|Y}(k|y) = \frac{p_K(k)f_{Y|K}(y|k)}{f_Y(y)}$$

$$f_Y(y) = \sum_{k'} p_K(k')f_{Y|K}(y|k')$$

- Inference of Y given K

$$f_{Y|K}(y|k) = \frac{f_Y(y)p_{K|Y}(k|y)}{p_K(k)}$$

$$p_K(k) = \int f_Y(y')p_{K|Y}(k|y')dy'$$

Example: Signal Detection (1)

Inference of discrete K given continuous Y :

$$p_{K|Y}(k|y) = \frac{p_K(k)f_{Y|K}(y|k)}{f_Y(y)}, \quad f_Y(y) = \sum_{k'} p_K(k')f_{Y|K}(y|k')$$

Inference of discrete K given continuous Y :

$$p_{K|Y}(k|y) = \frac{p_K(k)f_{Y|K}(y|k)}{f_Y(y)}, \quad f_Y(y) = \sum_{k'} p_K(k')f_{Y|K}(y|k')$$

- K : -1, +1, original signal, equally likely. $p_K(1) = 1/2, p_K(-1) = 1/2$.

Inference of discrete K given continuous Y :

$$p_{K|Y}(k|y) = \frac{p_K(k)f_{Y|K}(y|k)}{f_Y(y)}, \quad f_Y(y) = \sum_{k'} p_K(k')f_{Y|K}(y|k')$$

- K : -1, +1, original signal, equally likely. $p_K(1) = 1/2, p_K(-1) = 1/2$.
- Y : measured signal with Gaussian noise, $Y = K + W, W \sim N(0, 1)$

Inference of discrete K given continuous Y :

$$p_{K|Y}(k|y) = \frac{p_K(k)f_{Y|K}(y|k)}{f_Y(y)}, \quad f_Y(y) = \sum_{k'} p_K(k')f_{Y|K}(y|k')$$

- K : -1, +1, original signal, equally likely. $p_K(1) = 1/2, p_K(-1) = 1/2$.
- Y : measured signal with Gaussian noise, $Y = K + W, W \sim N(0, 1)$
- Your received signal = 0.7. What's your guess about the original signal?

Example: Signal Detection (1)

Inference of discrete K given continuous Y :

$$p_{K|Y}(k|y) = \frac{p_K(k)f_{Y|K}(y|k)}{f_Y(y)}, \quad f_Y(y) = \sum_{k'} p_K(k')f_{Y|K}(y|k')$$

- K : -1, +1, original signal, equally likely. $p_K(1) = 1/2, p_K(-1) = 1/2$.
- Y : measured signal with Gaussian noise, $Y = K + W, W \sim N(0, 1)$
- Your received signal = 0.7. What's your guess about the original signal?
- Your received signal = -0.2. What's your guess about the original signal?

Example: Signal Detection (1)

Inference of discrete K given continuous Y :

$$p_{K|Y}(k|y) = \frac{p_K(k)f_{Y|K}(y|k)}{f_Y(y)}, \quad f_Y(y) = \sum_{k'} p_K(k')f_{Y|K}(y|k')$$

- K : -1, +1, original signal, equally likely. $p_K(1) = 1/2, p_K(-1) = 1/2$.
- Y : measured signal with Gaussian noise, $Y = K + W, W \sim N(0, 1)$
- Your received signal = 0.7. What's your guess about the original signal? **+1**
- Your received signal = -0.2. What's your guess about the original signal? **-1**

Example: Signal Detection (2)

- $Y|K = 1 \sim N(1, 1)$ and $Y|K = -1 \sim N(-1, 1)$.

Example: Signal Detection (2)

- $Y|K = 1 \sim N(1, 1)$ and $Y|K = -1 \sim N(-1, 1)$.

$$f_{Y|K}(y|k) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(y-k)^2}, \quad k = 1, -1$$

Example: Signal Detection (2)

- $Y|K = 1 \sim N(1, 1)$ and $Y|K = -1 \sim N(-1, 1)$.

$$f_{Y|K}(y|k) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(y-k)^2}, \quad k = 1, -1$$

$$f_Y(y) = \frac{1}{2} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(y+1)^2} + \frac{1}{2} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(y-1)^2}$$

Example: Signal Detection (2)

- $Y|K = 1 \sim N(1, 1)$ and $Y|K = -1 \sim N(-1, 1)$.

$$f_{Y|K}(y|k) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(y-k)^2}, \quad k = 1, -1$$

$$f_Y(y) = \frac{1}{2} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(y+1)^2} + \frac{1}{2} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(y-1)^2}$$

- Probability that $K = 1$, given $Y = y$? After some algebra,

$$p_{K|Y}(1|y) = \frac{1}{1 + e^{-2y}}$$

Example: Signal Detection (2)


- $Y|K = 1 \sim N(1, 1)$ and $Y|K = -1 \sim N(-1, 1)$.

$$f_{Y|K}(y|k) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(y-k)^2}, \quad k = 1, -1$$

$$f_Y(y) = \frac{1}{2} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(y+1)^2} + \frac{1}{2} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(y-1)^2}$$

- Probability that $K = 1$, given $Y = y$? After some algebra,

$$p_{K|Y}(1|y) = \frac{1}{1 + e^{-2y}}$$



Questions?

- 1) What is PDF and CDF?
- 2) Why do we need CDF?
- 3) What are joint/marginal/conditional PDFs?
- 4) Explain memorylessness of exponential random variables.
- 5) Explain the version of Bayes' rule for continuous and mixed random variables.