

Lecture 3: Random Variable, Part I

Yi, Yung (이웅)

EE210: Probability and Introductory Random Processes
KAIST EE

April 19, 2021

April 19, 2021 1 / 1

Roadmap

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

April 19, 2021 2 / 1


- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

L3(1)

April 19, 2021 3 / 1

Random Variable: Idea

- In reality, many outcomes are **numerical**, e.g., stock price.
- Even if not, very convenient if we map numerical values to random outcomes, e.g., '0' for male and '1' for female.



(b) Two rolls of tetrahedral dice

L3(1)

April 19, 2021 4 / 1

- Mathematically, a random variable X is a **function** which maps from Ω to \mathbb{R} .
- Notation.** Random variable X , numerical value x .
- Different random variables can be defined on the same sample space.
- For a fixed value x , we can associate an **event** that a random variable X has the value x , i.e., $\{\omega \in \Omega \mid X(\omega) = x\}$
- Assume that values x are discrete¹ such as $1, 2, 3, \dots$.
For notational convenience,

$$p_X(x) \triangleq \mathbb{P}(X = x) \triangleq \mathbb{P}(\{\omega \in \Omega \mid X(\omega) = x\})$$
- For a discrete random variable X , we call $p_X(x)$ **probability mass function** (PMF).

¹Finite or countably infinite.

- (1) Random variable: Idea and formal definition
- (2) **Popular discrete random variables**
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

- Only **binary** values

$$X = \begin{cases} 0, & \text{w.p. } 1 - p, \\ 1, & \text{w.p. } p \end{cases}$$


In other words, $p_X(0) = 1 - p$ and $p_X(1) = p$ from our PMF notation.

- Models a trial that results in binary results, e.g., success/failure, head/tail
- Very useful for an **indicator rv** of an event A . Define a rv $\mathbf{1}_A$ as:

$$\mathbf{1}_A = \begin{cases} 1, & \text{if } A \text{ occurs,} \\ 0, & \text{otherwise} \end{cases}$$

¹w.p.: with probability
L3(2)


- integers a, b , where $a \leq b$
- Choose a number out of $\Omega = \{a, a + 1, \dots, b\}$ uniformly at random.
- $p_X(i) = \frac{1}{b-a+1}$, $i \in \Omega$



- Models complete **ignorance** (I don't know anything about X)

- Models the number of **successes** in a given number of **independent** trials
- n independent trials, where one trial has the success probability p .

$$p_X(k) = \binom{n}{k} p^k (1-p)^{n-k}$$




¹ $\binom{n}{k} = \frac{n!}{k!(n-k)!}$, which we read ' n choose k '.
L3(2)

April 19, 2021 9 / 1

- Infinitely many independent Bernoulli trials, where each trial has success probability p
- Random variable: number of trials until the **first success**.

$$p_X(k) = (1-p)^{k-1} p$$

- Models **waiting** times until something happens.



- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

L3(3)

April 19, 2021 11 / 1

- Average

Definition

$$\mathbb{E}[X] = \sum_x x p_X(x)$$

- $p_X(x)$: relative frequency of value x (trials with x /total trials)
- **Example.** Bernoulli rv with p

$$\mathbb{E}[X] = 1 \times p + 0 \times (1 - p) = p = p_X(1)$$

L3(3)

April 19, 2021 12 / 1

Not very surprising. Easy to prove using the definition.


- If $X \geq 0$, $\mathbb{E}[X] \geq 0$.
- If $a \leq X \leq b$, $a \leq \mathbb{E}[X] \leq b$.
- For a constant c , $\mathbb{E}[c] = c$.

L3(3)

April 19, 2021 13 / 1

Expectation of a function of a RV

- For a rv X , $Y = g(X)$ is also a r.v.
- $\mathbb{E}[Y] = \mathbb{E}[g(X)] = \sum_x g(x)p_X(x)$
- Compute $\mathbb{E}[Y]$ for the following:



$$4 \times (0.4 + 0.3) + 3 \times (0.1 + 0.2) = 2.8 + 0.9 = 3.7$$

Linearity of Expectation

$$\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$$

L3(3)

April 19, 2021 14 / 1

- Measures how much the **spread** of a PMF is.
- What about $\mathbb{E}[X - \mu]$, where $\mu = \mathbb{E}[X]$? Zero
- Then, what about $\mathbb{E}[(X - \mu)^2]$?

Variance, Standard Deviation

$$\text{var}[X] = \mathbb{E}[(X - \mu)^2]$$

$$\sigma_X = \sqrt{\text{var}[X]}$$

L3(3)

April 19, 2021

15 / 1

Variance: Useful Property

- $\text{var}[X] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$

$$\text{var}[X] = \mathbb{E}[X^2 - 2\mu X + \mu^2]$$

$$= \mathbb{E}[X^2] - 2\mu\mathbb{E}[X] + \mu^2 = \mathbb{E}[X^2] - \mu^2$$
- $Y = X + b$, $\text{var}[Y] = \text{var}[X]$

$$\text{var}[Y] = \mathbb{E}[(X + b)^2] - (\mathbb{E}[X + b])^2$$
- $Y = aX$, $\text{var}[Y] = a^2\text{var}[X]$

$$\text{var}[Y] = \mathbb{E}[a^2X^2] - (a\mathbb{E}[X])^2$$

Example: Variance of a Bernoulli rv (p)

$$\mu = \mathbb{E}[X] = 1 \times p + 0 \times (1 - p) = p$$

$$\mathbb{E}[X^2] = 1 \times p + 0 \times (1 - p) = p$$

$$\begin{aligned} \text{var}[X] &= \mathbb{E}[X^2] - \mu^2 = p - p^2 \\ &= p(1 - p) \end{aligned}$$

L3(3)

April 19, 2021

16 / 1

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

L3(4)

April 19, 2021 17 / 1

Joint PMF

- Joint PMF.** For two random variables X, Y , consider two events $\{X = x\}$ and $\{Y = y\}$, and

$$p_{X,Y}(x, y) \triangleq \mathbb{P}(\{X = x\} \cap \{Y = y\})$$

- $\sum_x \sum_y p_{X,Y}(x, y) = 1$

- Marginal PMF.**

$$p_X(x) = \sum_y p_{X,Y}(x, y),$$

$$p_Y(y) = \sum_x p_{X,Y}(x, y)$$

Example.

VIDEO PAUSE

	y				
	4	1/20	2/20	2/20	
	3	2/20	4/20	1/20	2/20
	2		1/20	3/20	1/20
	1		1/20		
		1	2	3	4
		x			

$$p_{X,Y}(1, 3) = 2/20$$

$$p_X(4) = 2/20 + 1/20 = 3/20$$

$$\mathbb{P}(X = Y) = 1/20 + 4/20 + 3/20 = 8/20$$

L3(4)

April 19, 2021 18 / 1

- Consider a rv $Z = g(X, Y)$. (Ex) $X + Y, X^2 + Y^2$. Then, PMF of Z is:

$$p_Z(z) = \mathbb{P}(g(X, Y) = z) = \sum_{(x,y): g(x,y)=z} p_{X,Y}(x, y)$$

- Similarly,

$$\mathbb{E}[Z] = \mathbb{E}[g(X, Y)] = \sum_x \sum_y g(x, y) p_{X,Y}(x, y)$$

L3(4)

April 19, 2021 19 / 1

Linearity of Expectation for Multiple RVs

- Remember: $\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$

- Similarly,

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

(easy to prove, using the definition.)

- $\mathbb{E}[X_1 + \dots + X_n] = \mathbb{E}[X_1] + \dots + \mathbb{E}[X_n]$
- $\mathbb{E}[2X + 3Y - Z] = 2\mathbb{E}[X] + 3\mathbb{E}[Y] - \mathbb{E}[Z]$

- Example.** Mean of a binomial rv Y with (n, p)

- Y : number of successes in n Bernoulli trials with p
- $Y = X_1 + \dots + X_n$, where X_i is a Bernoulli rv.

- $\mathbb{E}[Y] = n\mathbb{E}[X_i] = n\mathbb{P}(X_i = 1) = np$

Message. When some rv X is write as a linear combination of other rvs, it is often easy to handle X .

L3(4)

April 19, 2021 20 / 1

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) **Conditioning for random variables**
- (6) Independence for random variables

L3(5)

April 19, 2021 21 / 1

Conditional PMF: Conditioning on an event


Remember two probability laws: $\mathbb{P}(\cdot)$ and $\mathbb{P}(\cdot|A)$ for an event A .

- | | |
|--|---|
| <ul style="list-style-type: none"> • $p_X(x) \triangleq \mathbb{P}(X = x)$ • $\mathbb{E}[X] = \sum_x x p_X(x)$ • $\mathbb{E}[g(X)] = \sum_x g(x) p_X(x)$ • $\text{var}[X] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$ | <ul style="list-style-type: none"> • $p_{X A}(x) \triangleq \mathbb{P}(X = x A)$ • $\mathbb{E}[X A] \triangleq \sum_x x p_{X A}(x)$ • $\mathbb{E}[g(X) A] \triangleq \sum_x g(x) p_{X A}(x)$ • $\text{var}[X A] \triangleq \mathbb{E}[X^2 A] - (\mathbb{E}[X A])^2$ • (Note) $p_{X A}(x)$, $\mathbb{E}[X A]$, $\mathbb{E}[g(X) A]$, and $\text{var}[X A]$ are all just notations! |
|--|---|

L3(5)


April 19, 2021 22 / 1

$$A = \{X \geq 2\}$$



$$\mathbb{E}[X] = \frac{1}{4}(1 + 2 + 3 + 4) = 2.5$$

$$\begin{aligned} \text{var}[X] &= \mathbb{E}[X^2] - (\mathbb{E}[X])^2 \\ &= \frac{1}{4}(1 + 2^2 + 3^2 + 4^2) - 2.5^2 \end{aligned}$$



$$\mathbb{E}[X|A] = \frac{1}{3}(2 + 3 + 4) = 3$$

$$\begin{aligned} \text{var}[X|A] &= \mathbb{E}[X^2|A] - (\mathbb{E}[X|A])^2 \\ &= \frac{1}{3}(2^2 + 3^2 + 4^2) - 3^2 = 2/3 \end{aligned}$$

L3(5)

April 19, 2021 23 / 1

What do we mean by “conditioning on a rv”? Consider $A = \{Y = y\}$ for a rv Y .

- $p_{X|A}(x) \triangleq \mathbb{P}(X = x|A)$
- $\mathbb{E}[X|A] \triangleq \sum_x x p_{X|A}(x)$
- $\mathbb{E}[g(X)|A] \triangleq \sum_x g(x) p_{X|A}(x)$
- $\text{var}[X|A] \triangleq \mathbb{E}[X^2|A] - (\mathbb{E}[X|A])^2$
- $p_{X|Y}(x|y) \triangleq \mathbb{P}(X = x|Y = y)$
- $\mathbb{E}[X|Y = y] \triangleq \sum_x x p_{X|Y}(x|y)$
- $\mathbb{E}[g(X)|Y = y] \triangleq \sum_x g(x) p_{X|Y}(x|y)$
- $\text{var}[X|Y = y] \triangleq \mathbb{E}[X^2|Y = y] - (\mathbb{E}[X|Y = y])^2$

L3(5)

April 19, 2021 24 / 1

VIDEO PAUSE

- Conditional PMF

$$p_{X|Y}(x|y) \triangleq \mathbb{P}(X = x | Y = y) = \frac{p_{X,Y}(x, y)}{p_Y(y)}$$

for y such that $p_Y(y) > 0$.

- $\sum_x p_{X|Y}(x|y) = 1$

- Multiplication rule.

$$\begin{aligned} p_{X,Y}(x, y) &= p_Y(y)p_{X|Y}(x|y) \\ &= p_X(x)p_{Y|X}(y|x) \end{aligned}$$

- $p_{X,Y,Z}(x, y, z) = p_X(x)p_{Y|X}(y|x)p_{Z|X,Y}(z|x, y)$

y				
4	1/20	2/20	2/20	
3	2/20	4/20	1/20	2/20
2		1/20	3/20	1/20
1		1/20		
	1	2	3	4
	x			

$$p_{X|Y}(2|2) = \frac{1}{1+3+1}$$

$$p_{X|Y}(3|2) = \frac{3}{1+3+1}$$

$$\mathbb{E}[X|Y = 3] = 1(2/9) + 2(4/9) + 3(1/9) + 4(2/9)$$

L3(5)



April 19, 2021 25 / 1

Remind: Total Probability Theorem (from Lecture 2)

- Partition of Ω into A_1, A_2, A_3
- Known: $\mathbb{P}(A_i)$ and $\mathbb{P}(B|A_i)$
- What is $\mathbb{P}(B)$? (probability of result)

Total Probability Theorem

$$\mathbb{P}(B) = \sum_i \mathbb{P}(A_i)\mathbb{P}(B|A_i)$$



L3(5)



April 19, 2021 26 / 1

Total Probability Theorem: $B = \{X = x\}$

- Partition of Ω into A_1, A_2, A_3

Total Probability Theorem

$$p_X(x) = \sum_i \mathbb{P}(A_i) \mathbb{P}(X = x | A_i) = \sum_i \mathbb{P}(A_i) p_{X|A_i}(x)$$



Total Expectation Theorem for $\{A_i\}$



- Partition of Ω into A_1, A_2, A_3

Total Probability Theorem

$$p_X(x) = \sum_i \mathbb{P}(A_i) \mathbb{P}(X = x | A_i) = \sum_i \mathbb{P}(A_i) p_{X|A_i}(x)$$

Total Expectation Theorem

$$\mathbb{E}[X] = \sum_i \mathbb{P}(A_i) \mathbb{E}[X | A_i]$$




- Partition of Ω into A_1, A_2, A_3

Total Expectation Theorem

$$\mathbb{E}[X] = \sum_i \mathbb{P}(A_i) \mathbb{E}[X|A_i]$$

Total Expectation Theorem

$$\mathbb{E}[X] = \sum_y \mathbb{P}(Y = y) \mathbb{E}[X|Y = y] = \sum_y p_Y(y) \mathbb{E}[X|Y = y]$$



L3(5)

April 19, 2021 29 / 1

Example 1: Total Expectation Theorem


- Using the definition of expectation,

$$\begin{aligned} \mathbb{E}[X] &= \frac{1}{9}(0 + 1 + 2) + \frac{2}{9}(6 + 7 + 8) \\ &= \frac{3 + 12 + 14 + 16}{9} = 5 \end{aligned}$$

- Let's use TET, for which consider

$$A_1 = \{X \in \{0, 1, 2\}\}, A_2 = \{X \in \{6, 7, 8\}\}$$

$$\begin{aligned} \mathbb{E}[X] &= \sum_{i=1,2} \mathbb{P}(A_i) \mathbb{E}[X|A_i] \\ &= \frac{1}{3} \cdot 1 + \frac{2}{3} \cdot 7 = 5 \end{aligned}$$



L3(5)

April 19, 2021 30 / 1

- Some random variable often does not have **memory**.
- Definition.** A random variable X is called **memoryless** if, for any $n, m \geq 0$,

$$\mathbb{P}(X > n + m | X > m) = \mathbb{P}(X > n)$$

- Meaning.** Conditioned on $X > m$, $X - m$'s distribution is the same as the original X .

$$\mathbb{P}(X - m > n | X > m) = \mathbb{P}(X > n)$$

- Suppose that X is the time of waiting for a bus and X is memoryless. At the bus stop, I have waited for the bus for 10 mins. Then, the time until the bus arrival does not depend on how much I have waited for a bus. **No memory.**

- Theorem.** Any geometric random variable is **memoryless**.
- Remind.** Geometric rv X with parameter p

$$\mathbb{P}(X = k) = (1 - p)^{k-1}p, \quad \mathbb{P}(X > k) = 1 - \sum_{k'=1}^k (1 - p)^{k'-1}p = (1 - p)^k$$

- Proof.**

$$\begin{aligned} \mathbb{P}(X > n + m | X > m) &= \frac{\mathbb{P}(X > n + m \text{ and } X > m)}{\mathbb{P}(X > m)} = \frac{\mathbb{P}(X > n + m)}{\mathbb{P}(X > m)} \\ &= \frac{(1 - p)^{n+m}}{(1 - p)^m} = (1 - p)^n = \mathbb{P}(X > n) \end{aligned}$$

- Meaning.** Conditioned on $X > m$, $X - m$ is geometric with the same parameter.

Example 2: Mean of Geometric rv

- Write softwares over and over, and each time w.p. p of working correctly (independent from prev. programs).
- X : number of trials until the program works correctly.
- (Q) mean of X
- X is geometric
- Direct computation is boring.

$$\mathbb{E}[X] = \sum_{k=1}^{\infty} k(1-p)^{k-1}p$$

- Total expectation theorem and memorylessness helps a lot.

- $A_1 = \{X = 1\}$ (first try is success),
 $A_2 = \{X > 1\}$ (first try is failure).

$$\begin{aligned}\mathbb{E}[X] &= 1 + \mathbb{E}[X - 1] \\ &= 1 + \mathbb{P}(A_1)\mathbb{E}[X - 1|X = 1] \\ &\quad + \mathbb{P}(A_2)\mathbb{E}[X - 1|X > 1] \\ &= 1 + (1 - p)\mathbb{E}[X]\end{aligned}$$

$$\mathbb{E}[X] = 1 + (1 - p)\frac{1}{p} = 1/p.$$

Roadmap

- (1) Random variable: Idea and formal definition
- (2) Popular discrete random variables
- (3) Summarizing random variables: Expectation and Variance
- (4) (Functions of) multiple random variables
- (5) Conditioning for random variables
- (6) Independence for random variables

- Two events

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$$

$$\mathbb{P}(A \cap B | C) = \mathbb{P}(A | C) \cdot \mathbb{P}(B | C)$$

- A rv and an event

$$\mathbb{P}(\{X = x\} \cap B) = \mathbb{P}(X = x) \cdot \mathbb{P}(B), \quad \text{for all } x$$

$$\mathbb{P}(\{X = x\} \cap B | C) = \mathbb{P}(X = x | C) \cdot \mathbb{P}(B | C), \quad \text{for all } x$$

- Two rvs

$$\mathbb{P}(\{X = x\} \cap \{Y = y\}) = \mathbb{P}(X = x) \cdot \mathbb{P}(Y = y), \quad \text{for all } x, y$$

$$p_{X,Y}(x, y) = p_X(x) \cdot p_Y(y)$$

$$\mathbb{P}(\{X = x\} \cap \{Y = y\} | Z = z) = \mathbb{P}(X = x | Z = z) \cdot \mathbb{P}(Y = y | Z = z), \quad \text{for all } x, y$$

$$p_{X,Y|Z}(x, y) = p_{X|Z}(x) \cdot p_{Y|Z}(y)$$

L3(6)

April 19, 2021 35 / 1

Example


- $X \perp\!\!\!\perp Y$?

$$p_{X,Y}(1, 1) = 0, \quad p_X(1) = 3/20$$

$$p_Y(1) = 1/20.$$

- $X \perp\!\!\!\perp Y | \{X \leq 2 \text{ and } Y \geq 3\}$?

- Yes.



$Y = 4 \ (1/3)$	$1/9$	$2/9$
$Y = 3 \ (2/3)$	$2/9$	$4/9$
	$X = 1 \ (1/3)$	$X = 2 \ (2/3)$

L3(6)

April 19, 2021 36 / 1

- Always true.
 $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$
- Generally, $\mathbb{E}[g(X, Y)] \neq g(\mathbb{E}[X], \mathbb{E}[Y])$
- However, if $X \perp\!\!\!\perp Y$,

$$\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$$

$$\mathbb{E}[g(X)h(Y)] = \mathbb{E}[g(X)]\mathbb{E}[h(Y)]$$

- **Proof.**

$$\begin{aligned}\mathbb{E}[g(X)h(Y)] &= \sum_x \sum_y g(x)h(y)p_{X,Y}(x, y) \\ &= \sum_x x p_X(x) \sum_y y p_Y(y)\end{aligned}$$

- Always true.
 $\text{var}[aX] = a^2\text{var}[X], \text{var}[X + a] = \text{var}[X]$
- Generally, $\text{var}[X + Y] \neq \text{var}[X] + \text{var}[Y]$
- However, if $X \perp\!\!\!\perp Y$,
 $\text{var}[X + Y] = \text{var}[X] + \text{var}[Y]$
- Practice.
 - $X = Y \implies \text{var}[X + Y] = 4\text{var}[X]$
 - $X = -Y \implies \text{var}[X + Y] = 0$
 - $X \perp\!\!\!\perp Y \implies \text{var}[X - 3Y] = \text{var}[X] + 9\text{var}[Y]$

L3(6)

April 19, 2021 37 / 1

$\text{var}[X + Y] \neq \text{var}[X] + \text{var}[Y]$

- Why not generally true?

$$\begin{aligned}\text{var}[X + Y] &= \mathbb{E}[(X + Y)^2] - (\mathbb{E}[X + Y])^2 \\ &= \mathbb{E}[X^2 + Y^2 + 2XY] - ((\mathbb{E}[X])^2 + (\mathbb{E}[Y])^2 + 2\mathbb{E}[X]\mathbb{E}[Y]) \\ &= \text{var}[X] + \text{var}[Y] + 2(\mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y])\end{aligned}$$

- $X \perp\!\!\!\perp Y$ is a sufficient condition for $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$
- Also, a necessary condition? we will see later, when we study **covariance**.

L3(6)

April 19, 2021 38 / 1

- n people throw their hats in a box and then pick one at random
- X : number of people with their own hat
- $\mathbb{E}[X]$? $\text{var}[X]$?
- All permutations are equally likely as $1/n!$. Thus, this equals to picking one hat at a time.
- **Key step 1.** Define a rv $X_i = 1$ if i selects own hat and 0 otherwise.

$$X = \sum_{i=1}^n X_i.$$

- $\{X_i\}, i = 1, 2, \dots, n$: identically distributed (symmetry)

- $\mathbb{E}[X] = n\mathbb{E}[X_1] = n\mathbb{P}(X_1 = 1) = n \times \frac{1}{n} = 1$.
- **Key step 2.** Are X_i s are independent? If yes, easy to get $\text{var}(X)$.
- Assume $n = 2$. Then, $X_1 = 1 \rightarrow X_2 = 1$, and $X_1 = 0 \rightarrow X_2 = 0$. Thus, **dependent**.

$$\begin{aligned} \text{var}(X) &= \mathbb{E}[X^2] - (\mathbb{E}[X])^2 \\ &= \mathbb{E}\left[\sum_i X_i^2 + \sum_{i,j:i \neq j} X_i X_j\right] - (\mathbb{E}[X])^2 \end{aligned}$$

$$\mathbb{E}[X_i^2] = 1 \times \frac{1}{n} + 0 \times \frac{n-1}{n} = \frac{1}{n}$$

$$\mathbb{E}[X_i X_j] = \mathbb{E}[X_1 X_2] = 1 \times \mathbb{P}(X_1 X_2 = 1) = \mathbb{P}(X_1 = 1)\mathbb{P}(X_2 = 1|X_1 = 1), \quad (i \neq j)$$

- $\mathbb{E}[X^2] = n\mathbb{E}[X_1^2] + n(n-1)\mathbb{E}[X_1 X_2] = n\frac{1}{n} + n(n-1)\frac{1}{n(n-1)} = 2$
- $\text{var}(X) = 2 - 1 = 1$

Questions?

L3(6)

April 19, 2021 41 / 1

Review Questions

- 1) What is Random Variable? Why is it useful?
- 2) What is PMF (Probability Mass Function)?
- 3) Explain Bernoulli, Binomial, Poisson, Geometric rvs, when they are used and what their PMFs are.
- 4) What are joint and marginal PMFS?
- 5) Describe and explain the total probability/expectation theorem for random variables?
- 6) When is it useful to use total probability/expectation theorem?
- 7) What is conditional independence?

L3(6)

April 19, 2021 42 / 1