

## Lecture 1: Probabilistic Model

Yi, Yung (이웅)

EE210: Probability and Introductory Random Processes  
KAIST EE

April 19, 2021

- (1) Probabilistic Model
  - Mathematical description of uncertain situations
- (2) Sample Space, Event, Probability Law
  - Elements of probability theory
- (3) Probability Axioms
  - 3 axioms for the completeness of a theory

April 19, 2021 1 / 1

April 19, 2021 2 / 1

- (1) Probabilistic Model
- (2) Sample Space, Event, Probability Law
- (3) Probability Axioms

**Modeling:** Understand reality with a simple (mathematical) model

- Experiment
  - Flip two coins
- Observation: a random outcome
  - for example,  $(H, H)$
- All outcomes
  - $\{(H, H), (H, T), (T, H), (T, T)\}$

- 
- **Our goal:** Build up a **probabilistic model** for an experiment with random outcomes
  - **Probabilistic model?**
    - Assign a number to each outcome or a set of outcomes
    - Mathematical description of an uncertain situation
  - Which model is good or bad?

**Goal:** Build up a probabilistic model. Hmm... How?

The first thing: What are the *elements* of a probabilistic model?

### Elements of Probabilistic Model

1. All outcomes of my interest: **Sample Space  $\Omega$**
2. Assigned numbers to each outcome of  $\Omega$ : **Probability Law  $\mathbb{P}(\cdot)$**

**Question:** What are the conditions of  $\Omega$  and  $\mathbb{P}(\cdot)$  under which their induced probability model becomes "legitimate"?

- (1) Probabilistic Model
- (2) **Sample Space, Event, Probability Law**
- (3) Probability Axioms

L1(1)

April 19, 2021 5 / 1

L1(2)

April 19, 2021 6 / 1

## 1. Sample Space $\Omega$

### Examples: Sample Space $\Omega$


The set of all outcomes of **my interest**

- (1) Mutually exclusive
- (2) Collectively exhaustive
- (3) At the **right granularity**  
(not too concrete, not too abstract)

1. Toss a coin. What about this?  
 $\Omega = \{H, T, HT\}$
2. Toss a coin. What about this?  $\Omega = \{H\}$
3. (a) Just figuring out prob. of H or T.  
 $\Rightarrow \Omega = \{H, T\}$   
  
(b) The impact of the weather (rain or no rain) on the coin's behavior.  
  
 $\Rightarrow \Omega = \{(H, R), (T, R), (H, NR), (T, NR)\},$   
  
R(Rain), NR(No Rain).


- o **Discrete case:** Two rolls of a tetrahedral die

$$\Omega = \{(1, 1), (1, 2), \dots, (4, 4)\}$$



- o **Continuous case:** Dropping a needle in a plain

$$\Omega = \{(x, y) \in \mathbb{R}^2 \mid 0 \leq x, y \leq 1\}$$



L1(2)

April 19, 2021 7 / 1

L1(2)

April 19, 2021 8 / 1

- Assign numbers to what? Each outcome?
- What is the probability of dropping a needle at  $(0.5, 0.5)$  over the  $1 \times 1$  plane?
- Assign numbers to each **subset** of  $\Omega$
- a subset of  $\Omega$ : **an event**
- $\mathbb{P}(A)$ : Probability of an event  $A$ .
  - This is where probability meets set theory.
- Roll a dice. What is the probability of odd numbers?  
 $\mathbb{P}(\{1, 3, 5\})$ , where  $\{1, 3, 5\} \subset \Omega$  is an event.

- (1) Probabilistic Model
- (2) Sample Space, Event, Probability Law
- (3) **Probability Axioms**

L1(2)

April 19, 2021 9 / 1

L1(3)

April 19, 2021 10 / 1

How should we construct  $\mathbb{P}(\cdot)$ ?

## Probability Axioms

- Need to construct  $\mathbb{P}(\cdot)$  that naturally satisfies the intention of a probability theory designer just like you. What about the followings as starting points?
  - $\mathbb{P}(A) \geq 0$  for any event  $A \subset \Omega$
  - $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$
  - $\mathbb{P}(A \cup B) \leq \mathbb{P}(A) + \mathbb{P}(B)$
  - For two disjoint<sup>1</sup> events  $A$  and  $B$ ,  $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B)$
  - $\mathbb{P}(\Omega) = 1$  (Why not  $\mathbb{P}(\Omega) = 10$ ?)
  - $\mathbb{P}(\emptyset) = 0$
  - If  $A \subset B$ ,  $\mathbb{P}(A) \leq \mathbb{P}(B)$
  - many others

<sup>1</sup>Their intersection is empty.

L1(3)

April 19, 2021 11 / 1

L1(3)

April 19, 2021 12 / 1

- Surprisingly, we need just the following three rules (called **axioms**):

## Probability Axioms: Version 1

A1. **Nonnegativity**:  $\mathbb{P}(A) \geq 0$  for any event  $A \subset \Omega$

A2. **Normalization**:  $\mathbb{P}(\Omega) = 1$

A3. **(Finite) additivity**: For two disjoint events  $A$  and  $B$ ,  $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B)$

- No other things are necessary, and we can prove all other things from the above axioms.
- Note that coming up with the above axioms is far from trivial.

A1: Nonnegativity, A2: Normalization, A3: Finite additivity

Prove the following properties using the axioms:

VIDEO PAUSE

1. For any event  $A$ ,  $\mathbb{P}(A) \leq 1$

$$1 \stackrel{A2}{=} \mathbb{P}(\Omega) = \mathbb{P}(A \cup A^c) \stackrel{A3}{=} \mathbb{P}(A) + \mathbb{P}(A^c) \stackrel{A1}{\implies} \mathbb{P}(A) = 1 - \mathbb{P}(A^c) \leq 1$$

2.  $\mathbb{P}(\emptyset) = 0$

$$\mathbb{P}(\Omega \cup \emptyset) \stackrel{A3}{=} \mathbb{P}(\Omega) + \mathbb{P}(\emptyset) \stackrel{A2}{=} 1 + \mathbb{P}(\emptyset) \stackrel{\text{from 1.}}{\implies} \mathbb{P}(\emptyset) = 0$$

3. If  $A \subset B$ ,  $\mathbb{P}(A) \leq \mathbb{P}(B)$

$$\mathbb{P}(B) \stackrel{A3}{=} \mathbb{P}(A) + \mathbb{P}(B \setminus A) \stackrel{A1}{\geq} \mathbb{P}(A)$$

1. Specify the sample space
2. Specify a probability law  
- from my earlier belief, from data, from expert's opinion
3. Identify an event of interest
4. Calculate

Toss a (biased) coin

1.  $\Omega = \{H, T\}$
2.  $\mathbb{P}(\{H\}) = 1/4$ ,  $\mathbb{P}(\{T\}) = 3/4$ ,
3. probability of head or tail
4.  $1/4$ ,  $3/4$

- $\Omega = \{1, 2, 3, \dots\}$ ,  $\mathbb{P}(\{n\}) = \frac{1}{2^n}$ ,  $n = 1, 2, \dots$
- Is the above probability law legitimate? seems OK

$$\mathbb{P}(\Omega) = \frac{1}{2} + \frac{1}{2^2} + \dots = \frac{1/2}{1 - 1/2} = 1$$

- $\mathbb{P}(\text{even numbers})?$

$$\begin{aligned} \mathbb{P}(\text{even}) &= \mathbb{P}(\{2, 4, 6, \dots\}) \\ &= \frac{1}{2^2} + \frac{1}{2^4} + \frac{1}{2^6} + \dots = 1/3 \end{aligned}$$

- Is the above right? If not, why?
  - Wrong: Finite additivity axiom does not allow this.

#### Probability Axioms: Version 1 2

- A1. Nonnegativity:  $\mathbb{P}(A) \geq 0$  for any event  $A \subset \Omega$
- A2. Normalization:  $\mathbb{P}(\Omega) = 1$
- A3. (Finite) additivity: For two disjoint events  $A$  and  $B$ ,  $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B)$
- A4. Countable additivity: If  $A_1, A_2, A_3, \dots$  is an infinite sequence of disjoint events, then  $\mathbb{P}(A_1 \cup A_2 \cup \dots) = \mathbb{P}(A_1) + \mathbb{P}(A_2) + \dots$ .

- A narrow view: A branch of math
  - axioms  $\rightarrow$  theorems
  - Mathematicians work very hard to find the smallest set of necessary axioms (just like atoms in physics)
- Frequencies:  $\mathbb{P}(H) = 1/2$ 
  - Understanding an uncertain situation: fractions of successes out of many experiments
- Beliefs:  $\mathbb{P}(\text{He is reelected}) = 0.7$

Anyway, we believe that probabilistic reasoning is very helpful to understand the world with many uncertain situations.

## Questions?

L1(3)

April 19, 2021 17 / 1

L1(3)

April 19, 2021 18 / 1

You build up the very basics of a probabilistic model.

What else do we need to build up?

- 1) Please explain what a probabilistic model is and why we need it.
- 2) What is the mathematical definition of event?
- 3) What are the key elements of the probabilistic model?
- 4) Please list up the probability axioms and explain them.
- 5) Why do we need countable additivity in the probability axioms?

L1(3)

April 19, 2021 19 / 1

L1(3)

April 19, 2021 20 / 1