Does the length of an outburst predict the wait for the next? A preregistered forecast test for SS Cygni

Draft for review, 9 November 2026. Release and submission are Barry W. Brook's decisions.

Barry W. Brook*[1], Claude Opus 5.5 (Anthropic) and GPT-6 Codex (OpenAI)[2]

Draft for review; release and submission are Barry W. Brook's decision.

1. University of Tasmania, Hobart, Tasmania, Australia
2. AI research collaborators; roles are stated in "Roles of AI systems and humans"

Corresponding author: Barry W. Brook

Abstract

Visual observers have recorded the outbursts of the dwarf nova SS Cygni almost continuously since 1896. Earlier analyses reported that longer outbursts tend to be followed by longer quiet intervals, but those were in-sample correlations, and the best-known version partly reflected how cycles were defined. We asked a narrower forecasting question. Once an outburst has ended, does its duration improve a probabilistic forecast of the next onset, beyond what the recent quiet intervals already say? We extracted outbursts from archived visual estimates with a threshold-crossing method qualified on synthetic histories. We fitted lognormal forecast clocks to 1910–1959 only, froze them, published the protocol with a DOI, and then scored observation-time retrospective forecasts issued when the recorded observations confirmed each outburst had ended, from 1960 to 2017. Adding duration improved the log score by 0.0165 nats per forecast (90% cluster interval 0.0049 to 0.0281; 441 forecasts). The predeclared criterion was met. The gain corresponds to a 1.7% higher geometric mean likelihood of the observed outcomes. The evaluation was completed after a documented post-access repair to the parser, for a missing-uncertainty token that no forecast uses. Two predeclared secondary comparisons were inconclusive. One was the later segment, 1993–2017 (0.0057; −0.0107 to 0.0221). The other was a rule built on the 12-day long/short outburst classes (0.0025; −0.0022 to 0.0072). Exploratory fits made after the result favour a long/short class indicator over continuous duration. They also show that the class rule's score is sensitive to a shared level adjustment against the fixed baseline. These fits supply no further confirmation. The result establishes incremental predictive information in the completed outburst for one star, under one extraction and one timing baseline. It does not establish physical memory, identify a mechanism, or carry over to other dwarf novae.

1. Introduction

Dwarf novae are close binaries in which a white dwarf accretes from a low-mass companion through a disc. Every few weeks to months the disc brightens by several magnitudes for days to weeks and then fades again. The disc instability model explains these outbursts as a limit cycle. Hydrogen ionisation makes the disc thermally unstable, so it switches between a cool low-viscosity state that stores mass and a hot high-viscosity state that drains it (Lasota 2001; Hameury 2020). Recurrence is quasi-periodic, not periodic. Whether a completed outburst improves a next-onset forecast is a measurable predictive question; relating any gain to the surviving disc state requires a separate physical model.

SS Cygni is the natural test case. It is the brightest dwarf nova in the northern sky. Its light curve, built from visual estimates archived by the American Association of Variable Star Observers (AAVSO), runs almost unbroken from 1896. Bath and van Paradijs described SS Cyg as a relaxation oscillator, with outburst energy related to the following waiting period (Bath and van Paradijs 1983). Cannizzo and Mattei (CM92) then analysed 705 outbursts from 1896 to 1992 (Cannizzo and Mattei 1992). They showed that a correlation between an outburst's duration and the "following cycle" largely reflects the cycle containing the duration. They recommended the quiet interval instead.

On that footing CM92 reported the following correlations:

- duration with the following quiescence, r = 0.223;
- duration with the preceding quiescence, 0.282;
- successive durations, −0.400, a strong alternation of long and short outbursts;
- durations split cleanly into two classes at about 12 days.

They read the positive correlation between successive quiet intervals as a hint of secular variation. A 1990 study by Hempelmann and Kurths (*Astronomy and Astrophysics* 232, 356) found short-range dependence and possible long-range memory in a symbolic sequence of outburst widths. Searches for low-dimensional chaos found no attractor (Cannizzo and Goodings 1988). Cannizzo's disc-instability calculations tuned to SS Cyg reproduce the alternation (Cannizzo 1993). In those calculations a long outburst drains more of the disc than a short one, and the depleted outer disc tends to make the next outburst short. Moving averages of the real light curve also showed longer recurrence times where long outbursts were relatively more common. The calculations reproduce that relation through slow changes in disc parameters, not through changes in the mass-transfer rate.

All of this work describes associations within the record it analysed. None of it asks whether the duration of a

completed outburst improves forecasts of the next onset on data not used to build the forecast. That question differs from the correlations. A forecast may use only information available when it is issued. It must beat a baseline that already knows the recent rhythm of quiet intervals. A proper scoring rule must reward it for the whole predictive distribution, not for a point estimate (Gneiting and Raftery 2007). Evaluation must also respect time order (Bürkner, Gabry, and Vehtari 2020). The only held-out timing forecast of a dwarf nova we found is a machine-learning study of the Kepler light curve of V363 Lyr by Grigorievskiy and colleagues, in the 2014 ITISE proceedings. It scored point forecasts of onset position on a single test outburst. A search for short-term photometric precursors of SS Cyg outbursts found none (Price et al. 2007).

We report a preregistered, frozen-parameter test of one question. Does a forecast that knows the completed outburst's duration assign higher probability to the next extracted onset than one that knows only the recent quiet intervals? To our knowledge this is the first preregistered probabilistic evaluation of dwarf-nova recurrence forecasts on held-out data, and the first held-out test of whether the completed outburst's duration adds forecast skill beyond a specified timing-history baseline. Our bounded search included a citation screen, with gaps in ADS access and some full texts (Supplement S3.8). Section 2 sets out the evidence and design, Section 3 the results, and Section 4 what they do and do not support. The supplement holds the definitions, methods, sensitivities and the record of corrections made during review.

2. Evidence

2.1 Observations and outburst extraction

We used AAVSO International Database visual estimates of SS Cyg, keeping only rows in the `Vis.` band with validation flag `V`. The 1960–2017 evaluation window contains 249,356 such rows, out of 500,479 deduplicated archive rows. Earlier rows supplied the development fit (1910–1959) and the causal history before 1960. Raw observations and observer identities are not redistributed.

Outbursts were extracted by a threshold-crossing method, X4, at visual magnitude 10, CM92's threshold (Supplement S2.1). For each candidate episode, X4 forms a posterior for the times at which the brightening and fading curves cross the threshold. It includes a shared systematic error so that the crossing intervals do not narrow artificially when many observers report. An episode is confirmed by two observers, or by one observer bright on two nights, once well past the threshold, with no contradicting faint report. Before any real-star forecast was scored, X4 and a coverage rule, C1/7 d, were qualified on synthetic histories with realistic observer cadence. The coverage rule treats a gap of more than seven days between visits as a loss of coverage. The qualification was conditional: the unseen waveform family failed benchmark qualification before X4 recovery was scored, and diagnostics at magnitude 9.5 failed in later eras. The magnitude-10 extraction is therefore part of the estimand, not a neutral measurement.

2.2 Forecasts, outcomes and models

Each observation-time confirmation that an outburst has ended issues one forecast. Its predictors come only from the 400 days of observations before that moment. A test with deliberately poisoned future rows enforces this. The predictors are the following:

- **duration:** the completed outburst's onset-to-end time, from the prefix extraction;
- **previous gap:** the preceding quiet interval, if covered;
- **recent level:** the mean of up to six earlier covered log quiet intervals.

This is an observation-time retrospective comparison: we use reported dates, while the historical upload and revision times needed for an operational replay are unknown. Both timing predictors are standardised using the frozen development centres and scales; an unavailable recent level is set to zero after standardisation (Supplement S2.4).

The outcome is the next extracted onset bracket. The bracket comes from the full record up to the end of the block, and serves only for scoring. A forecast is right-censored at the earliest of coverage loss, the next forecast's issue time, or the end of the block.

Each model is a lognormal clock for the quiet interval, conditioned on survival to the age at issue:

- **M0:** a constant;
- **M2:** M0 plus the standardised log previous gap and the recent level;
- **M3:** M2 plus β times the standardised log duration, with β constrained to be non-negative.

We fitted M2 on 1910–1959, then fitted β with M2's parameters held fixed, and froze all parameters. The development fit gave $\beta = 0.0466$ per standard deviation of log duration. Forecasts without a covered previous gap use M0 for both models and contribute exactly zero to the contrast.

2.3 The primary test and the access record

The single confirmatory statistic is the mean paired log-score gain of M3 over M2 across all score-eligible 1960–2017 issuances, including the fallback zeros. Its uncertainty comes from consecutive blocks of six times 365.25 days, a small-sample-corrected cluster-robust standard error and Student's t with nine degrees of freedom. The result counts as confirmed if the one-sided 95% lower bound exceeds zero. The protocol named the era split, the long/short class rule, several sensitivity variants and calibration diagnostics as secondary analyses that cannot replace the primary.

The protocol was published with a DOI, 10.5281/zenodo.24053338, minutes before the first request for evaluation data. That first attempt stopped before any ledger or score was computed. The archive returns an uncertainty field as the literal token `None` for missing values, which the parser rejected. The uncertainty field enters no extraction, feature, routing or score. An independently reviewed repair accepted that exact token without altering or dropping any observation. Synthetic invariance tests, the full test suite and a new rehearsal passed before the run continued. No magnitude, outcome, count or score informed the repair. The failed attempt is preserved. The evaluation therefore cannot be described as untouched, and the DOI timestamps the original protocol, not the repair. Data from 2018 onward remain closed.

Table 1. Pivotal inputs, what each shows directly, and what a stronger claim would also need.

Input	Shows directly	Used for	A stronger claim would need
AAVSO visual estimates, 1910–2017	Brightness reports by many observers	All events, predictors and outcomes	Independent photometry or a second extraction
X4 extraction at magnitude 10, C1/7 d	Onset and end brackets, coverage	Durations, gaps, outcome brackets	Qualification on unseen waveforms and other thresholds
Frozen M0, M2, M3 (1910–1959 fit)	Forecast distributions	The scored comparison	Comparison with stronger latent-state baselines
Protocol DOI and access record	Order of publication and data access	Status of the test	A fully untouched evaluation, now impossible for 1960–2017
CM92 Table 4	Published in-sample correlations, 1896–1992	Prior direction and expected size	Not applicable: prior art, not evidence for the result

3. Results

3.1 The primary comparison

The predeclared criterion was met (Fig. 2). Across 441 issuances, adding the completed outburst's duration raised the mean log score by 0.0165 nats per issuance, with 90% cluster interval 0.0049 to 0.0281. Exponentiated, this is a 1.7% higher geometric mean likelihood of the scored outcomes. It is neither an accuracy percentage nor an arithmetic probability gain. Timing models were used on 436 issuances and the shared fallback on 5. The outcomes comprised 435 onset brackets and 6 right-censored cases. The baseline itself is not trivial: M2 beat the constant M0 by 0.117 nats per issuance (0.066 to 0.167). The duration gain is about 14% of that baseline gain, a ratio that should not be read as variance explained.

8 of the ten blocks have positive means (Fig. 2A). Leaving out any one block gives means between 0.0132 and 0.0194. With the estimate and critical value held fixed, the strict failure boundary is approximately 1.42705 times the reported standard error; an exact 43% increase fails. This arithmetic does not validate the cluster dependence assumption. Every protocol sensitivity kept a positive lower bound. That includes uniform end uncertainty (0.0166, lower bound 0.0044), the previous-gap-only baseline (0.0170, 0.0054), a two-day issue delay (0.0165, 0.0049), the joint-valid rows (0.0167, 0.0049) and three-year clusters (lower bound 0.0059). These probe the same comparison and are not further confirmations.

Fig. 1 shows one forecast, selected by observation time rather than by score. It illustrates the information available at issue.

3.2 Predeclared qualifications

The two era segments differ in estimate but not decisively (Fig. 2B). The segment 1960–1992 gave 0.0240 (0.0091 to 0.0389; 261 issuances). These years were held out of the fit, but they overlap the record behind CM92's published association. The segment 1993–2017, outside the published sign analyses we know of, gave 0.0057 (−0.0107 to 0.0221; 180 issuances). That is inconclusive. The intervals overlap, no era-difference test was specified, and the result is not evidence that the gain disappeared.

The predeclared class rule used a 12-day long/short indicator plus a within-class duration term. It gained 0.0025 (−0.0022 to 0.0072), also inconclusive. Section 3.4 examines sensitivity to its coding and forecast level.

Calibration is a separate question from relative skill. The realised sample overcovers (Fig. 3). M3's nominal 50% central intervals contained between 0.55 and 0.62 of outcomes. Its nominal 90% intervals contained between 0.92 and 0.98. These are partial-identification bounds given the extracted brackets and censoring, not confidence intervals for a population coverage rate. The positive paired gain does not imply sharply calibrated operational forecasts.

3.3 Exploratory contrasts with a simulation reference and a shifted level

The development fit was in-sample and modest: a likelihood-ratio statistic of 5.19 against M2. In the operating-characteristic simulations, the reference injected effect gave a mean gain of 0.0045 nats, and at the published-size effect 11 of 40 simulated evaluations confirmed. The realised gain is larger than this model-specific reference. The reference carries Monte Carlo uncertainty and was not a target, so this is not a claim that the effect exceeds a known physical size.

A post-result decomposition, exploratory and conditional on evaluation-fitted adjustments, describes a level component without identifying its cause. Evaluation-era durations and gaps are both shorter than in 1910–1959. The best common log-location shift for M2 is −0.0468. That shift reduces the untruncated clock scale by about 4.6%; it is not an estimate of bias in the remaining-wait forecasts. Because mean standardised log duration is negative in the evaluation era, the duration term partly supplies this level correction. That accounts for 0.0037 nats (0.0001 to 0.0073). After the duration is recentred at its evaluation mean, the score contrast is 0.0128 (0.0036 to 0.0221). A single global mean does not remove drift within the period, and the two parts are a descriptive, approximately additive split, not a causal partition. The intervals condition on the fitted centre and level.

3.4 Exploratory: long/short class and alternation

All results in this subsection were computed after the primary result, using the evaluation outcomes, and are labelled exploratory (Fig. 4). With M2's historical parameters fixed and a free common level, a long/short indicator alone reaches a likelihood-ratio statistic of 14.7 against the level-only model. Continuous log duration reaches 11.7. These equal-dimensional models are not nested. Adding linear within-class duration to the indicator changes the statistic by 0.17. The fit therefore favours a parsimonious class reading. It is not a test showing that duration within a class adds nothing. In the development years the two forms fit about equally well. The fitted class contrast corresponds to a 14% larger untruncated clock scale after long outbursts, conditional on the other covariates. This is a model-scale ratio, not an observed difference in mean waiting

times. Era-specific refits give larger class contrasts in 1960–1992 than in 1993–2017, without an era-change test.

The predeclared class rule coded the class as an uncentred 0/1 indicator with a non-negative coefficient, added to a baseline whose level was fixed. The class component could therefore only lengthen forecasts after long outbursts, never shorten them after short ones. The rule's centred within-class term could still shorten some short-outburst forecasts. When both rules receive M2's evaluation-fitted common level, the frozen class rule gains about 0.009 nats (0.0041 to 0.0140). That shows sensitivity to coding and baseline calibration. It neither rescues the original inconclusive result nor validates a corrected rule. The lesson for future protocols is that indicator coding and intercept freedom must be specified together.

Alternation is strong in the forecast record; these pairs include one repeat-flagged issuance, rather than forming an independent unique-outburst catalogue. The lag-one correlation of log duration is -0.52. A long outburst followed a long one in 0.15 of pairs and followed a short one in 0.66. Fitted separately on the evaluation data, with evaluation M2 refitted and fixed, the current duration offset is positive (0.042) and the previous outburst's is negative (-0.038). Fitted jointly, both conditional support intervals include zero. These working-likelihood ranges hold the refitted M2 fixed and do not propagate baseline, dependence or measurement uncertainty. The negative previous-duration coefficient is compatible with alternation combined with a positive current association. Its causes have not been separated.

4. Discussion

4.1 What follows

For SS Cyg, the completed outburst carries usable information about when the next will begin. That information goes beyond the recent quiet intervals that a timing-only forecaster already knows, and it survived a frozen, preregistered test on 58 years of held-out record. CM92's in-sample association thus has a forecast counterpart. The 1.7% geometric likelihood gain quantifies incremental skill under this comparison. Its value for telescope scheduling and its relationship to a physical disc state were not measured.

Two physical readings are compatible with the result and we cannot separate them. In the cycle-to-cycle reading, a long outburst drains more of the disc, and the drained disc both alternates the next outburst's type and alters the interval before it. Cannizzo's model provides the first half of this account; we did not find in it a reported contrast in quiescence after long versus short outbursts (Cannizzo 1993). In the slow-state reading, duration acts as a proxy for a slowly varying disc or transfer state that the recent-gap level does not fully capture. The observed covariation of recurrence time with the long-to-short ratio points this way. The negative previous-duration coefficient is compatible with alternation, but a negative net lag does not exclude a dominant slow component coexisting with a stronger alternating one (Supplement S3.4). Neither reading is a mechanism test, and the word "memory" should not be applied to the result without a model that distinguishes them.

The weaker later segment is the most important open qualification. Its estimate is a quarter of the earlier segment's, and the exploratory class contrast is smaller there too. The record gives no basis for choosing between real change, a change in observing that alters extraction, and sampling variation in 180 forecasts. The design did not specify an era test, so we do not claim one.

Measurement limits remain. The duration and the following gap share the estimated end time, so end-time error couples them. The coefficient bound and matched routing were designed against the negative coupling seen in simulations. An exact construction shows the coupling can also produce positive information (Supplement S2.4). The uniform-end and delayed-issue sensitivities kept positive lower bounds, which probes this route without excluding it. Informative observing, missed outbursts and incomplete propagation of measurement error are further limits.

The study also carries lessons about method. The cycle convention that misled earlier work (CM92) has a forecasting analogue. Any predictor computed from data after the issue time contaminates the score, so we enforced observation-time features with poisoned-row tests. A component restriction in a frozen model can affect a secondary rule's score, as the exploratory class-rule adjustment illustrates without identifying a unique cause. A post-access format failure, however innocent, permanently changes what a test can be called.

Recording it openly is the only remedy.

4.2 Which conclusions depend on which assumptions

Table 2. Dependence of each main conclusion on its assumptions.

Calculated result	Rests on	Interpretation limits
The frozen duration model improves the score of extracted-onset forecasts	Fixed parameters and information set; cluster interval	One extraction and timing baseline; dependence assumptions and shared end-time error limit interpretation
Global recentering leaves a positive descriptive score contrast	Evaluation-fitted mean adjustment	Does not remove within-period drift or define a causal partition; intervals condition on the adjustment
The class indicator has greater exploratory in-sample likelihood	Fixed historical timing nuisance and a free level	Nonnested fitted alternatives; no exclusive attribution, within-class absence test or new forecast confirmation
The later-segment gain is inconclusive	Its interval includes zero	No era-difference or equivalence test; no evidence of disappearance
X4 passed a conditional synthetic recovery screen	Magnitude 10, C1/7 d and one qualified waveform/cadence pair	No real-event completeness guarantee; unseen waveform recovery was not tested

4.3 What would sharpen the answer

Four kinds of evidence would most improve these inferences.

- **A new frozen protocol for the closed data.** At the earlier issuance rate, the 2018-onward SS Cyg record would hold roughly 65 forecasts, a planning assumption and not a count or a power result. A new design study should quantify its contribution to a multi-star test. That test should use a class rule whose indicator coding and intercept freedom are defined together, and treat the anomalous 2021 SS Cyg event (Kimura et al. 2021) as a declared stress case.
- **Other long-record dwarf novae.** U Gem-type systems with long, well-sampled histories could test whether the result generalises. Their extraction must be qualified first.
- **An independent extraction for recent decades.** Photoelectric, CCD and space photometry would test sensitivity to the extraction and observing record in the later segment.
- **Stronger timing baselines.** Latent-state models of the quiet-interval sequence would test whether duration remains informative once a slowly varying state is modelled explicitly.

5. Conclusions

The completed outburst's duration improved frozen probabilistic forecasts of SS Cyg's next onset over a timing-only baseline, by 0.0165 nats per forecast across 441 held-out forecasts from 1960 to 2017. The preregistered criterion was met after a documented post-access parser repair. The gain was smaller and inconclusive in 1993–2017, and the predeclared long/short class rule was inconclusive. Exploratory fits favour a class reading, while CM92 had already reported the association, the alternation and the split. The result concerns extracted onsets for one star under one extraction and baseline. Whether it reflects cycle-to-cycle disc depletion or a slowly varying state, and whether it holds for other dwarf novae, remain open.

Methods in brief

- **Definitions, readings and units:** Supplement S1.

- **Extraction, coverage, issuance, outcome and censoring rules, models, fitting, scoring and interval construction:** S2.
- **Sensitivities, calibration, development comparison, exploratory calculations and the slow-driver counterexample:** S3.
- **Corrections made during review:** S4.
- **Reproduction commands and the evidence map:** S5.

The claims ledger (`manuscript/claims.csv`) gives every quoted number with its evidence file, its verifying commit, a reproduction command and its limits.

Roles of AI systems and humans

This work was conducted by AI systems with a human scientific lead, and the record is public.

- **Barry W. Brook** (University of Tasmania) set the scientific direction and the data boundaries. He chose to publish the protocol on Zenodo before opening the evaluation data, and keeps the 2018-onward record closed. He reviews the final products and decides on release and submission.
- **GPT-6 Pro** (OpenAI, through ChatGPT) produced the seed ideation and the first framework.
- **Claude Opus 5.5** (Anthropic, through Claude Code) did the following:
  - wrote the novelty and physical-framing audit;
  - developed and diagnosed the extraction methods, and wrote the X4 extractor and the first observation-time feature code;
  - proposed or reviewed each protocol;
  - reviewed the sealed result and ran the post-result exploratory analyses (A10j);
  - re-read the primary sources;
  - drafted this article, the supplement, the brief and the agent guide.

  A separate Claude Sonnet subagent refreshed the novelty search.
- **GPT-6 Codex** (OpenAI, through Codex) did the following:
  - implemented the forecasting, scoring, simulation and sealed-evaluation code;
  - executed the sealed evaluation and the format repair;
  - wrote the independent numerical audits of the result and the exploratory claims (A10k), which narrowed several of Claude's interpretations;
  - built the figures and the document toolchain;
  - reviewed every result-bearing passage in the first drafts, checked the narrowed novelty claim independently and inspected the rendered products (A10m).

Model names are the labels the systems reported on the project's correspondence board. Agreement between models is not treated as independent validation. Independence rests on separately implemented computational routes, which the audit records name. The human–AI correspondence is published with the repository.

Data and code

The repository https://github.com/BWBrook/dwarf-nova-memory holds the protocols, code, frozen parameters, aggregate evidence, figures, claims ledger and the full access record. Code is MIT licensed; text, figures and derived data are CC BY-SA 4.0. The evaluation protocol is archived at https://doi.org/10.5281/zenodo.24053338.

AAVSO observations are available from the AAVSO International Database (https://www.aavso.org). We do not redistribute raw observations, observer identities, event catalogues or per-forecast scores. The supplement (S5) gives the commands that reproduce every reported number from the retained run without new archive requests.

We acknowledge the worldwide AAVSO observer community and the International Database for the visual photometry and metadata used in this study. Observer credit and co-authorship will be reviewed before release against the AAVSO guidelines.

References

Bath, G. T., and J. van Paradijs. 1983. "Outburst Period–Energy Relations in Cataclysmic Novae." *Nature* 305 (5929): 33–36. https://doi.org/10.1038/305033a0.

Bürkner, Paul-Christian, Jonah Gabry, and Aki Vehtari. 2020. "Approximate Leave-Future-Out Cross-Validation for Bayesian Time Series Models." *Journal of Statistical Computation and Simulation* 90 (14): 2499–2523. https://doi.org/10.1080/00949655.2020.1783262.

Cannizzo, John K. 1993. "The Accretion Disk Limit Cycle Model: Toward an Understanding of the Long-Term Behavior of SS Cygni." *The Astrophysical Journal* 419: 318. https://doi.org/10.1086/173486.

Cannizzo, John K., and D. A. Goodings. 1988. "Chaos in SS Cygni?" *The Astrophysical Journal* 334: L31. https://doi.org/10.1086/185306.

Cannizzo, John K., and Janet A. Mattei. 1992. "On the Long-Term Behavior of SS Cygni." *The Astrophysical Journal* 401: 642. https://doi.org/10.1086/172092.

Gneiting, Tilmann, and Adrian E Raftery. 2007. "Strictly Proper Scoring Rules, Prediction, and Estimation." *Journal of the American Statistical Association* 102 (477): 359–78. https://doi.org/10.1198/016214506000001437.

Hameury, J.M. 2020. "A Review of the Disc Instability Model for Dwarf Novae, Soft X-Ray Transients and Related Objects." *Advances in Space Research* 66 (5): 1004–24. https://doi.org/10.1016/j.asr.2019.10.022.

Kimura, Mariko, Shinya Yamada, Nozomi Nakaniwa, Yoshihiro Makita, Hitoshi Negoro, Megumi Shidatsu, Taichi Kato, et al. 2021. "On the Nature of the Anomalous Event in 2021 in the Dwarf Nova SS Cygni and Its Multi-Wavelength Transition." *Publications of the Astronomical Society of Japan* 73 (5): 1262–79. https://doi.org/10.1093/pasj/psab073.

Lasota, Jean-Pierre. 2001. "The Disc Instability Model of Dwarf Novae and Low-Mass X-Ray Binary Transients." *New Astronomy Reviews* 45 (7): 449–508. https://doi.org/10.1016/s1387-6473(01)00112-9.

Price, A., A. A. Henden, G. Foster, V. Petriew, R. Huziak, R. James, M. D. Koppelman, et al. 2007. "SS Cygni Outburst Predictors and Long Term Quasi-Periodic Behavior." *Publications of the Astronomical Society of the Pacific* 119 (862): 1361–66. https://doi.org/10.1086/524413.

Fig. 1.




Figure 1: One forecast in the opened SS Cyg record, selected as the first eligible issuance by observation time rather than score. The upper panel shows observer-day display summaries, with detections separated from faint limits; arrows point toward fainter magnitudes. Prefix observations and the fitted crossing brackets define the completed-outburst duration available at issue. Display summaries are calculated separately on each side of issue. The lower panel shows the saved frozen timing and timing-plus-duration distributions, conditioned once on survival to issue. The later onset bracket supplies the scoring outcome. Reported observation times are known; historical upload and revision times are not. This example illustrates the retrospective information set, rather than validating the extraction or forecasting model. The completed comparison followed a post-access format repair.

Fig. 2.




Figure 2: Frozen forecast gain and its predeclared qualifications. The primary paired log-score gain is 0.0165 nats per issuance, with interval [0.0049, 0.0281], over 441 forecasts. Horizontal intervals in the lower panel use the predeclared cluster procedure; only the primary comparison is confirmatory. The later segment and frozen type/within-type rule remain inconclusive. The upper panel shows descriptive means for the fixed blocks, without block-specific confidence intervals; calendar labels are approximate. The result concerns incremental information beyond the specified timing baseline, following the recorded post-access format repair, and does not identify a physical mechanism.

Fig. 3.




Figure 3: Observed coverage of central predictive intervals for the constant, timing and timing-plus-duration forecasts. Each segment runs from definite to possible coverage given the extracted onset bracket or right censoring; filled and open endpoints distinguish these bounds. Dashed lines mark the nominal fractions. These are partial-identification bounds in the realised sample, conditional on the extraction and scoring population. They are not confidence intervals for population coverage or guarantees about physical onset times. The sample shows overcoverage; the primary paired gain does not imply sharply calibrated operational predictions.

334 **Fig. 4.**




Figure 4: Exploratory post-result comparisons, fitted using the evaluation outcomes. With a free level and historical timing nuisance parameters held fixed, the class-only fit has likelihood-ratio statistic 14.7, compared with 11.7 for continuous log duration; these equal-dimensional models are nonnested. Adding a linear within-class term to class gives 0.17. This is neither exclusive attribution to class nor an absence or equivalence test for within-class effects. The right panel gives conditional long/short ratios of the untruncated working-clock scale, rather than observed-gap or remaining-wait ratios. Era points are descriptive refits, not an era-difference test. No panel supplies a further held-out confirmation.