

IDENTIFYING PERINATAL HEALTH RISK USING MECHINE LEANING TECHNIQUES

Introduction

The perinatal period represents a vital period for maternal and infant health. Progress in perinatal health has been made in the past decades worldwide, represented by the decreased maternal and neonatal mortality, low birthweight (LBW), and preterm birth (PTB) rates. Recent studies have shown that artificial intelligence (AI) has the potential to accelerate this progress. This review aims to present an overview of perinatal health indicators, summarize evidence on the current state of evidence for AI application in perinatal health, and discuss future directions for this field.

Perinatal health indicators

Maternal mortality and morbidity; fetal, neonatal, and infant mortality; LBW (defined as birthweight <2500 g);¹ and PTB (defined as gestational age at birth <37 weeks)¹ are common indicators of perinatal health that are of public health concern and have significant implications for both the mother and child.

Some of the perinatal health indicators are included in the Sustainable Development Goals (SDGs) of the World Health Organization. SDG 3 that relates to good health and well-being includes goals to reduce maternal and neonatal mortality. One of the targets of SDG 3 is to reduce maternal mortality ratio to less than 70 per 100,000 births by 2030, with no country having a maternal mortality ratio of more than twice the global average. Another target is to reduce neonatal mortality in all countries to at least 12 per 1000 live births. Besides

this, a target of SDG 3 is to increase health financing and training of the health workforce, which relates to the use of current methods that are relatively cheap but efficient to improve perinatal health and well-being. This indicates that adequate antenatal care, skilled care at birth, and postnatal maternal and infant care are important for perinatal health and well-being.² This has been illustrated by a study that found a midwife-led continuity of care to reduce PTBs by nearly 24%.³ The American College of Obstetrics and Gynecology recommends provision of postpartum care as a continuum rather than a single encounter;⁴ therefore, one needs to focus on the woman not only during pregnancy but throughout the life course⁵ and incorporate improved health care system as a component of maternal and child care.

Maternal indicators

There is considerable difference in maternal mortality and morbidity rates between high-income countries (HICs) and low-income countries (LICs). In 2017, the maternal mortality ratio was estimated to be 11 per 100,000 live births in HIC, whereas in LIC it was 462 per 100,000 live births with the sub-Saharan Africa having the highest estimate at 534 per 100,000 live births.⁶ Notably, about 90% maternal deaths occur in LICs and lower-middle-income countries.⁷ Most maternal deaths occur due to obstetric hemorrhage, infections, hypertensive disorders during pregnancy, preexisting chronic health conditions, complications from delivery, and unsafe abortion.

Although maternal mortality has reduced in most HICs mainly due to skilled attendant care, severe maternal morbidity (SMM) is still a concern in most countries. In LIC, in addition to obstetric causes such as hemorrhage, hypertensive disorders during pregnancy, and sepsis due to obstetric infections, non-obstetric cause such as sepsis is a major contributor of maternal mortality.¹⁰ There are long-term implications of SMM—for example, preeclampsia is associated with an increased risk for cardiovascular disease for the mother.¹¹ For the

infants, SMM is associated with increased risk for fetal death, PTB, and LBW.

Child health indicators

Like maternal mortality, neonatal and infant mortality rates have reduced worldwide, with significant differences between HIC and LIC. In HIC, the neonatal mortality rate¹ is 3.0 per 1000 live births, and the infant mortality rate¹ is 4.0 per 1000 live births. In contrast, in LIC, the corresponding rates are 27 per 1000 live births and 48 per 1000 live births, respectively.¹⁴ Neonatal mortality constitutes one-third of deaths under 5 years of age, and like maternal morbidity it is related to lack of quality or skilled care at birth and treatment postpartum.³ The leading causes of neonatal/infant mortality are PTB; LBW; congenital anomalies; pregnancy complications such as infections, diabetes, or hypertensive disorders during pregnancy; intrapartum-related complications;³ and sudden and unexpected infant deaths.

Other indicators of perinatal health

Other indicators of perinatal health include congenital anomalies, which are related to perinatal health indicators such as PTB and neonatal and infant mortality.²⁶ Globally, congenital anomalies including chromosomal (e.g. aneuploidy), single gene defects, and non-chromosomal anomalies (e.g. congenital heart defects) are the second leading cause of neonatal death after birth asphyxia and birth trauma,²⁷ whereas in the United States it is the leading cause of infant mortality.²⁸ Besides the long-term physical, cognitive, and social impact on the child²⁹ and parental quality of life,³⁰ it is associated with substantial economic burden; in the United States, the cost of birth defect-associated hospitalizations was estimated to be US\$22.9 billion in 2013.³¹ Besides this, it is a significant contributor to PTB (more than 2.5 times higher risk) and LBW (more than 3.5 times higher risk).³² Common risk factors for congenital anomalies include chromosomal (most common being aneuploidy) or genetic, smoking, alcohol, substance abuse, diabetes mellitus, multiple gestation,³³ advanced maternal age, environmental teratogens, infections such as Zika and rubella,²⁷ and increased body mass index.

Future directions and role of AI in perinatal health

AI for perinatal health is in its nascent stages. If developed methodically and in collaboration with perinatal health specialists, AI research can benefit clinical aspects of perinatal health and even

research into applied AI at large. With more efficient AI tools serving perinatal health care researchers and clinicians that better predict conditions more precisely and sooner than current methods, AI can have stronger clinical impact. However, even more important than the need to develop robust AI is the need to develop explainable AI or AI that is transparent and able to be understood by humans.⁴⁸ In matters of health, development and validation of AI models should bring with it trust about how the model is actually processing and functioning.⁴⁹ Much more subtly and crucially, this AI should be understandable by clinicians since clinicians are tasked with diagnosis and treatment. For these reasons, a vision of AI for health and perinatal health care especially hinges on efficiency and “explainability.”

Thus, AI for perinatal health should have three goals for future research. First, AI research should aim to amass population-representative, routine clinical data for health research at large and specifically perinatal research. AI is a powerful tool that can be carefully built to include multimodal data but often requires large sample size to demonstrate strong performance. Thus, routinely collected, administrative primary and secondary care electronic health record (EHR) data that are representative of the population and collected systematically offer a solid foundation to begin AI research.⁵⁰ However, it is worth noting the inherent limitations of EHR data such as imprecise measurement, informed presence bias,⁵¹ and missing data.

Second, current state-of-the-art AI for prediction and classification in health care research should be modified and applied to the field of perinatal health. Research in AI for health has been growing in popularity. There has been much work in prediction of events and conditions in neurology, cardiology, and other medical focus areas.⁵² As an exemplar, in neurology, multimodal data—high-dimensional brain imaging and genetics—can be used in a deep-

learning model for improved prediction of epilepsy.⁵³ Another example is its use in anesthesiology where it can be used to improve the depth of anesthesia monitoring, pain management, and prediction of operative, postoperative, and critical care-related events.⁵⁴ Furthermore, in cardiovascular medicine, it can be used in the management of heart failure, including identification of patients at risk, development of risk assessment tools, and the use of multimodal EHR data for clinical decision support in these patients.⁵⁵ Another use in cardiology is in electrocardiography AI research where convolutional neural networks can be used to identify subclinical disease or prognostic factors of disease.⁵⁵ Prediction of PTB and LBW would be an important research contribution in perinatal health. A 2013 study found only an estimated 5% relative reduction in PTBs for years 2010 to 2015 resulting in US\$3 billion savings in total economic cost if evidence-based interventions were implemented in countries with very high human development index.⁵⁶ Also, two-thirds of the variation in total risk for PTB is still unaccounted for despite decades of research. This highlights the potential for research into factors that can lead to greater reduction in PTB and economic burden too. In addition, multi-task learning is an exciting area in ML research that involves simultaneous learning of multiple tasks by a shared model.⁵⁷ Pregnancy is a unique and finite period that involves shared maternal-fetal exposures resulting in pregnancy and infant outcomes. In this context, multi-task learning can be applied to study exposures common or unique to both the mother and fetus, interactions between them, and fetal outcomes such as live birth, stillbirth, congenital anomalies, LBW, and maternal outcomes such as obstetric hemorrhage, intrapartum infections using a single model.

Furthermore, real-time electronic health recording and predictive modeling have found initial success in the field of pregnancy health⁴⁰ and can especially be applied to health care systems in LIC. In HIC, 81% of the population is estimated to be living in urban areas, whereas in LIC this proportion is 33%.¹⁴ Women in rural areas,

irrespective of development status, have poor access to medical care and resources.⁵⁸ In addition, in HIC, the number of beds per 1000 people is 5.3, whereas in LIC it is 0.8 and in more than 55% of the countries the rate of nursing and midwifery is 10 per 10,000 population.⁷ In these settings, mHealth can help pregnant women with conditions such as preeclampsia to identify signs and symptoms and seek timely help. In low-resource settings where health care professionals mostly rely on last menstrual period to estimate gestational age, AI-based tools can enable a cost-effective and more accurate estimation of gestational age. AI can facilitate screening process for SMM in HICs as well as LICs for faster review based on a list of criteria.

Finally, AI for perinatal health requires methods for explaining the decision-making processes of its models. Recently, with an explosion of AI methods such as machine, deep, and reinforcement learning-based models for prediction tasks, the need to ask the question: “What is AI actually learning?” remains ever important. Answering this question requires the development of tools for responsibly gauging and vetting AI models. In the past decade, some tools to explain the importance of input variables in image and static data such as the Shapley Additive Explanations (SHAP)⁵⁹ and even explainers for longitudinal data⁶⁰ have allowed subject specialists to query what the model is attributing as signal. These methods have even been used in the field of perinatal health to understand input features strongly associated to the condition of autism spectrum disorder.⁶¹ When these tools illuminate the importance of a particular feature, specialists can sort these explanations into four categories as shown in [Figure 1](#). While features in squares 1 and 4 demonstrate agreement and little controversy between the subject area specialists and AI explainers, features sorted into the remaining two squares suggest further analyses to understand the actual importance of those features. AI understanding and clinical understanding might not always be in line with one another, and many such explanations may reside in squares

2 and 3; the resolution of these disagreements and progress in AI-driven perinatal health care require clinicians and AI specialists working side by side.

		Clinicians	
		Important	Not important
AI Explainer	Important	1	2
	Not important	3	4

Figure 1.

Explanations found important/non-important by clinicians (horizontal) and artificial intelligence (AI) explainers (vertical).

With these three goals, collaboration between perinatal health specialists and AI experts can yield robust AI tools along with trust in the tools. Both are necessary for progress and more importantly, clinical acceptance in areas such as perinatal health care.

Conclusion

In this review, we have summarized and highlighted the potential of AI-based algorithms and methods for the development of novel prediction models, better diagnosis, early identification, and monitoring of women during pregnancy, labor, and postpartum to advance research, clinical practice, and policies, and ensure optimal perinatal health.

Although this is a comprehensive and broad review of perinatal health indicators using AI where we have proposed a threefold approach to the use of AI in perinatal health research, it is not a systematic review that is a structured synthesis of evidence based on a priori protocol. Therefore, it is subject to biases such as publication bias and reviewer bias regarding choice of studies included and biases due to non-assessment of quality of studies included.

While our three goals for adaptation of AI for perinatal health care pave the way for novel findings for the field, we note that AI without societal acceptance and integration is insufficient for impacting clinical care. While robust performing AI and explainable AI can provide explanations for these black-box tools, integrating AI tools into the clinical setting and building community trust require engagement from every faction of society: from health care professionals to governmental workers and policy makers. Therefore, a multidisciplinary approach involving health care professionals, epidemiologists, statisticians, AI scientists, community health

workers, organizations and institutions (local, national, and global), engineers, local governments, policy makers, and most importantly the mother herself is required to improve health of mothers and children in developed and developing nations alike.