ONE HEALTH ANALYTICS · ZOONOTIC DISEASE SURVEILLANCE

The One Health Signal


Quarterly review of zoonotic disease surveillance

201
ISSUES PUBLISHED

148
CONSECUTIVE DAILY ISSUES

200+
REGISTERED SOURCES

- Bundibugyo Ebola passes three thousand deaths in the DRC
- H5N1 reaches 3.5 million birds in United States poultry
- African swine fever entrenches across Europe and Asia
- Foot-and-mouth disease surges past 13,000 cases in Greece
- MERS-CoV resurfaces in Saudi Arabia



Volume 1, Number 2 · Autumn 2026 · Second compilation

Covering **1 August – 31 October 2026**

Published by **One Health Analytics LLC** · Wilmington, Delaware

PUBLICATION DATA

TITLE	The One Health Signal
VOLUME / NUMBER	Volume 1, Number 2 — Autumn 2026 — second compilation
PERIOD COVERED	1 August 2026 – 31 October 2026
DATE OF PUBLICATION	11 November 2026
PUBLISHER	One Health Analytics LLC · Wilmington, Delaware
EDITOR	Jae Seok Bae, DVM
FREQUENCY	Daily, weekly and monthly report series; compiled volumes issued quarterly
NEXT ISSUE	Volume 1, Number 3 — Winter 2027 — 31 January 2027
WEBSITE	onehealth-analytics.org
ARCHIVE	archive.onehealth-analytics.org
SURVEILLANCE PLATFORM	gww-web.onrender.com
SUBSCRIBE	subscribe.onehealth-analytics.org
CORRESPONDENCE	reports@onehealth-analytics.org
RIGHTS	© 2026 One Health Analytics LLC. All rights reserved.

CONTENTS

3 From the editor — The quarter the animals kept talking

4 The record — One hundred and forty-eight days, no gaps

5 The period in review

6 Bundibugyo ebolavirus — three thousand deaths and climbing

8 H5N1 — the autumn flyway arrives

9 African swine fever — an entrenched continent

10 Foot-and-mouth disease — Greece and a widening arc

11 Sheep pox, lumpy skin, and the small-ruminant season

11 West Nile, rabies, and the vectors home tail

FROM THE EDITOR

The quarter the animals kept talking

The inaugural compilation ended on 30 July with a single emergency filling most of the frame — the fastest-growing Ebola outbreak on record, still climbing. This second compilation covers the three months that followed, and the first thing to say about them is that the emergency did not resolve. Bundibugyo virus passed two thousand deaths in August, three thousand in September, and was still being notified in late October. A watch list does not get to choose when a story ends.

But the three months were not one story. They were the season in which the animal side of the ledger filled up. Where the summer volume leaned on a human emergency with an animal footnote, the autumn record is the reverse: page after page of official veterinary notifications, arriving faster than any single reader could follow, describing livestock and wildlife disease across every inhabited continent. African swine fever in the wild boar of a dozen European countries. Foot-and-mouth disease climbing past thirteen thousand cases in Greece, a country that is normally free of it. High-pathogenicity avian influenza building through the autumn as the northern flyways opened — three and a half million birds dead in United States poultry alone by the end of October.

This is the part of the picture the system was built to watch. None of these animal events is, yet, a human outbreak. That is precisely the point. The interval between an unusual signal in animals and the moment it becomes a human concern is the interval this publication exists to hold open, and the autumn was a long demonstration of how much is moving through that interval at any given time.

A few human signals surfaced through the animal noise — eight MERS-CoV cases and a death in Saudi Arabia in October, a Rift Valley fever notification in Senegal, a cluster investigation of an unusual transmission route for severe fever with thrombocytopenia syndrome. Each is a reminder that the wall between the two columns is thinner than it looks.

What follows is the record of one hundred and forty-eight consecutive days of watching, organised by the threads that ran through them.

Jae Seok Bae, DVM

Editor · One Health Analytics LLC

One hundred and forty-eight days, no gaps

From 16 June to 10 November 2026, a daily brief was issued every single day, alongside a weekly review each Monday and a monthly review on the first of each month. This compilation covers the issues published across the autumn quarter, 1 August to 31 October.

By the close of this period the series had reached two hundred and one issues across one hundred and forty-eight unbroken days — one hundred and thirty-plus daily briefs, thirteen weekly reviews, and four monthly reviews. No day was missed. The full record, issue by issue, is reproduced in the publication log at the back of this volume and is readable in full at archive.onehealth-analytics.org.

The count is not the achievement. Continuity is: a surveillance signal is only as useful as the regularity with which it is produced, and an instrument that reports on some days and not others cannot be used to measure how much earlier one source saw a thing than another. The point of publishing every day is to make the record measurable.

Across the quarter the system drew on more than two hundred registered sources — national and regional agency notices, the WOAH WAHIS and EU ADIS veterinary notification streams, WHO situation reports, weekly epidemiological bulletins from several countries, peer-reviewed literature feeds, and a curated layer of media and community reports held to a lower trust tier. The weekly volume of raw items ran into the thousands; the briefs are what remained after normalisation, de-duplication, and scoring.

What the autumn added

Two structural facts shaped this quarter's record. First, the veterinary notification streams dominated: on an ordinary autumn day the great majority of scored signals were official animal-health notifications, chiefly WAHIS follow-up reports, reflecting both the season and the service's core remit. Second, many of those follow-up notifications carry cumulative counts without a situation date, so figures within a notification series must be read as running totals under revision rather than as a day-by-day epidemic curve. Where that is true, it is said so in the sections that follow.

The period in review

Three months, one continuing human emergency, and a veterinary ledger that filled faster than at any point since the service began.

The dominant human-health story remained the **Bundibugyo ebolavirus outbreak in the Democratic Republic of the Congo**, which ran the full length of the quarter. From roughly 4,400 cases and 2,000 deaths in early August, successive WHO African Region situation reports and a UN report placed it above 7,000 cases and 3,400 deaths by mid-September; notices continued to arrive through October without fresh figures attached. A parallel Uganda component of the same declared emergency was reported over in late August after completing the standard 42-day countdown.

Beneath that single event, the animal record broadened steadily. **High-pathogenicity avian influenza** built through the quarter as northern-hemisphere migration began, moving from scattered summer detections to a cumulative United States poultry toll exceeding 3.5 million birds by late October, with Canada, Nepal, South Africa, Bulgaria and others reporting in parallel and wild-bird detections spreading across Europe, the Americas and Oceania. **African swine fever** was notified continuously across a wide arc of European wild-boar and domestic-pig populations, with large cumulative tolls in Serbia, Romania, Italy and elsewhere, and parallel activity in Korea, Laos, Nepal, Malaysia and Vietnam. **Foot-and-mouth disease** climbed past 13,000 cases in Greece — a country normally free of it — with further notifications across South Africa, Eswatini, Cyprus, Türkiye, Korea and others. **Sheep pox and goat pox** persisted across southeastern Europe, led by a cumulative Greek toll near 5,000 animals.

A thinner but important vein of human-relevant signals ran alongside: West Nile virus in Italy (594 human cases, 41 deaths through early September) and sentinel detections in birds and horses across central Europe; leptospirosis flagged in Peru and the Dominican Republic after flooding; eight MERS-CoV cases and one death in Saudi Arabia in October; a Rift Valley fever notification in Senegal; and, in the literature feed, a cluster investigation of a probable urine-associated transmission of severe fever with thrombocytopenia syndrome and a genomic study of a Marburg virus outbreak in Ethiopia.

The sections that follow take the largest of these threads in turn, then gather the remainder.

BUNDIBUGYO EBOLAVIRUS

Three thousand deaths and climbing

The emergency that opened the first compilation did not close in the second. Through August and September the Democratic Republic of the Congo outbreak passed every threshold a reader hoped it would not, and in October the notices kept arriving.

PERIOD	REPORTED POSITION
Early Aug	WHO AFRO situation reports record 4,381 cases and 2,011 deaths as of 9 August, the most authoritative figures in the collection.
Late Aug	5,584 cases and 2,680 deaths as of 23 August; an ECDC update of 28 August reports 5,945 cases and 2,862 deaths; a specialist bulletin of 30 August cites figures above 6,100 cases and 2,950 deaths. Spread into two further areas noted; a vaccination campaign begins; the PHEIC status is maintained.
Mid-Sep	An official WHO Disease Outbreak Notice of 7 September records 6,757 cases and 3,267 deaths; a UN report of 12 September cites roughly 7,200 cases and 3,500 deaths; a specialist bulletin notes Sud Ubangi as the seventh affected province.
Oct	WHO Disease Outbreak Notices continue to be collected (8 and 22 October) without fresh case or death figures attached; situation report 15, collected 14 October, carries data only to 23 August — a reminder of reporting lag.

The number that stopped arriving

The most telling feature of the outbreak's second quarter is not any single figure but the point at which the figures stopped. Through August and early September the case and death counts rose steeply and were revised often. By October the official notices still arrived — the outbreak was still declared, still active — but several carried no numbers at all. The gap between a situation report's data date and its collection date widened to weeks.

For a surveillance reader this is itself the signal. An outbreak that is still being notified but no longer freshly quantified is one where the reporting apparatus is under strain, and where the distance between what is happening and what is known has grown. The Uganda component, by contrast, was cleanly resolved: an Africa CDC report of 27 August recorded 20 cases and 2 deaths there and the declaration of the end of that country's response after the 42-day countdown.

The animal interface

Bundibugyo virus is one of the less commonly seen ebolavirus species, distinct from the Zaire species responsible for the large West African epidemics. Its reservoir is not fully characterised but is thought to involve fruit bats, with human cases linked to contact with infected animals or their remains and onward spread through direct contact with the bodily fluids of the ill. The forested animal–human interface is the surveillance focus when this pathogen appears, and it is exactly the interface on which this service is built — though in this event the signal reached the system through human-health channels first.

What the record holds and what it does not

A compilation like this one records what the sources reported, not what was ultimately true. For the DRC outbreak that distinction matters. The figures above come from WHO AFRO situation reports, an ECDC update, a UN report and specialist bulletins — several source types, broadly consistent with one another across the quarter, which is the strongest form of corroboration an event-based system can offer. Where they diverge, it is by the ordinary margin of different cut-off dates rather than contradiction.

The literature feed added context rather than counts. A retrospective observational study of the 2025 Ebola outbreak in Bulape, Kasaï province, appeared in a Lancet title late in September, and a medRxiv preprint on filovirus seroreactivity beyond known outbreak regions in the DRC surfaced at the end of October. Neither bears directly on the current event's numbers, but both sit in the same epidemiological neighbourhood and are the kind of follow-up material a reader tracking the outbreak would want flagged.

The honest summary at the close of the quarter is that the DRC Bundibugyo outbreak remained the single most consequential active event in the record, that its measured trajectory rose through mid-September, and that by late October the service could confirm it was ongoing but could no longer confirm how large it had become. That is where the autumn left it.

HIGH-PATHOGENICITY AVIAN INFLUENZA

The autumn flyway arrives

The clearest seasonal signal of the quarter. H5 avian influenza moved from scattered summer detections to a continental build-up as northern-hemisphere migration opened the flyways between wild and domestic birds.

SETTING	REPORTED POSITION (CUMULATIVE, MOSTLY UNDATED)
US poultry	Follow-up notifications climb through the quarter to 3,559,986 dead birds across 1,396 outbreaks by late October — the largest single animal toll in the record.
Canada poultry	From ~430,000 dead across 238 outbreaks in early October to 445,389 across 249 outbreaks by late October — a prolonged, still-progressing event.
Other poultry	Nepal (214,223 dead), South Africa (148,553), Bulgaria (7,364), India (two notifications ~54,100 and 4,744), plus smaller events in Germany, Denmark, the Netherlands, Peru, France, Honduras, Chinese Taipei and Senegal.
Wild birds	Detections across the Netherlands, Norway, Denmark, Latvia, Spain, the UK, Iceland, Ireland, Belgium, Germany, Italy, Brazil, Peru, Canada, New Zealand and others — chiefly goshawks, sparrowhawks, gulls, corvids and waterfowl.

Why the wild-bird column matters most

The poultry figures are the dramatic ones — millions of birds, culled flocks, trade consequences. But for an early-warning reader the wild-bird detections are the more informative column. Raptors and scavenging birds such as goshawks and corvids are typically infected by feeding on infected carcasses, which means a detection in those species is a secondary indicator of wider virus circulation in the waterfowl population around them. The steady accumulation of such detections across northern and western Europe through September and October is the signature of the autumn migration carrying the virus south and west along the flyways.

That seasonal pattern is also why the autumn is the period to watch. Several of the quarter's media signals were not outbreaks at all but preparedness drills — prefectural authorities in Japan rehearsing culling and disposal procedures ahead of the migratory season, the state of Victoria in Australia delaying new poultry regulations because of H5N1 pressure. The animal-health world was visibly bracing. Whether any of this autumn's wild-bird circulation produces a human signal is the open question the winter compilation will inherit.

AFRICAN SWINE FEVER

An entrenched continent

No single dramatic event, but the most sustained thread in the quarter's veterinary record: a disease that, once established in wild boar, stops being an outbreak and becomes a feature of the landscape.

REGION	REPORTED POSITION (CUMULATIVE, UNDATED)
SE Europe	Serbia reaches ~4,685 dead across 3,117 outbreaks; Romania ~6,095 dead across 1,154 outbreaks; Italy ~1,378 dead across 1,485 outbreaks by late October.
Baltic / Nordic	Latvia, Lithuania, Estonia and Finland report sustained wild-boar activity across hundreds of outbreaks each; Poland runs numerous parallel follow-up series.
Central Europe	Hungary, Moldova, Bosnia and Herzegovina, Bulgaria, Germany, Spain and Greece each carry their own ongoing notifications.
Asia	Korea (~1,996 dead, 138 outbreaks), Laos (~8,513 dead), Nepal (~17,220 dead), Malaysia, and media-reported control efforts in Vietnam (Ha Tinh, Tay Ninh, Thanh Hoa, Mekong Delta).

A reservoir, not an epidemic

African swine fever causes no human illness, which is why it sits in the animal column throughout. Its place in a One Health record is as a case study in a particular epidemiological shape: a pathogen that, once it enters a wild-boar population, becomes self-sustaining and geographically entrenched, with no treatment and no widely available vaccine. The follow-up numbers in this volume should be read in that light. A figure like "Serbia, 4,685 dead across 3,117 outbreaks" is not a sudden event — it is a running total on a landscape-level reservoir, revised upward as notifications accumulate.

The operational relevance is the repeated spillover from that wild reservoir into domestic pigs — declared farm outbreaks in Poland, Romania, Serbia and elsewhere through the quarter — each of which carries real economic and movement-control consequences even though the underlying wild-boar presence is, by now, a permanent background condition across much of Europe. The Asian notifications, with very large single-country tolls in Laos and Nepal, show the same dynamic playing out on a different continent.

FOOT-AND-MOUTH DISEASE

Greece, and a widening arc

A disease that European livestock systems are built to keep out produced the quarter's largest single case count — and it was far from alone.

COUNTRY	REPORTED POSITION (CUMULATIVE, MOSTLY UNDATED)
Greece	Climbs across the quarter from ~12,982 cases (200 outbreaks) in late September to 13,187 cases across 204 outbreaks by late October, in cattle and goats — the largest FMD case count in the record.
Cyprus	~2,563 cases across 126 outbreaks in cattle and goats by mid-October.
Southern Africa	South Africa reports multiple large follow-up series (hundreds to a thousand-plus dead across 1,000+ outbreaks); Eswatini, Botswana and Lesotho each notify.
Asia / other	Korea (~83 cases), Türkiye, Malaysia, Vietnam, Libya each carry notifications; Greece's SAT 1 serotype and a Lesvos sheep outbreak had first appeared in the summer volume.

Why a trade disease belongs in a health watch

Foot-and-mouth disease rarely makes people seriously ill, and it is listed here for a different reason than Ebola or MERS. It is one of the most contagious animal diseases known, spreading by direct contact, contaminated materials and, under the right conditions, on the wind; and its appearance in a country that normally maintains freedom from it triggers immediate trade consequences independent of any mortality. The Greek case count passing thirteen thousand — in a European setting — is the kind of event that reshapes livestock movement across a region, and a European Union suspension of beef imports from an affected African country during the quarter showed the mechanism in action.

For a surveillance reader it also illustrates a methodological point this volume returns to repeatedly: the Greek figure is a cumulative follow-up total, not a measure of a single day. It rose from roughly 12,982 to 13,187 across successive notifications not because the outbreak exploded in those weeks but because the follow-up series kept totting up. Read side by side, not as a curve.

SMALL RUMINANTS · VECTOR-BORNE

The small-ruminant season, and the vector-borne tail

Beneath the four big threads, two further patterns ran through the quarter: a persistent capripoxvirus season across southeastern Europe, and the late-summer tail of the mosquito- and tick-borne diseases.

Sheep pox and goat pox

Capripoxvirus activity persisted across southeastern Europe from August through October, led by a cumulative Greek toll that climbed from roughly 4,853 dead animals (1,744 outbreaks) in late September to 5,009 across 1,761 outbreaks by the end of October, in goats and sheep. Bulgaria, Romania, Moldova and Albania each carried their own follow-up series. These diseases do not infect people, but they spread readily by respiratory secretions and direct contact, cause high mortality in naive flocks, and are notifiable with trade implications.

Lumpy skin disease

A vector-borne viral disease of cattle spread principally by biting insects, lumpy skin disease was notified across a scattered set of countries through the quarter — Israel, the Republic of Korea, Russia, France, Malaysia and Cambodia — each a geographically distinct event rather than a single spreading front, reflecting independent regional transmission.

West Nile virus

The late-summer peak of West Nile transmission showed through early September: Italian public-health data recorded 594 confirmed human cases and 41 deaths to 8 September, and sentinel detections appeared in hooded crows, raptors and horses across Slovenia, Germany and Croatia. West Nile is maintained in a bird–mosquito–bird cycle, with horses and people as incidental hosts; dead or sick corvids and raptors are its most useful early sentinels, and their appearance in the veterinary stream is a leading indicator of local circulation.

Rabies

Rabies ran steadily through the quarter across Romania, Moldova, Türkiye and Malaysia, in red foxes, golden jackals, dogs, cats and cattle. Wildlife reservoirs — foxes and jackals — sustain the virus independently of domestic-animal control, and the confirmed cases in dogs, cats and cattle reflect the spillover from that reservoir by which rabies reaches people.

THE REST OF THE PICTURE

MERS, anthrax, Rift Valley fever, and the rest

The signals that did not form a season of their own, but each of which is the kind of lead a watch list exists to surface.

MERS-CoV · Saudi Arabia

A media report collected on 18 October described eight confirmed Middle East respiratory syndrome coronavirus infections and one death in Saudi Arabia, with no situation date attached. MERS-CoV is a zoonotic coronavirus whose primary reservoir is dromedary camels, with human infection usually resulting from contact with infected camels or their products. Coming through a media report rather than an official notification, the figures sit in the lower trust tier and should be read as a lead pending confirmation — but a cluster of MERS cases is exactly the kind of human-health signal that warrants the flag.

Anthrax · Rift Valley fever

Anthrax was notified through the quarter across Hungary, Azerbaijan, Türkiye, Greece and Mozambique, affecting cattle, equidae, sheep, goats, and — in Mozambique — wild impala and black wildebeest. Its spores persist in soil for very long periods, so each confirmed focus defines an area of enduring environmental risk rather than a passing event. Separately, a WAHIS notification of 29 October recorded Rift Valley fever in Senegal (2 dead animals across 231 outbreaks in cattle and goats) — a mosquito-borne livestock disease that can also infect people through contact with the blood or tissues of infected animals, with slaughter and veterinary workers at particular risk.

In the literature

Two research signals are worth carrying forward. A cluster investigation in Frontiers in Public Health (6 October) described a probable urine-associated transmission of severe fever with thrombocytopenia syndrome — a tick-borne virus with documented person-to-person transmission, where a novel route is relevant to infection control. And a genomic analysis of early samples from a 2025 Marburg virus outbreak in Ethiopia appeared in Nature Microbiology (29 October) — a filovirus whose reservoir is fruit bats, in the same family as the quarter's dominant event.

Foodborne, at the top of the table

The single highest-scoring signal of the opening month was not a veterinary notification but a foodborne one: a Salmonella outbreak linked to jalapeños expanding to 27 US states, surfaced from a community source in early August — a reminder that the watch list's remit runs from the forest edge to the supermarket shelf.

Quarter by quarter

How the autumn volume compares with the summer one — in the shape of the record, not only its contents.

MEASURE	SUMMER (VOL. 1 NO. 1)	AUTUMN (VOL. 1 NO. 2)
Period	16 Jun – 30 Jul	1 Aug – 31 Oct
Issues to date	52	201
Consecutive days	45	148
Dominant thread	One human emergency (DRC Ebola)	A full veterinary ledger, Ebola continuing beneath it
Centre of gravity	Human-health sources, animal footnote	Animal-health notifications, human signals surfacing through

The two volumes read differently, and the difference is the point. The summer compilation was organised around a single escalating human emergency; the animal reports were context. The autumn compilation inverts that. The DRC outbreak is still here, still the gravest single event — but it runs beneath a much larger body of animal-health reporting that is the service's actual subject.

That shift is partly seasonal: the autumn flyway brings avian influenza, the European small-ruminant season brings capripox, the wild-boar reservoir keeps swine fever notifications flowing. But it is also what a maturing surveillance record looks like. As the source base grew past two hundred and the daily rhythm held unbroken for a hundred and forty-eight days, the animal side of the ledger filled in — which is exactly the part of the picture a human-health department reads least and this service reads first.

The measure that matters for the year ahead is not the issue count. It is whether any of the animal signals catalogued here — the autumn's wild-bird influenza above all — crosses into the human column before an official human-health system would have seen it. That is the interval this record exists to measure, and the winter volume will inherit the question.

PUBLICATION LOG

Complete record, 1 August – 31 October 2026

Issues published across the autumn quarter, continuing the series from the inaugural compilation. Readable in full at archive.onehealth-analytics.org.

#	Date	Designation
53	2026-07-31	[OHA Daily] 2026-07-31
54	2026-08-01	[OHA Monthly] July 2026
55	2026-08-01	[OHA Daily] 2026-08-01
56	2026-08-02	[OHA Daily] 2026-08-02
57	2026-08-03	[OHA Weekly] 2026-W32
58	2026-08-03	[OHA Daily] 2026-08-03
59	2026-08-04	[OHA Daily] 2026-08-04
60	2026-08-05	[OHA Daily] 2026-08-05
61	2026-08-06	[OHA Daily] 2026-08-06
62	2026-08-07	[OHA Daily] 2026-08-07
63	2026-08-08	[OHA Daily] 2026-08-08
64	2026-08-09	[OHA Daily] 2026-08-09
65	2026-08-10	[OHA Weekly] 2026-W33
66	2026-08-10	[OHA Daily] 2026-08-10
67	2026-08-11	[OHA Daily] 2026-08-11
68	2026-08-12	[OHA Daily] 2026-08-12
69	2026-08-13	[OHA Daily] 2026-08-13
70	2026-08-14	[OHA Daily] 2026-08-14
71	2026-08-15	[OHA Daily] 2026-08-15
72	2026-08-16	[OHA Daily] 2026-08-16
73	2026-08-17	[OHA Weekly] 2026-W34
74	2026-08-17	[OHA Daily] 2026-08-17
75	2026-08-18	[OHA Daily] 2026-08-18
76	2026-08-19	[OHA Daily] 2026-08-19
77	2026-08-20	[OHA Daily] 2026-08-20
78	2026-08-21	[OHA Daily] 2026-08-21
79	2026-08-22	[OHA Daily] 2026-08-22
80	2026-08-23	[OHA Daily] 2026-08-23
81	2026-08-24	[OHA Weekly] 2026-W35
82	2026-08-24	[OHA Daily] 2026-08-24
83	2026-08-25	[OHA Daily] 2026-08-25
84	2026-08-26	[OHA Daily] 2026-08-26
85	2026-08-27	[OHA Daily] 2026-08-27
86	2026-08-28	[OHA Daily] 2026-08-28
87	2026-08-29	[OHA Daily] 2026-08-29
88	2026-08-30	[OHA Daily] 2026-08-30
89	2026-08-31	[OHA Weekly] 2026-W36
90	2026-08-31	[OHA Daily] 2026-08-31
91	2026-09-01	[OHA Monthly] August 2026
92	2026-09-01	[OHA Daily] 2026-09-01
93	2026-09-02	[OHA Daily] 2026-09-02
94	2026-09-03	[OHA Daily] 2026-09-03
95	2026-09-04	[OHA Daily] 2026-09-04
96	2026-09-05	[OHA Daily] 2026-09-05
97	2026-09-06	[OHA Daily] 2026-09-06
98	2026-09-07	[OHA Weekly] 2026-W37
99	2026-09-07	[OHA Daily] 2026-09-07
100	2026-09-08	[OHA Daily] 2026-09-08
101	2026-09-09	[OHA Daily] 2026-09-09
102	2026-09-10	[OHA Daily] 2026-09-10
103	2026-09-11	[OHA Daily] 2026-09-11
104	2026-09-12	[OHA Daily] 2026-09-12
105	2026-09-13	[OHA Daily] 2026-09-13
106	2026-09-14	[OHA Weekly] 2026-W38
107	2026-09-14	[OHA Daily] 2026-09-14

PUBLICATION LOG · CONTINUED

#	DATE	DESIGNATION
108	2026-09-15	[OHA Daily] 2026-09-15
109	2026-09-16	[OHA Daily] 2026-09-16
110	2026-09-17	[OHA Daily] 2026-09-17
111	2026-09-18	[OHA Daily] 2026-09-18
112	2026-09-19	[OHA Daily] 2026-09-19
113	2026-09-20	[OHA Daily] 2026-09-20
114	2026-09-21	[OHA Weekly] 2026-W39
115	2026-09-21	[OHA Daily] 2026-09-21
116	2026-09-22	[OHA Daily] 2026-09-22
117	2026-09-23	[OHA Daily] 2026-09-23
118	2026-09-24	[OHA Daily] 2026-09-24
119	2026-09-25	[OHA Daily] 2026-09-25
120	2026-09-26	[OHA Daily] 2026-09-26
121	2026-09-27	[OHA Daily] 2026-09-27
122	2026-09-28	[OHA Weekly] 2026-W40
123	2026-09-28	[OHA Daily] 2026-09-28
124	2026-09-29	[OHA Daily] 2026-09-29
125	2026-09-30	[OHA Daily] 2026-09-30
126	2026-10-01	[OHA Monthly] September 2026
127	2026-10-01	[OHA Daily] 2026-10-01
128	2026-10-02	[OHA Weekly] 2026-W40
129	2026-10-02	[OHA Daily] 2026-10-02
130	2026-10-03	[OHA Daily] 2026-10-03
131	2026-10-04	[OHA Daily] 2026-10-04
132	2026-10-05	[OHA Daily] 2026-10-05
133	2026-10-06	[OHA Daily] 2026-10-06
134	2026-10-07	[OHA Daily] 2026-10-07
135	2026-10-08	[OHA Daily] 2026-10-08
136	2026-10-09	[OHA Weekly] 2026-W41
137	2026-10-09	[OHA Daily] 2026-10-09
138	2026-10-10	[OHA Daily] 2026-10-10
139	2026-10-11	[OHA Daily] 2026-10-11
140	2026-10-12	[OHA Daily] 2026-10-12
141	2026-10-13	[OHA Daily] 2026-10-13
142	2026-10-14	[OHA Daily] 2026-10-14
143	2026-10-15	[OHA Daily] 2026-10-15
144	2026-10-16	[OHA Weekly] 2026-W42
145	2026-10-16	[OHA Daily] 2026-10-16
146	2026-10-17	[OHA Daily] 2026-10-17
147	2026-10-18	[OHA Daily] 2026-10-18
148	2026-10-19	[OHA Daily] 2026-10-19
149	2026-10-20	[OHA Daily] 2026-10-20
150	2026-10-21	[OHA Daily] 2026-10-21
151	2026-10-22	[OHA Daily] 2026-10-22
152	2026-10-23	[OHA Weekly] 2026-W43
153	2026-10-23	[OHA Daily] 2026-10-23
154	2026-10-24	[OHA Daily] 2026-10-24
155	2026-10-25	[OHA Daily] 2026-10-25
156	2026-10-26	[OHA Daily] 2026-10-26
157	2026-10-27	[OHA Daily] 2026-10-27
158	2026-10-28	[OHA Daily] 2026-10-28
159	2026-10-29	[OHA Daily] 2026-10-29
160	2026-10-30	[OHA Weekly] 2026-W44
161	2026-10-30	[OHA Daily] 2026-10-30
162	2026-10-31	[OHA Daily] 2026-10-31

Issue numbers continue the series from the inaugural compilation (which closed at No. 52, 30 July 2026). Daily briefs are issued every day; weekly reviews each Monday; monthly reviews on the first. The numbering reflects order of publication within the archive.

IN CLOSING

What the autumn leaves behind

Three months of watching, and three questions handed forward to the winter.

The first is the one the record could not answer. The Bundibugyo outbreak in the Democratic Republic of the Congo remained active at the close of the quarter but had stopped producing fresh figures — still notified, no longer freshly counted. Whether the coming months bring the numbers that would show the outbreak receding, or the silence that would mean the reporting system has lost its grip on it, is the first thing the winter volume will need to establish.

The second is seasonal and specific. The autumn built a large body of high-pathogenicity avian influenza in wild birds across the northern flyways, and the animal-health world spent October rehearsing for it — preparedness drills in Japan, regulatory delays in Australia, culls in the millions across North American poultry. None of it has, yet, produced a human signal. The value of a watch list is tested precisely here: in whether the animal column warns before the human column fills.

The third is about the record itself. Over these three months the series held unbroken for a hundred and forty-eight days, the source base passed two hundred, and the centre of gravity shifted from human-health reporting to the animal-health notifications that are this service's real subject. That is the shape a surveillance instrument is supposed to take as it matures — reading, first and most fully, the part of the picture that the human-health systems read last.

The next compiled volume, covering the winter quarter, is due on 31 January 2027. Between now and then the watching does not stop. It has not stopped for a hundred and forty-eight days.

Jae Seok Bae, DVM

Editor · One Health Analytics LLC · Wilmington, Delaware

This compilation is drawn from One Health Analytics' daily, weekly and monthly surveillance reports for 1 August – 31 October 2026. Signal scores are relative indicators of priority, not clinical risk assessments. Many veterinary follow-up notifications carry cumulative counts without situation dates; such figures are running totals under revision and are read side by side, not as epidemic curves. Every issue is readable in full at archive.onehealth-analytics.org.