Certificate Bounds for Backtracking Frank–Wolfe under Function-Dependent Curvature

Yaroslav P. Lyakhov Stepan Suvorov

17 October 2026

Abstract

We study convex minimization over a compact convex set when the Hessian norm is bounded by $H_0 + H_1(f - f_Q^*)$, relative to the constrained minimum. For ordinary quadratic backtracking Frank–Wolfe, we prove a bound of $O(H_0 D^2/\varepsilon + (1 + H_1 D^2)\log(U_0/\varepsilon))$ updates to an observable certificate, together with an explicit initialization term. The method receives neither curvature parameters nor the optimal value. A phase argument controls estimates that lag behind falling local curvature, and all trial function evaluations cost at most $1 + 2N$ plus an explicit logarithmic overhead after N updates. Standard away steps inherit the convex bound. Under a geometric error bound they yield logarithmic certificate accuracy dependence, with a separate initial curvature contribution. A differential comparison gives the optimal pointwise Frank–Wolfe gap envelope, with one-dimensional attaining objectives. When $H_0 = 0$, the solution set is an exposed face on which the gradient is constant; backtracking away-step Frank–Wolfe then terminates exactly in finitely many updates without a separate geometric-growth assumption. A quadratic counterexample separates this endpoint from every $H_0 > 0$. Exact segment minimization and known-curvature short steps provide comparisons in the same linear-oracle model. Controlled experiments with analytically verified objectives and real Poisson data separate initialization, active-face geometry, and stopping cost. They show no uniform advantage of certificate-scaled initialization and expose regimes where the scalar certificate substantially lags primal accuracy. The results concern an existing backtracking method; they establish neither a new algorithm nor optimal oracle complexity.

1 Introduction

Frank–Wolfe (FW) methods solve linear subproblems in place of projections. Backtracking makes their quadratic step models usable without a prescribed smoothness constant, and away steps improve progress on polytopes [1, 2, 3]. A global curvature bound can nevertheless discard substantial information when curvature decreases with the objective value. We consider the affine loss-suboptimality Hessian condition studied in [4], specialized to a constrained convex problem. It is also the exponent-one case of the loss-dependent smoothness family in [27]. More general function-value bounds on Hessians already appear in self-bounding regularity frameworks [5, 6]; the condition itself is not a contribution of this paper.

Our question is quantitative: what does ordinary backtracking guarantee for an observable stopping rule under this condition, including its initial estimate and all trial evaluations? Theorem 2.3 answers it with separate H_0/ε and $H_1 \log(1/\varepsilon)$ contributions. The main device is a potential for estimates that remain too large as the certificate enters lower dyadic levels. It does not change the algorithm or restrict the LMO to a sublevel set. Theorem 3.1 extends the result by standard atom-removal counting. Under a geometric gap inequality, Theorem 3.3 gives logarithmic accuracy dependence with an explicit initial H_1 cost. Theorem 3.2 gives the optimal pointwise FW-gap envelope for this class, with attaining one-dimensional objectives. Proposition 4.1 characterizes the zero-base case $H_0 = 0$: its entire solution set is an exposed

face with a common gradient. Theorem 4.2 then upgrades approximate stopping to finite exact termination of backtracking AFW, with an explicit bound independent of requested accuracy. This uses known identification mechanisms [7, 8]; neither a new AFW algorithm nor general identification theory is asserted. Complete proofs, boundary examples and counterexamples are provided.

Related analyses. Function-dependent smoothness and two-regime gradient-method bounds precede this work [4, 9, 10]. Vyguzov and Stonyakin [11] analyze FW under gradient-norm dependent smoothness. On a compact convex domain, $f - f_Q^* \le D \|\nabla f\|$ converts our condition to one such Hessian bound, but a global bound on the largest gradient does not retain the separated dependence studied here. Pedregosa et al. [2, Theorems 1, 3, and 4] already give backtracking trial counts, an averaged-gradient Fenchel dual guarantee, and linear AFW convergence with accepted local curvature estimates. Our scalar lower-bound maximum differs from that Fenchel dual; their values cannot simply be identified. The result here specializes local curvature to a certificate threshold and explicitly accounts for lag. It does not claim priority for backtracking, dual certificates, linear AFW convergence, or the telescoping identity for rejected trials. Best-so-far tangent bounds and their computable stopping gaps are also classical [28, Method 1]. Optimistic FW and primal–dual short steps explicitly optimize computable gaps under global smoothness [29, Proposition 4.1]; they do not evaluate the inherited halving/doubling rule analyzed here.

Other related frameworks include generalized self-concordant FW, adaptive Bregman steps, and adaptive conditional gradient sliding [12, 13, 14]. Extended-curvature bounds and restarted error-bound schemes already provide two-regime and geometric rates [15, 16]; linear objective- and FW-gap convergence beyond global gradient smoothness is also known for log-homogeneous barriers [17]. Auto-conditioned FW supplies a practical comparator with different preprocessing and oracle accounting [18]. Gradient-difference initialization with backtracking [19] and line-search-free conditional gradients [20] provide further parameter-adaptive alternatives. ALFCG estimates smoothness without line search but retains global component or average smoothness assumptions and a stochastic stationarity criterion [30, Section 3]. Recent damped Newton–FW uses Hessian information and assumes positive definite Hessians [31, Section 1]; this does not cover our zero-base optimum, where the Hessian vanishes. Appendix E specifies the differences used in our comparison. A direct global-smoothness substitution is generally weaker, but that observation is not a claim that no equivalent consequence can be obtained from another adaptive analysis.

The empirical study addresses three issues relevant to interpreting these bounds: whether certificate initialization helps independently of a starting-point change; whether away steps resolve active-face obstructions; and whether failure to stop reflects remaining primal error or a conservative certificate. Oracle budgets are kept separate, and failed runs are retained.

2 Ordinary backtracking and a computable certificate

Let $Q \subset \mathbb{R}^d$ be nonempty, compact and convex, with diameter $D > 0$. Let f be convex and C^2 on a neighborhood of Q, $f_Q^* = \min_Q f$, and $\delta(x) = f(x) - f_Q^*$. Assume

$$\left\|\nabla^2 f(z)\right\|_2 \le H_0 + H_1 \delta(z), \qquad z \in Q, \quad H_0, H_1 \ge 0. \tag{1}$$

The reference value in this assumption is the constrained optimum. A bound relative to a smaller unconstrained infimum does not imply the same constrained bound without modifying H_0. If $D = 0$ the only feasible point solves the problem.

At a source x_k, obtain an exact linear minimization oracle (LMO), and form the FW gap

and accumulated certificate:

$$s_k \in \arg\min_{v\in Q} \langle \nabla f(x_k), v\rangle\,, \quad G_k = \langle \nabla f(x_k), x_k - s_k\rangle\,, \tag{2}$$

$$\ell_k = \max_{0\le i\le k}\{f(x_i) - G_i\}, \quad U_k = f(x_k) - \ell_k. \tag{3}$$

Convexity gives $0 \le \delta_k \le U_k \le G_k$. A valid known lower bound can be included in ℓ_k, but is not required. A zero gap certifies optimality. We stop at $U_k \le \varepsilon$. The best-so-far tangent construction is classical [28, Method 1, maximization convention]; the new analysis concerns its curvature-dependent cost, not the certificate itself.

The algorithm. Choose $M > 0$ and set $L_{-1} = M$. At each nonterminal source, put $d_k = s_k - x_k$, $q_k = \|d_k\|$, and start with $L = L_{k-1}/2$. Repeat

$$\alpha = \min\{1, G_k/(Lq_k^2)\}, \qquad f(x_k + \alpha d_k) \le f(x_k) - \alpha G_k + \tfrac{1}{2}L\alpha^2 q_k^2 \tag{4}$$

until the inequality passes, doubling L after each rejection. Store $L_k = L$, update $x_{k+1} = x_k + \alpha d_k$, and reuse the accepted value. Only trial function values are recomputed during the search. This is the standard quadratic backtracking scheme [2]. For M fixed, the algorithm does not need D, H_0, H_1, or f_Q^*. The optional $M = U_0/D^2$ uses a diameter bound and the first paid function, gradient and LMO calls.

Lemma 2.1 (Local acceptance)**.** *For a feasible descent direction d with $g = -\langle \nabla f(x), d\rangle > 0$ and feasible cap $a_{\max} > 0$, the test in (4) passes at $\alpha = \min\{a_{\max}, g/(L\|d\|^2)\}$ whenever $L \ge H_0 + H_1\delta(x)$. Every accepted trial, regardless of this sufficient threshold, satisfies*

$$f(x) - f(x + \alpha d) \ge \tfrac{1}{2}\alpha g. \tag{5}$$

Thus the search terminates and

$$L_k \le \max\{L_{k-1}/2,\ 2(H_0 + H_1\delta_k)\}. \tag{6}$$

The assertion also allows $a_{\max} > 1$.

Proof. Write $h(t) = f(x + td) - f(x)$ and $K = H_0 + H_1\delta(x)$. While $h \le 0$, the Hessian bound gives $h''(t) \le K\|d\|^2$. Initially $h'(0) = -g < 0$. If $K > 0$, a first return to zero before $\min\{a_{\max}, 2g/(K\|d\|^2)\}$ contradicts $h(t) \le -gt + K\|d\|^2 t^2/2 < 0$. The same integration proves the quadratic upper bound through the prescribed trial; the zero-K case follows with a linear upper model. Its curvature is at most L, so the test passes. For any accepted trial, $L\alpha\|d\|^2 \le g$, proving (5). If there is a rejection, the last rejected estimate is less than K, so the accepted one is less than $2K$; otherwise it is $L_{k-1}/2$. □

Consequently f_k and U_k are nonincreasing. The analysis uses curvature only on a descent segment, not a uniform upper model on every segment point.

Lemma 2.2 (Lag potential)**.** *Let source phase indices be nondecreasing in $\{0, \dots, J-1\}$. Let positive caps obey $C_j/2 \le C_{j+1} \le C_j$. Suppose an update with $L_k > C_j$ in its source phase must have $L_k = L_{k-1}/2$. Then the number of such lag updates is at most*

$$\lceil \log_2 \max\{1, M/C_0\}\rceil + J - 1.$$

The statement applies to every prefix and allows initially skipped phases.

Proof. Before an update in phase j, use the nonnegative integer potential

$$p = \max\{0, \lceil \log_2(L_{k-1}/C_j)\rceil\}.$$

A lag update reduces it by one; a non-lag update leaves zero potential before a phase change. A forward jump of m phases raises it by at most m. Starting from virtual phase zero, all jumps, including an initial jump, sum to at most $J - 1$. Nonnegativity proves the count. □

Theorem 2.3 (Observable stopping and total function cost)**.** *For $0 < \varepsilon < U_0$, let*

$$J = \lceil \log_2(U_0/\varepsilon) \rceil, \qquad P = \left\lceil \log_2 \max\{1, MD^2/U_0\} \right\rceil.$$

Ordinary backtracking reaches $U_N \le \varepsilon$ within

$$N \le P + \left\lceil (6 + 16H_1D^2)J + 32H_0D^2/\varepsilon \right\rceil \tag{7}$$

updates. It uses $N + 1$ gradients and LMOs, including the final check. For its actual $N \ge 1$, every function evaluation, including all rejected trials, is counted in

$$N_f \le 1 + 2N + \lceil \log_2 \max\{1, 2(H_0 + H_1U_0)/M\} \rceil. \tag{8}$$

For $N = 0$, one function, gradient and LMO evaluation suffice. These are exact-oracle and exact-arithmetic bounds.

Proof. Put $b_j = U_0/2^j$ and assign a nonterminal source to phase j when $b_j/2 < U_k \le b_j$, for $j = 0, \dots, J-1$. Use

$$C_j = \max\{2(H_0 + H_1 b_j),\ b_j/(2D^2)\}.$$

The caps decrease by at most a factor two. By Lemma 2.1, an accepted estimate above C_j must be a first-trial halving. Since $C_0 \ge U_0/(2D^2)$, Lemma 2.2 bounds lag updates by $P + J$.

For a non-lag update in phase j, $G_k \ge U_k > b_j/2$ and $q_k \le D$ imply

$$f_k - f_{k+1} \ge p_j := \min\{b_j/4,\ b_j^2/(8C_jD^2)\}.$$

Every update decreases f, and the total decrease within the phase is at most its initial suboptimality, hence at most b_j. Thus its non-lag updates number at most

$$\lfloor b_j/p_j \rfloor + 1 \le 5 + 16H_1D^2 + 16H_0D^2/b_j.$$

Summing and using $\sum_{j<J} 1/b_j < 2/\varepsilon$ proves (7). This bounds every unfinished prefix, which also establishes termination.

The local envelope and monotone U give $L_k \le \max\{M/2^{k+1}, 2(H_0 + H_1U_0)\}$. If r_k denotes rejections, $L_k = 2^{r_k - 1}L_{k-1}$, so exactly

$$\sum_{k<N} (r_k + 1) = 2N + \log_2(L_{N-1}/M)$$

trial values are computed. Add the initial value and use the envelope to obtain (8). Accepted values are cached. □

The last identity is standard [2]. For $M = U_0/D^2$, $P = 0$, and the function-count overhead is independent of ε. This normalization is optional for the order of the bound.

Proposition 2.4 (Initialization is visible even for affine objectives)**.** *Let $Q = [-1, 1]$, $f(x) = -cx$, $c > 0$, $x_0 = -1$, and $L_{-1} = M > 0$. For $0 < \varepsilon < 2c$, halving backtracking uses exactly*

$$N = \left\lceil \log_2 \left(1 + \frac{M}{c}(1 - \varepsilon/(2c)) \right) \right\rceil$$

updates. Thus halving U_0 may require $\Theta(\log(M/c))$ updates.

Proof. Every test accepts its first trial: $L_k = M/2^{k+1}$, $x_{k+1} = \min\{1, x_k + c/L_k\}$. The tangent lower bound is always $-c$, so $x_N = \min\{1, -1 + (2c/M)(2^N - 1)\}$ and $U_N = c(1 - x_N)$. Solving the stopping inequality gives the formula. □

This lower bound concerns the specified halving rule, not arbitrary first-order algorithms. With $M = U_0/D^2 = c/2$ the first step solves it.

Comparison with a global bound. For $f(x) = \cosh(x) - 1$ on $[-R, R]$, $H_0 = H_1 = 1$, while a global gradient-Lipschitz constant is $L_Q = \cosh R$. Starting at R gives $U_0 = 2R \sinh R$. For fixed tolerance and fixed M, our update bound grows as $O(R^3 + R^2/\varepsilon)$ as $R \to \infty$, rather than the exponential $L_Q D^2/\varepsilon$ term obtained by substituting $H_0 = L_Q, H_1 = 0$ in the same certificate theorem. This compares upper bounds, not a runtime or oracle lower bound; exact segment minimization solves this particular one-dimensional example in one step. Appendix A gives further step comparisons.

3 Away steps and geometric progress

Let $Q = \operatorname{conv}(\mathcal{A})$ for a finite atom set. Maintain $x = \sum_{a \in S} \lambda_a a$ with positive weights summing to one. Besides the full FW atom s, choose $v \in \arg\max_{a \in S} \langle \nabla f(x), a \rangle$. Use the larger-gap direction among $s - x$ and $x - v$, choosing FW on a tie. Its gap g satisfies $g \geq G \geq U$ and its norm is at most D. The feasible cap is 1 for FW and $\lambda_v/(1 - \lambda_v)$ for away; a singleton takes FW. Apply the same halving/doubling test with $\alpha = \min\{a_{\max}, g/(L \|d\|^2)\}$. This is standard away-step FW (AFW) [3, 2]. The certificate still uses the full FW gap. Active-set search is additional work, separate from full LMO calls.

An away step at its cap removes an atom and is called a drop; other steps are good. If m good updates occur, drops number at most the number of FW updates plus $|S_0| - 1$, hence

$$N \leq 2m + |S_0| - 1. \tag{9}$$

This follows because FW adds at most one atom and a drop removes one, while the active set remains nonempty. Full FW steps can remove more atoms and only improve the bound.

Theorem 3.1 (Convex AFW certificate bound)**.** *With J, P as in Theorem 2.3, AFW reaches $U_N \leq \varepsilon$ within*

$$N \leq 2\left[P + \left\lceil (6 + 16H_1 D^2)J + 32H_0 D^2/\varepsilon \right\rceil\right] + |S_0| - 1.$$

It uses $N + 1$ gradients and full LMOs. Its actual function-call count satisfies (8)*.*

Proof. Use the caps and phase potential from Theorem 2.3, counting lag across all updates. On a good away step the quadratic minimizer is not clipped, giving decrease $g^2/(2L \|d\|^2)$. A full FW step gives at least $g/2$; a nonfull FW step has the same quadratic bound. Thus controlled good steps have the same lower progress p_j as in that theorem. Drops also decrease f, so the phase budget still bounds the good steps by the ordinary FW count. Apply (9). Lemma 2.1, the accepted-estimate envelope and the trial-count identity remain valid across drops, proving the function bound. □

3.1 From primal progress to the observable threshold

For geometric convergence assume the selected gap satisfies

$$g_k^2 \geq 2\kappa\delta_k, \qquad \kappa > 0. \tag{10}$$

This is an extra hypothesis. For example strong convexity $\mu > 0$ and pyramidal width W give $\kappa = \mu W^2/4$. Quadratic growth relative to a possibly non-singleton solution set, $\delta(x) \geq (\gamma/2) \operatorname{dist}(x, X^*)^2$ on the initial sublevel set, gives $\kappa = \gamma W^2/16$. These are established polytope scaling implications; Appendix B records the constants and attribution [3, 16].

Freezing curvature at the source gives the two-branch quadratic envelope in Appendix B.1, by the established descent inversion [21, Eq. (15)]. Retaining its dependence on the function along the segment gives the exact envelope for our class.

Theorem 3.2 (Optimal pointwise FW-gap envelope)**.** *Suppose $H_1 > 0$, and write $a = D\sqrt{H_1}$ and $c = H_0/H_1$. At every feasible source,*

$$U \le G \le E(\delta) := \begin{cases} D\sqrt{2H_0\delta + H_1\delta^2}, & \delta \le c(\cosh a - 1), \\ a\delta \coth a + ac \tanh(a/2), & \delta > c(\cosh a - 1). \end{cases} \tag{11}$$

For every $D > 0$, $H_0 \ge 0$, $H_1 > 0$ and $\delta > 0$, a convex C^2 objective on a one-dimensional interval of diameter D attains equality. Thus no smaller bound depending only on these four quantities is valid for this class. The inverse envelope is

$$\eta_\varepsilon = \begin{cases} \dfrac{(\varepsilon/D)^2}{H_0 + \sqrt{H_0^2 + H_1(\varepsilon/D)^2}}, & \varepsilon \le ac \sinh a,\ H_0 > 0, \\ \dfrac{\varepsilon}{a} \tanh a - c(1 - \operatorname{sech} a), & \text{otherwise.} \end{cases} \tag{12}$$

Hence $\delta \le \eta_\varepsilon$ implies $U \le \varepsilon$. For $H_1 = 0$, put $A = D^2 H_0$: the limiting envelope is $\sqrt{2A\delta}$ for $2\delta \le A$, and $\delta + A/2$ otherwise; its inverse is $\varepsilon^2/(2A)$ for $\varepsilon \le A$, and $\varepsilon - A/2$ otherwise. At $H_0 = H_1 = 0$ these are $E(\delta) = \delta$ and $\eta_\varepsilon = \varepsilon$.

Proof. Set $h(t) = f(x + t(s - x)) - f_Q^*$ for $0 \le t \le 1$. Then $h \ge 0$, $h(0) = \delta$, $h'(0) = -G$, and $h'' \le D^2(H_0 + H_1 h) = a^2(h + c)$. Variation of constants, whose kernel $\sinh(a(t-u))/a$ is nonnegative for $u \le t$, gives

$$0 \le h(t) \le (\delta + c)\cosh(at) - (G/a)\sinh(at) - c.$$

Consequently, for every $0 < t \le 1$,

$$G \le a\frac{(\delta + c)\cosh(at) - c}{\sinh(at)}.$$

For $c > 0$ its derivative changes sign at $\cosh(at) = 1 + \delta/c$. If this point lies in the segment, the minimum is $a\sqrt{\delta(\delta + 2c)}$; otherwise it is the value at $t = 1$. For $c = 0$ the expression is decreasing and its minimum is $a\delta \coth a$. These are (11). The increasing branches meet at $\delta = c(\cosh a - 1)$, where $E = ac \sinh a$. Inversion gives (12). Appendix B.1 constructs attaining objectives for both branches, including convex C^2 extensions beyond the interval. For $H_1 = 0$, integrate $h'' \le D^2 H_0$ and minimize $\delta/t + D^2 H_0 t/2$ on $(0, 1]$. □

The word "optimal" here concerns a pointwise inequality within the specified function class, not oracle complexity or bibliographic priority. At $H_0 = 0$ the relation is linear, $G \le a\coth(a)\delta$. For fixed $H_0 > 0$ and small tolerance, $\eta_\varepsilon = \Theta(\varepsilon^2)$; this exponent cannot be removed in general: $f(x) = x^2/2$ on $[-1, 1]$, at $x = t > 0$, gives $\delta = t^2/2$ and $G = t(1 + t)$. At the first observation $U = G$. This is not an iteration lower bound or a restriction on stronger history-dependent certificates.

Theorem 3.3 (Geometric AFW certificate bound)**.** *Assume $H_0 + H_1 > 0$ and (10). Set*

$$J_* = \lceil \log_2(U_0/\eta_\varepsilon) \rceil, \qquad P_* = \lceil \log_2 \max\{1, M/[2(H_0 + H_1 U_0)]\} \rceil.$$

For $0 < \varepsilon < U_0$, the same algorithm reaches $U_N \le \varepsilon$ within

$$N \le 2\left\lceil P_* + (6 + 4H_0 D^2/\kappa)J_* + 8H_1 D^2 U_0/\kappa \right\rceil + |S_0| - 1. \tag{13}$$

Equation (8) remains valid. None of H_0, H_1, κ, f_Q^* is an algorithm input.




Figure 1: Pointwise theoretical bounds. The differential envelope retains the changing function value along the segment; the frozen quadratic comparison is (23). At zero base curvature, the exact linear coefficient is $\sqrt{b}\coth\sqrt{b}$. These are inequalities over a function class, not optimization trajectories or timing results.

Proof. Analyze true-error phases $b_j/2 < \delta_k \le b_j$ with $b_j = U_0/2^j$, until $\delta \le \eta_\varepsilon$. Their positive caps $C_j = 2(H_0 + H_1 b_j)$ decrease by at most a factor two. The initial true error may skip phases; Lemma 2.2 still bounds lag updates by $P_* + J_*$. On a controlled good step,

$$f_k - f_{k+1} \ge \delta_k \min\{1/2, \kappa/(C_j D^2)\} > \min\{b_j/4, \kappa b_j/(2C_j D^2)\}.$$

For a full FW step use $g \ge G \ge \delta$; for a nonclipped step use (10). The phase budget b_j permits at most $5 + 4H_0 D^2/\kappa + 4H_1 D^2 b_j/\kappa$ controlled good updates. Sum $\sum_{j<J_*} b_j < 2U_0$, add lag updates and apply (9). Theorem 3.2 ensures the actual certificate has passed by this point. □

For fixed problem and initialization the accuracy dependence is logarithmic. The explicit H_1 phase cost is proportional to U_0; H_1 also enters J_* through (12). Both AFW bounds apply when the geometric hypothesis holds; the initial U_0/κ contribution means the geometric bound is not uniformly tighter than the convex one. One need not assume $H_0 > 0$: the caps are positive at positive phase levels when $H_1 > 0$. A nonaffine example is $f = t + (\max\{t, 0\})^3$, $t = 1/2 - x$, on $[-1/2, 1/2]$, with $H_0 = 0, H_1 = 6$ and $\kappa = 1/16$ (Appendix B.1). In fact $H_0 = 0$ has stronger structural consequences, proved next without assuming the geometric inequality (10).

4 Zero base curvature: structure and finite termination

The endpoint $H_0 = 0$ is qualitatively different from a small positive H_0. The statements below concern the constrained gap $\delta = f - f_Q^*$, with the curvature condition holding throughout Q.

Proposition 4.1 (Exposed optimal face and a common gradient)**.** *Assume $H_0 = 0$. For any $x^* \in X^* := \arg\min_Q f$, put $g^* = \nabla f(x^*)$. Then*

$$X^* = \{x \in Q : \langle g^*, x - x^* \rangle = 0\}. \tag{14}$$

The gradient is the same g^* at every optimum, and for $x \in Q$,

$$\|\nabla f(x) - g^*\| \le \tfrac{1}{2} H_1 D \delta(x). \tag{15}$$

In particular, an ambient stationary optimum $g^* = 0$ forces f to be constant on Q. A nonconstant function in this zero-base class cannot have a relative-interior optimum either.

Proof. Optimality and convexity give $w := \langle g^*, x - x^* \rangle \geq 0$ and $\delta(x) \geq w$. For $h(t) = \delta(x^* + t(x - x^*))$, the curvature condition gives $h'' \leq H_1 \|x - x^*\|^2 h$, with $h(0) = 0$ and $h'(0) = w$. The same positive-kernel comparison as in Theorem 3.2 yields

$$0 \leq h(t) \leq w \frac{\sinh(rt)}{r}, \qquad r = \sqrt{H_1}\, \|x - x^*\|,$$

where the quotient is t at $r = 0$. Thus $w = 0$ implies $h(1) = 0$, proving (14). A segment joining two optima lies in X^*; its Hessian is zero by the assumed norm bound, so integration shows that their gradients coincide. Along a segment from x^* to x, convexity gives $h(t) \leq t\delta(x)$. Integrating the Hessian norm therefore gives

$$\|\nabla f(x) - g^*\| \leq H_1 \|x - x^*\| \int_0^1 h(t)\, dt \leq \tfrac{1}{2} H_1 D \delta(x).$$

At a relative-interior optimum, the linear functional g^* is constant on the affine hull of Q, so (14) gives $X^* = Q$. □

Suppose now that $Q = \mathrm{conv}(\mathcal{A})$ is a polytope with a finite atom set, and that f is nonconstant on Q. Write $\mathcal{A}_* := \mathcal{A} \cap X^*$ and

$$m := \min_{v \in \mathcal{A} \setminus \mathcal{A}_*} \langle g^*, v - x^* \rangle > 0. \tag{16}$$

For any active representation $x = \sum \lambda_v v$, its mass outside the optimal face obeys

$$\theta := \sum_{v \notin \mathcal{A}_*} \lambda_v \leq \frac{\langle g^*, x - x^* \rangle}{m} \leq \frac{\delta(x)}{m}. \tag{17}$$

Replacing its nonoptimal atoms by any point of X^* shows $\mathrm{dist}(x, X^*) \leq D\theta$. Hence the model itself implies the global weak-sharpness bound

$$\delta(x) \geq (m/D)\, \mathrm{dist}(x, X^*).$$

An independent quadratic-growth hypothesis is unnecessary here. The finite atom assumption matters for this error bound: for the affine function $f(x) = 1 - x_1$ on the unit disk, $\delta(\cos t, \sin t)/\,\mathrm{dist}((\cos t, \sin t), X^*) = |\sin(t/2)| \to 0$. Zero base curvature on a general compact convex set therefore does not imply weak sharpness.

Theorem 4.2 (Finite exact termination of backtracking AFW)**.** *Assume $H_0 = 0$, a finite atom set, and $U_0 > 0$. Set*

$$b = H_1 D^2, \quad c_b = \begin{cases} \sqrt{b} \coth \sqrt{b}, & b > 0, \\ 1, & b = 0, \end{cases} \qquad \overline{L} = \max\{M, 2H_1 U_0\},$$

$$\tau = \min\left\{ \frac{U_0}{2}, \frac{m}{4(1 + b + c_b)} \right\}, \qquad J_0 = \lceil \log_2(U_0/\tau) \rceil. \tag{18}$$

Define the two initialization potentials

$$P = \left\lceil \log_2 \max\{1, MD^2/U_0\} \right\rceil, \qquad Q_0 = \left\lceil \log_2 \max\{1, 8\overline{L} D^2 \tau / m^2\} \right\rceil.$$

The standard backtracking AFW algorithm, continued with stopping tolerance zero, reaches an exact optimum and $U_N = 0$ within

$$N \leq 2\lceil P + \lceil (6 + 16b) J_0 \rceil + Q_0 \rceil + |S_0| - 1 \tag{19}$$

updates. After the first source with $U \le \tau$, every nonoptimal update selects away, and at most Q_0 further good updates occur before only maximal drops of nonoptimal atoms remain. The gradient and full-LMO counts are $N+1$, and

$$N_f \le 1 + 2N + \lceil \log_2 \max\{1, 2H_1U_0/M\} \rceil .$$

Neither m, H_1, f_Q^* nor the threshold τ is an algorithm input. No hypothesis (10) *or positive strong-convexity constant is required. These are exact-arithmetic statements.*

Proof. The accepted-curvature envelope gives $L_k \le \overline{L}$ for all updates. Consider a nonoptimal source with $U \le \tau$; then $0 < \delta \le \tau$. By (15), the difference between the current and optimal gradient scores of any two atoms is at most $b\delta/2$. Thus every nonoptimal atom has a strictly larger current score than every optimal atom. An LMO returns an optimal atom. A nonoptimal source has some active nonoptimal atom, and an away maximizer v is necessarily nonoptimal. Its gap satisfies

$$g_{\mathrm{A}} = \langle \nabla f(x), v - x \rangle \ge m - \delta - b\delta/2 \ge 3m/4.$$

Meanwhile Theorem 3.2 gives $G \le c_b\delta \le m/4$. The algorithm therefore selects away. Its atom weight is at most $\theta \le \delta/m < 1/2$, so

$$\alpha_{\max} \le 2\delta/m.$$

Put $R = m^2/(8D^2\tau)$. Since

$$2H_1\tau/R = 16b\tau^2/m^2 \le b/(1+b+c_b)^2 \le 1,$$

an accepted estimate above R must be a first-trial halving; an acceptance after rejection is below $2H_1\delta \le R$. At entry the inherited estimate is at most $\overline{L}$, so there are at most Q_0 such halvings. An accepted estimate at most R remains at most R thereafter by the local envelope, and satisfies

$$L_k \|d\|^2 \alpha_{\max} \le 2RD^2\delta/m \le m/4 < g_{\mathrm{A}}.$$

Every such trial is clipped at the feasible cap and drops a nonoptimal atom. The objective and U decrease, so the local conditions remain valid. After finitely many drops all active atoms are optimal. Then $G = 0$ and the paid final check gives $U = 0$.

To count the preceding updates, apply the phase proof of Theorem 3.1 up to its first source $U \le \tau$. It bounds the total number of good updates before that source by $P + \lceil (6+16b)J_0 \rceil$. The local argument permits at most Q_0 further good updates. The global atom-removal count (9) proves (19). The function-call identity and envelope give the stated oracle bound, including rejections. □

Here J_0 does not depend on M. For a fixed objective and start, large M therefore contributes only $O(\log M)$ through P, Q_0, with no additional factor H_1D^2 on that initialization cost.

Finite active-set identification under strict complementarity is established theory [7, Theorem 3.1 and Appendix] and is not a new general phenomenon. Garber [8, Theorems 4–5] also gives finite-convergence and face-identification results for line-search FW variants. Here the zero-base model itself forces the exposed face to equal the entire solution set; identifying that face solves the problem exactly. The displayed threshold and trial count specialize this mechanism to the inherited halving/doubling curvature estimate, without requiring convergence to a particular optimum. This does not assert priority over every identification theorem.

Why the endpoint matters. The conclusion does not extend to an arbitrary $H_0 > 0$. On $[-1, 1]$, take $f(x) = hx^2/2$, $h > 0$, $0 < x_0 < 1$, and $M = 3h$. Backtracking accepts $L_k = 3h/2$ at every update: the first trial accepts initially, and each later trial $3h/4$ fails for this nonzero quadratic step. The selected direction is FW (in the usual two-endpoint representation), and

$$x_{k+1} = x_k - f'(x_k)/(3h/2) = x_k/3.$$

Hence no finite iterate is optimal, however small $h = H_0 > 0$ is. Finite identification of an optimal face does not in general solve a nonconstant objective on that face.

5 Experiments: initialization, geometry and stopping

The experiments test finite-budget behavior rather than tightness of the worst-case constants. All methods use the same scalar certificate unless a stopping control is explicitly identified. We distinguish function, gradient and full LMO budgets; no arbitrary conversion weights or wall-clock ranking are imposed. AFW active-set scans and atom scores are recorded separately. Capped runs remain in summaries.

5.1 Count-data objectives with verified curvature bounds

The main real-data protocols use Seoul Bike Sharing Demand [22] and Metro Interstate Traffic Volume [23], with the unmodified official files, hashes and CC BY 4.0 attribution retained. A stable chronological sort and the first 80% of rows define training data. Non-intercept features are transformed by $\tanh((v - \bar{v}_{\text{train}})/s_{\text{train}})$, using training statistics only; an intercept is included and constrained. Repeated source timestamps keep their original weights. We optimize a training objective, not a predictive-performance criterion.

With normalized response $t_i = y_i/\bar{y}_{\text{train}}$ and $c_i = t_i \log t_i - t_i$ (with $0 \log 0 = 0$), the objective is

$$f(x) = \frac{1}{m} \sum_i \{e^{a_i^\top x} - t_i a_i^\top x + c_i\} + \frac{\mu}{2} \|x\|^2, \qquad \|x\|_1 \le R, \quad \mu = 0.01. \tag{20}$$

All coordinates, including the intercept, are ridge-regularized. Since $\|a_i\|_\infty = 1$ and $\text{mean}(t_i) = 1$, valid constants relative to the constrained optimum are

$$H_1 = \max_i \|a_i\|^2, \qquad H_0 = \mu + H_1(1 + R), \qquad L_Q = \mu + e^R \lambda_{\max}(A^\top A/m). \tag{21}$$

The Hessian norm is at most $\mu + H_1 \,\text{mean}(e^{a_i^\top x})$. Since $f_Q^* \le f(0)$, the objective formula gives

$$\text{mean}(e^{a_i^\top x}) \le \delta + 1 + R.$$

These are analytic upper bounds, not fitted or minimal constants. On a compact domain a global Lipschitz bound also exists.

Table 1: Count-data protocols. R_* is the offline unconstrained reference ℓ_1 norm; tested radii are $R_*/2$ and $2R_*$. Widths are numerical reference brackets, without directed rounding. All $2d$ signed coordinate vertices are used at each radius.

Data	Training rows	d	R_*	H_1	Max. width
Seoul	7008	15	4.575821	8.897	4.2e-08
Metro	38563	12	1.431745	6.208	2.3e-09

An offline damped Newton reference estimates $R_* = \|x^*_{\text{unc}}\|_1$. The radii $R_*/2$ and $2R_*$ give active and interior controls, with independently recomputed constrained numerical references.

Each reference uses a feasible value as an upper bound and its convex tangent/LMO lower bound. These are numerical brackets without directed rounding, not formal interval certificates. They are never solver inputs. We report post-run error bounds to this numerical precision. All $2d$ signed coordinate vertices are used once at each radius: 108 dataset/radius/start cases in total. Radii and starts within a dataset share training data and are not independent datasets.

5.2 Common function budgets and failed runs

Four standard BT variants combine FW/AFW with $M = 1$ or U_0/D^2. Two informed short-step controls use $L = H_0 + H_1 U_k$ with FW/AFW, denoted Q. They receive the constants in (21); BT does not. The original six-method archive has 648 runs. We add 108 AC-FW runs on exactly the same cases, preserving all earlier records.

AC-FW is an independent implementation of Algorithms 1–2 of [18], with its Eq. (6), parameter $\delta = 1$. The compulsory initial LMO vertex move and both initial function and gradient calls are charged. A rejected trial consumes an outer iteration and a function call, retaining the old gradient; full LMOs are also charged. The cited global gradient-Lipschitz assumption on $\mathbb{R}^d$ is not satisfied by (20), so these compact-domain runs are an empirical comparison rather than a direct application of its global theorem.

The target is $U \leq 10^{-5}$. BT/Q have a cap of 1000 updates; AC has 1000 outer trials, including rejections. The primary comparison instead uses at most 1000 function calls: every nonterminal run can reach that budget before its cap, while successful runs are carried forward. Each prefix is the last fully paid source before the budget, reconstructed from the full trace rather than interpolated between sampled rows. Other oracle types are unrestricted in a single-oracle comparison.

Table 2: Certified starts at the common budget of 1000 function calls. Every nonterminal run reaches this budget before its cap; completed runs are carried forward. Seoul has 30 starts per cell, Metro 24. F/A: FW/AFW; subscripts $1, U$: initial $M = 1, U_0/D^2$; Q: known-curvature short step; AC: auto-conditioned FW with charged preprocessing.

Data	R/R_*	F_1	F_U	A_1	A_U	F_Q	A_Q	AC
Seoul	0.5	0	0	30	30	0	0	0
Seoul	2	0	0	0	0	0	0	0
Metro	0.5	2	3	24	24	1	24	13
Metro	2	0	0	0	0	0	0	0

At the active radii, both AFW-BT initializations certify every start at 1000 function calls. FW-BT certifies none on Seoul and only 2/24 or 3/24 on Metro. AC-FW obtains 13/24 active Metro successes and none on active Seoul. The known-curvature AFW-Q succeeds on all active Metro starts but none on active Seoul, illustrating conservatism of the analytic curvature parameters. At the interior radii no method in this seven-method scalar-certificate comparison reaches the target within 1000 function calls. End-of-cap results and censored call medians are reported separately in Table 5.

The references separate remaining error from stopping lag. At the interior Metro radius, all 24 starts of each BT variant have numerical reference error upper bound at most 10^{-5} by the original endpoint, but FW has no scalar-certificate successes and AFW has only 6/24 or 7/24. On interior Seoul, the corresponding reference-error condition holds for 12/30 and 18/30 AFW starts, while neither scalar BT variant stops. An unsuccessful certificate is therefore not equivalent to a large remaining primal error.




Figure 2: Median scalar certificates over all distinct vertices at function-call budgets. Curves include failed runs and charged AC-FW preprocessing; completed runs are carried forward. The dotted horizontal line is 10^{-5}. BT has no curvature inputs, Q receives analytic H_0, H_1. These are paired finite cases, not confidence bands for a population of problems.

5.3 Controlling initialization and objective units

A separate 2×2 study controls the original point versus the compulsory LMO vertex, and AC versus BT updates. All four cells pay the same two-source preprocessing, retain both tangent bounds, and start their first trial at the same secant curvature. Only AC/vertex is the native published comparator; the other cells isolate factors. The study contains 280 runs on 70 earlier matched cases and introduces no new datasets. Table 3 shows all three planted active families, including the unfavorable scaled family. Their construction and the active-face slack identity appear in Appendix D.

Table 3: Starting-point control on three planted active families, ten matched instances each. Counts are certified starts at 1000 function calls. O/V: original point/compulsory initial LMO vertex. The last column is the descriptive paired median BT-V/AC-V certificate ratio and 95% percentile-bootstrap interval, not a speedup estimate.

Family	AC-O	AC-V	BT-O	BT-V	Ratio [interval]
active	1	10	0	10	1.049 [0.935, 1.257]
active_scaled	0	0	0	0	1.530 [0.822, 1.750]
active_highdim	6	8	2	8	1.231 [0.798, 2.117]

On the active family, changing the starting point raises successes from 1 to 10 for AC and from 0 to 10 for BT. The descriptive BT/AC certificate-ratio interval from the vertex includes one. The scaled family remains unsolved by all four controls. Thus the earlier vertex-start comparison does not isolate an update-rule benefit.

Objective units provide another control. Under $\widetilde{f} = af + b$, $a > 0$, scaling M and ε by a preserves the exact iterates, trials and stopping index, since G, U, L scale together. Thus $M = U_0/D^2$ is automatically equivariant; fixing its numerical value instead changes effective initialization. A separate real-data radius/scale study verifies the counts in all 60 paired scale comparisons for each certificate-initialized method. This elementary property does not establish a faster algorithm.

5.4 Causal stopping controls on fixed BT paths

We repeat the 432 original BT paths from the count-data study, with exactly matching original endpoints and oracle counts. Five causal monitors give 2160 dependent stopping profiles. Disjoint blocks of $K = 8, 32, 128$ paid affine tangents provide a lower bound using one additional full LMO per completed block; the scalar envelope is retained. A separate standard control receives $\mu = 0.01$ and uses $f - \|\nabla f\|^2/(2\mu)$ as a lower bound. Appendix C proves validity and isolated oracle accounting; block sizes are fixed before outcomes and no monitor reads the reference.

Table 4: Isolated causal controls on 432 exactly repeated BT paths, giving 2160 dependent profiles. Counts use each profile's first stop or the original cap. More LMOs compares total direction plus certificate LMOs with scalar. Known μ receives the analytical ridge coefficient; blocks do not.

Profile	Certified	Earlier prefix	More total LMOs
scalar	129	0	0
block_8	129	0	432
block_32	129	0	425
block_128	129	0	380
known_mu	226	110	0

None of the tested blocks gives an earlier stop. The known-μ control gives 226 successes instead of 129 and 110 earlier prefixes, with no extra LMOs but additional model information. On interior Metro each BT variant now reaches 24/24 successes by the original endpoint; both AFW variants do so within 1000 function calls. Interior Seoul remains difficult: AFW obtains only 4/30 or 10/30 successes by the endpoint and none at 1000 function calls. These controls neither establish adaptation to unknown strong convexity nor evaluate the exact averaged-gradient Fenchel dual in [2]. They show that observable stopping cost is a separate practical issue.

6 Discussion

The phase argument replaces a single global curvature bound by a sequence of source-level bounds while retaining the full feasible-region LMO. Its output is a guarantee for the accumulated scalar certificate and a count of every rejected function trial. This distinction matters when a small primal error is insufficient for an algorithm to certify stopping. The initialization term is intrinsic to the specified halving rule, as the affine example shows.

The experiments support controls on initialization, objective units, and active geometry. They do not establish a universal practical ranking. Positive H_1 on a compact domain does not preclude a global Lipschitz constant, and the analytic constants can be conservative. The logarithmic AFW theorem for general positive base curvature requires its additional geometric inequality. At zero base curvature the model itself supplies an exposed optimal face and finite termination; this is a constrained boundary regime, not a general model of a nonconstant loss with an attained interior minimum. Exact-arithmetic bounds do not by themselves ensure finite-precision reliability; deterministic oracle corrections in Appendix C.1 only establish certificate validity. Extending the analysis to controlled noisy directions, pairwise swaps, and adaptive sliding with all inner linear oracles counted remains open.

References

[1] G. Braun, A. Carderera, C. W. Combettes, H. Hassani, A. Karbasi, A. Mokhtari, and S. Pokutta. Conditional Gradient Methods. arXiv:2211.14103v5, 2025.

[2] F. Pedregosa, G. Negiar, A. Askari, and M. Jaggi. Linearly Convergent Frank-Wolfe with Backtracking Line-Search. *AISTATS*, 2020. arXiv:1806.05123.

[3] S. Lacoste-Julien and M. Jaggi. On the Global Linear Convergence of Frank-Wolfe Optimization Variants. *NeurIPS*, 2015. arXiv:1511.05932.

[4] F. Alimisis, R. Islamov, and A. Lucchi. Why Do We Need Warm-up? A Theoretical Perspective. arXiv:2510.03164v2, 2026.

[5] S. Kale, J. D. Lee, C. De Sa, A. Sekhari, and K. Sridharan. From Gradient Flow on Population Loss to Learning with Stochastic Gradient Descent. arXiv:2210.06705v1, 2022.

[6] D. Y. Cao, A. Y. Chen, K. Sridharan, and B. Tang. Efficiently Escaping Saddle Points under Generalized Smoothness via Self-Bounding Regularity. arXiv:2503.04712v3, 2026.

[7] I. M. Bomze, F. Rinaldi, and D. Zeffiro. Active set complexity of the Away-step Frank-Wolfe Algorithm. arXiv:1912.11492, 2019.

[8] D. Garber. Revisiting Frank-Wolfe for Polytopes: Strict Complementarity and Sparsity. arXiv:2006.00558, 2020.

[9] A. Lobanov and A. Koloskova. Avoiding Bias in Clipped SGD for Overparameterized Models under Generalized Smoothness. arXiv:2605.14800v2, 2026.

[10] A. Lobanov. A Few Accelerated Algorithms for Convex Optimization under (H_0, H_1)-Smoothness. arXiv:2608.04884, 2026.

[11] A. A. Vyguzov and F. S. Stonyakin. Frank-Wolfe Algorithms for (L_0, L_1)-smooth functions. arXiv:2510.16468v5, 2026.

[12] P. Dvurechensky, K. Safin, S. Shtern, and M. Staudigl. Generalized Self-Concordant Analysis of Frank-Wolfe Algorithms. arXiv:2010.01009, 2020.

[13] S. Takahashi, S. Pokutta, and A. Takeda. Fast Frank–Wolfe Algorithms with Adaptive Bregman Step-Size for Weakly Convex Functions. *ICLR*, 2026. arXiv:2504.04330v4.

[14] S. Takahashi. Adaptive Conditional Gradient Sliding: Projection-Free and Line-Search-Free Acceleration. arXiv:2601.20443v2, 2026.

[15] E. Wirth, J. Peña, and S. Pokutta. Fast convergence of Frank-Wolfe algorithms on polytopes. arXiv:2406.18789v3, 2025.

[16] T. Kerdreux, A. d'Aspremont, and S. Pokutta. Restarting Frank-Wolfe: Faster Rates under Hölderian Error Bounds. arXiv:1810.02429v4, 2021.

[17] R. Zhao. New Analysis of an Away-Step Frank-Wolfe Method for Minimizing Logarithmically-Homogeneous Barriers. arXiv:2305.17808v4, 2023.

[18] K.-H. Giang-Tran, S. Shafiee, and N. Ho-Nguyen. Auto-Conditioned Frank-Wolfe Algorithms. arXiv:2605.15512, 2026.

[19] A. Khademi and A. Silveti-Falls. Adaptive Conditional Gradient Descent. arXiv:2510.11440v1, 2025.

[20] S. Yagishita and M. Ito. Simple linesearch-free first-order methods for nonconvex optimization. arXiv:2509.14670, 2025.

[21] S. Lacoste-Julien. Convergence Rate of Frank-Wolfe for Non-Convex Objectives. arXiv:1607.00345, 2016.

[22] Seoul Bike Sharing Demand [Dataset]. UCI Machine Learning Repository, 2020. doi:10.24432/C5F62R. CC BY 4.0.

[23] J. Hogue. Metro Interstate Traffic Volume [Dataset]. UCI Machine Learning Repository, 2019. doi:10.24432/C5X60B. CC BY 4.0.

[24] H. Karimi, J. Nutini, and M. Schmidt. Linear Convergence of Gradient and Proximal-Gradient Methods Under the Polyak–Łojasiewicz Condition. arXiv:1608.04636v4, 2020.

[25] W. Wolberg, O. Mangasarian, N. Street, and W. Street. Breast Cancer Wisconsin (Diagnostic) [Dataset]. UCI Machine Learning Repository, 1993. doi:10.24432/C5DW2B.

[26] H. Fanaee-T. Bike Sharing [Dataset]. UCI Machine Learning Repository, 2013. doi:10.24432/C5W894. CC BY 4.0.

[27] Y. Liu, Y. Ge, R. Pan, K. An, and T. Zhang. Theoretical Analysis on how Learning Rate Warmup Accelerates Convergence. arXiv:2509.07972v1, 2025.

[28] R. M. Freund and P. Grigas. New Analysis and Results for the Frank-Wolfe Method. arXiv:1307.0873, 2013.

[29] D. Martínez-Rubio and S. Pokutta. Beyond Short Steps in Frank-Wolfe Algorithms. arXiv:2501.18773v1, 2025.

[30] G. Yuan. Adaptive Lipschitz-Free Conditional Gradient Methods for Stochastic Composite Nonconvex Optimization. arXiv:2603.06369v1, 2026.

[31] S. Takahashi and A. Takeda. Beyond Polytopes: Damped Newton Frank–Wolfe Methods over Compact Convex Sets. arXiv:2609.34290v1, 2026.

A Comparison steps and certificate normalization

If H_0, H_1 are supplied, consider

$$\alpha = \min\{1, G/[(H_0 + H_1 U)q^2]\},$$

with a full step when its curvature is zero. Lemma 2.1's descent argument, with $U \geq \delta$, gives

$$f_k - f_{k+1} \geq \frac{1}{2}\min\{U_k, U_k^2/[D^2(H_0 + H_1 U_k)]\}.$$

In a phase $b/2 < U \leq b$, progress is at least $\min\{b/4, b^2/(8D^2H_0 + 4D^2H_1 b)\}$, because the right-hand side is increasing in $U > 0$. The phase budget therefore permits at most $5 + 4H_1D^2 + 8H_0D^2/b$ updates. Summing the same dyadic levels proves

$$N \leq \left\lceil (5 + 4H_1D^2)J + 16H_0D^2/\varepsilon \right\rceil. \tag{22}$$

There are $N+1$ calls of each oracle for this explicit step. Exact segment minimization performs at least as well as the comparison step, so it inherits (22) without receiving H_0, H_1. This does not bound the inner function or gradient calls needed for an exact scalar minimizer. The expanded technical report also derives a hyperbolic step with the same order; its elementary function is not needed for Theorem 2.3.

For completeness, certificate-scaled backtracking starts with $\theta_{-1} = 1$, tries $\theta_{k-1}/2$, sets $L = \theta U_k/D^2$, and doubles θ on a failed quadratic test. The sufficient threshold $T_k = D^2(H_0/U_k + H_1)$ is nondecreasing. By induction,

$$\theta_k \leq \max\{2^{-k-1}, 2T_k\}, \qquad L_k \leq \max\{U_k/(2^{k+1}D^2), 2(H_0 + H_1U_k)\}.$$

Indeed a first-trial acceptance halves the previous bound, and after a rejection the accepted estimate is less than $2T_k$. Thus $L_k \leq 2H_0 + (2H_1 + 1/(2D^2))U_k$. Applying the preceding phase argument with these effective constants gives

$$N \leq \left\lceil (7 + 8H_1D^2)J + 32H_0D^2/\varepsilon \right\rceil.$$

The recursion $\theta_k = 2^{r_k - 1}\theta_{k-1}$ gives

$$N_f \leq 1 + 2N + \left\lceil \log_2 \max\{1, 2D^2(H_0/\varepsilon + H_1)\} \right\rceil.$$

The last source before an update has $U > \varepsilon$, which bounds θ_{N-1}. This normalization requires D and is optional for the two-phase order. Theorems 2.3 and 3.1 concern ordinary backtracking, rather than this separate variant.

B Geometric implications and attaining examples

Let $p = \langle \nabla f(x), v - s \rangle$ be the pairwise gap over the full atom set and current active set. The selected FW/away gap is at least $p/2$. The standard polytope scaling inequality gives, for any optimal $x^* \neq x$,

$$p \geq W \frac{\langle \nabla f(x), x - x^* \rangle}{\|x - x^*\|}.$$

See [3, Theorem 3] and [16, Lemmas 3.4 and 3.6]. For quadratic growth, choose a nearest optimum. Convexity and the growth inequality imply $p \geq W\delta/\operatorname{dist}(x, X^*) \geq W\sqrt{\gamma\delta/2}$. Hence $g^2 \geq \gamma W^2 \delta/8$, giving $\kappa = \gamma W^2/16$. Monotone iterates stay in the initial sublevel set where growth is assumed. No uniqueness is required. For strong convexity, $\langle \nabla f(x), x - x^* \rangle \geq \delta + \mu \|x - x^*\|^2/2$; minimizing $\delta/r + \mu r/2$ over $r > 0$ gives $p \geq W\sqrt{2\mu\delta}$ and $\kappa = \mu W^2/4$. These geometric implications are prior results.

B.1 Gap-envelope equality and zero base curvature

The quadratic comparison. For $C = D^2(H_0 + H_1\delta)$, the feasible step $t = \min\{1, G/C\}$ and Lemma 2.1 give $\delta \geq G^2/(2C)$ if $G \leq C$, and $\delta \geq G - C/2$ otherwise. The full step is used when $C = 0$. Inverting this established descent inequality gives

$$G \leq \widehat{B}(\delta) := \begin{cases} \sqrt{2C\delta}, & 2\delta \leq C, \\ \delta + C/2, & 2\delta > C. \end{cases} \tag{23}$$

Theorem 3.2 retains curvature variation along the segment and strengthens this comparison when $H_1 > 0$.

Attainment at $H_1 = 0$. Let $Q = [-D/2, D/2]$, $x = D/2$ and $f(y) = h(y - z)^2/2$, $h > 0$. For $z \in Q$, $\delta = h(x - z)^2/2$ and $G = hD(x - z) = \sqrt{2hD^2\delta}$, attaining the square-root branch of (23) with $H_0 = h, H_1 = 0$. For $z < -D/2$, the constrained optimum is $-D/2$, $\delta = -hDz$ and $G = \delta + hD^2/2$, attaining its full-step branch. These examples also attain the $H_1 = 0$ limit of the exact envelope.

Attainment at $H_1 > 0$. Write $k = \sqrt{H_1}$, $a = kD$, $c = H_0/H_1$, and use $Q = [0, D]$ with source $x = 0$. If $c > 0$ and $0 < \delta \leq c(\cosh a - 1)$, set

$$r = k^{-1} \operatorname{arcosh}(1 + \delta/c), \qquad f(u) = c\big(\cosh(k(u - r)) - 1\big).$$

This is globally convex, $f_Q^* = f(r) = 0$, $f(0) = \delta$, $f'' = H_0 + H_1 f$, and the source gradient is negative. Thus its FW atom is D and $G = -Df'(0) = D\sqrt{2H_0\delta + H_1\delta^2}$.

For $\delta > c(\cosh a - 1)$, including all $\delta > 0$ when $c = 0$, write $E = E(\delta)$ and, for $u \leq D$, set

$$f(u) = (\delta + c)\cosh(ku) - \frac{E}{a}\sinh(ku) - c.$$

On $[0, D]$ it satisfies $f(D) = 0$ and

$$f'(D) = k\frac{c\cosh a - (\delta + c)}{\sinh a} < 0.$$

The identity $f'' = H_0 + H_1 f$ and this endpoint derivative imply that f is decreasing and nonnegative on the interval: explicitly, $f(u) + c = c\cosh(k(D - u)) - [f'(D)/k]\sinh(k(D - u))$ for $u \leq D$. It is convex on the entire left half-line. For $u > D$, extend by

$$f(u) = f'(D)(u - D) + \tfrac{1}{2}H_0(u - D)^2.$$

The two pieces have matching value, derivative and second derivative at D, so the extension is globally convex and C^2. Its constrained optimum is D, $f(0) = \delta$, and $G = -Df'(0) = E$. At $H_0 = 0$ this becomes $f(u) = \delta \sinh(k(D-u))/\sinh a$ on $u \le D$, with an affine extension beyond D. These constructions prove optimality for every parameter tuple, rather than only for the frozen quadratic model. Since (23) applies to each attainer, they also prove $E(\delta) \le \widehat{B}(\delta)$ and hence that the new threshold is never smaller than the quadratic one.

An exact global self-bounding corollary. Under the different assumption that the Hessian condition holds on all of $\mathbb{R}^d$ relative to a finite global infimum f^*, one has

$$\|\nabla f(x)\|^2 \le 2H_0(f(x) - f^*) + H_1(f(x) - f^*)^2. \tag{24}$$

Indeed apply the positive-kernel comparison on the entire ray $x - t\nabla f(x)/\|\nabla f(x)\|$, $t > 0$. Nonnegativity of $f - f^*$ bounds the initial directional derivative by

$$\sqrt{H_1} \inf_{t>0} \frac{(f(x) - f^* + H_0/H_1)\cosh(\sqrt{H_1}t) - H_0/H_1}{\sinh(\sqrt{H_1}t)} = \sqrt{2H_0(f(x) - f^*) + H_1(f(x) - f^*)^2}.$$

Use the quadratic limit if $H_1 = 0$, and a limit $t \to \infty$ if $H_0 = 0 < H_1$. The bound is attained by the global cosh objective above when $H_0, H_1 > 0$, by a quadratic when $H_1 = 0 < H_0$, and by $f(x) = A\exp(\sqrt{H_1}x)$, $A > 0$, when $H_0 = 0 < H_1$; the latter infimum is not attained. This gives exact constants for the global model. The bound in [4, Appendix G, Lemma 6] has the larger coefficients $9H_0/4$ and $27H_1/4$; that lemma does not claim sharpness. For the affine second-order bound, [6, Corollary 1] instead gives $\|\nabla f\|^2 \le 2(H_0 + H_1\delta)^2 \log(1 + H_1\delta/H_0)/H_1$ when $H_0, H_1 > 0$. The exact polynomial expression in (24) removes this logarithmic enlargement. The self-bounding principle itself is prior work [5, Lemma 11]; only the displayed constants and extremizers are being compared here. The constrained model alone does not imply (24): a nonconstant affine function on a compact interval has $H_0 = H_1 = 0$ and nonzero gradient.

A polynomial zero-base example. The globally convex C^2 example $f(x) = t + (\max\{t, 0\})^3$, $t = 1/2 - x \in [0, 1]$ on Q, has $f''(x) = 6t \le 6f(x)$, so $H_0 = 0, H_1 = 6$. It is not strongly convex. At a nonoptimal interior point both endpoint atoms are active, and the selected gap is $(1 + 3t^2)\max\{t, 1-t\} \ge 1/2$; the nonoptimal endpoint also satisfies this inequality. Since $\delta \le 2$, condition (10) holds with $\kappa = 1/16$. At zero error the geometric inequality is immediate. This shows the $H_0 = 0$ extension is nonvacuous.

C Lower-bound controls and deterministic oracle errors

Write $g_j = \nabla f(x_j)$ and $a_j = f(x_j) - \langle g_j, x_j\rangle$. For a completed block B of K consecutive paid sources,

$$b_B = \bar{a}_B + \min_{v \in Q} \langle \bar{g}_B, v\rangle \le f_Q^*.$$

Convexity gives $f(v) \ge a_j + \langle g_j, v\rangle$ for each source; average and minimize to prove validity. Disjoint blocks begin at source zero, and an incomplete block is not queried. The new lower bound is the maximum of ℓ_k and all completed b_B. Its certificate is nonincreasing and at most U_k on the same monotone path, so its first crossing occurs no later in updates, function or gradient calls. At source k, exactly $\lfloor (k+1)/K \rfloor$ additional full LMOs have been performed. Total LMO cost need not decrease. The monitors affect neither the line search nor the initial M; each reported cost is for one isolated monitor, excluding the other diagnostic monitors. An affordable budget source pays its entire scheduled verification.

These uniform block bounds differ from the averaged-gradient Fenchel dual in [2, Theorem 3]. When the global conjugate exists, Fenchel equality and Jensen give

$$b_B = -\overline{f^*(g_j)} - \sigma_Q(-\bar{g}_B) \le -f^*(\bar{g}_B) - \sigma_Q(-\bar{g}_B).$$

Here f^* denotes the conjugate, not the scalar optimum f_Q^*. No transfer of that paper's theorem or averaging weights is asserted. With a valid supplied strong-convexity constant μ, $f(x_j) - \|g_j\|^2/(2\mu) \le f_Q^*$ follows by minimizing the quadratic lower model over $\mathbb{R}^d$ [24, Appendix B]. Caching these bounds costs no LMO, but uses additional information. Native stopping checks on one synthetic interior instance verify all 20 method/profile pairs against the attainable-prefix construction and instrument actual oracle calls. They do not represent 2160 additional native solver runs.

C.1 Certificate validity with bounded deterministic errors

If $|\widehat{f} - f(x)| \le \nu$, $\|\widehat{g} - \nabla f(x)\| \le \rho$ and $\langle \widehat{g}, s\rangle \le \min_{v\in Q}\langle \widehat{g}, v\rangle + \eta$ with known bounds, then

$$\ell^{\text{safe}} = \widehat{f} + \langle \widehat{g}, s - x\rangle - \nu - \rho D - \eta \le f_Q^*.$$

For each $v \in Q$, convexity gives $f(v) \ge \widehat{f} - \nu + \langle \widehat{g}, v - x\rangle - \rho D \ge \ell^{\text{safe}}$; minimize over v. A maximum of such lower bounds remains valid, and $\widehat{f} + \nu - \max_j \ell_j^{\text{safe}}$ is an upper certificate. This is a validity statement, not a convergence theorem for noisy directions. Empirical variance is not a deterministic norm bound.

D Additional protocol details and controls

Features. Seoul uses eight numerical weather features (temperature, humidity, wind speed, visibility, dew point, solar radiation, rainfall and snowfall), sine/cosine of hour and weekday, holiday and functioning indicators. Metro uses temperature, rain, snow, cloud cover, sine/cosine of hour, weekday and day of year (period 365.25), and a holiday indicator. Other textual categories are omitted. Features constant on training data are dropped before the tanh transform; source encodings and feature names are recorded in the manifests. These deterministic choices precede the solver comparisons.

Native comparator. For $x \ne y$, the AC-FW estimate is

$$\ell(x,y) = \frac{2|f(y) - f(x) - \langle \nabla f(x), y - x\rangle|}{\|y - x\|^2},$$

and it is zero if $x = y$. Its update is the maximum of this trial estimate and $r_k L_k$, with $r_k = 1 - 1/((k+1)\log^2(k+3))$. Only strict objective decrease accepts a trial. After N outer trials the charged counts are $N_f = N_{\text{LMO}} = N + 2$, $N_g = N_{\text{accepted}} + 2$. Rejected trials count against the cap. All primary count-data starts are atoms, as in the cited initialization. Parameter-hidden tests on both count datasets instrument preprocessing, rejected trials and oracle calls independently of recorded counters.

Planted geometry. For the initialization controls, let

$$\phi(x) = m^{-1}\sum_i [\cosh(a_i^\top(x - x^*)) - 1] + \mu\|x - x^*\|^2/2, \quad f(x) = \phi(x) - \lambda v^\top(x - x^*),$$

where $\|x^*\|_1 = R$, $v \in \partial\|x^*\|_1$ and $\lambda = 0.15$. On Q, $f \ge \phi \ge 0 = f_Q^*$ and $\nabla f(x^*) = -\lambda v \ne 0$. Valid constants are $H_0 = \mu + \lambda_{\max}(A^\top A/m)$ and $H_1 = \max_i \|a_i\|^2$. The active and active-scaled families use $d = 12, m = 10$, $R = 1.5, \mu = 0.08$, with x^* entries $(0.6, -0.5, 0.4)$ at coordinates $(2, 6, 10)$; Gaussian row entries have standard deviation 0.22 or 0.9. The high-dimensional family has $d = 60, m = 40$ and standard deviation 0.22. Each has ten matched seeds. Off-support normal coordinates are uniform in $[-0.6, 0.6]$.

With $w = \operatorname{sign}(x^*)$, zero off support, the optimal-face slack $\sigma(x) = R - w^\top x$ obeys $\sigma(x + \alpha(s - x)) = (1 - \alpha)\sigma(x) + \alpha\sigma(s)$. Thus strict subunit steps to atoms on the face keep positive slack from an exterior start. A full initial vertex move can remove it. Moreover $f(x) \geq 0.4\lambda\sigma(x)$ on these planted examples. This familiar FW face obstruction explains why initialization is a necessary control, not a new acceleration mechanism.

Objective-unit study. The separate radius/scale archive contains 360 runs on the WDBC logistic control and UCI Bike Sharing Poisson objective [25, 26], three radii, five seeds, three positive scales and four BT methods. Accuracy is scaled with the objective. The 60 pairwise scale comparisons per certificate-initialized method compare final counters and sampled budget certificates, not bitwise identity of every intermediate iterate. All scalings of a problem are dependent controls. Additional studies and full baseline settings are retained in the expanded technical report and companion archive.

Table 5: Certification at 10^{-5} within 1000 updates. Each cell gives the certified count and median function calls to stop or cap, including failed runs. Seoul has 30 starts and Metro 24 per radius/method. F/A: FW/AFW; subscripts $1, U$: BT initialized at $1, U_0/D^2$; Q: known-H short step. AC counts 1000 outer trials, including rejections, and pays two preprocessing source evaluations. These medians are not times to success.

Data	R/R_*	F_1	F_U	A_1	A_U	F_Q	A_Q	AC
Seoul	0.5	1; 2001	1; 2003	30; 424	30; 399	0; 1001	0; 1001	0; 1002
Seoul	2	0; 2001	0; 1996	0; 2001	0; 1995.5	0; 1001	0; 1001	0; 1002
Metro	0.5	3; 2001	3; 2003	24; 99	24; 80.5	1; 1001	24; 423	13; 385
Metro	2	0; 2000	0; 2002	6; 2000	7; 2001	0; 1001	0; 1001	0; 1002

E What the closest bounds directly establish

The following comparison concerns specific stated results, not every possible consequence of their proofs. Khademi and Silveti-Falls [19, Theorems 4.4–4.5] combine gradient-difference estimates with backtracking. Their stated constrained rates retain a global smoothness/gradient contribution; they also explicitly discuss slow downward adaptation of ordinary backtracking. Thus that motivation is prior. Yagishita and Ito [20, Algorithm 3, Theorem 3.1] analyze a line-search-free composite conditional-gradient method with a nondecreasing estimate and a nonconvex minimum-gap guarantee. These are different methods and stopping statements, not evidence for a general priority claim. Pedregosa et al. [2, Theorem 3] bound a last-iterate Fenchel primal–dual gap using averaged accepted curvature estimates. Its dual lower value and the maximum scalar tangent lower bound have no general ordering for arbitrary histories: for $f = x^2/2$ on $[-1, 1]$, histories $(-1, 1)$ and $(-1, -1/2)$ reverse the order. These histories are not asserted to be algorithm trajectories and do not contradict that theorem. Its Theorem 4 already yields geometric progress from accepted local curvature, which our AFW proof specializes.

Substituting $L_Q = H_0 + H_1 \max_Q \delta$ in a global smoothness bound retains the global gap range. The smaller initial certificate U_0 does not bound curvature everywhere on Q; restricting the region to an initial sublevel set would change its LMO. Our first-return argument instead bounds only the tested descent segment. A phase count controls estimates above the current level.

The deterministic convex result in [9, Appendix C.4] already separates function-dependent-curvature regimes for gradient descent; it is not an LMO certificate theorem. The gradient-norm dependent FW analysis in [11, Theorem 5.1] contains a largest-gradient contribution after a direct substitution. Neither the use of phases nor unknown-curvature adaptation alone is presented as new here.

Takahashi et al. [13] use relative smoothness, a kernel, and segment scaling. A Euclidean kernel with a single global L_Q does not by itself give the separated dependence above. For adaptive sliding [14, Theorems 2.1 and 4.1], inner LMO effort is distinct from outer iterations, and the restarted theorem also requires local strong convexity and its scheduling parameters. Condition (1) alone does not give strong convexity: $\cosh(x_1) - 1$ on a square has a line of minimizers. No superiority over sliding is established.

Wirth et al. [15, Definition 2.3(b), Theorem 3.6] use uniform extended curvature and a vertex error bound. Kerdreux et al. [16, Proposition 4.2, Theorem 5.1] already combine burn-in, computable strong-Wolfe stopping, unknown error-bound parameters and cumulative LMO counts, with global away curvature. Zhao [17, Theorems 4.1–4.2] proves linear objective- and FW-gap rates for log-homogeneous barriers, using exact search or a local-Hessian step. Our function class and the evaluated algorithm differ. None of these comparisons rules out an equivalent refined specialization of another adaptive proof.

F Reproducibility and evidence checks

The companion package includes source snapshots, raw data attribution, CSV results and SHA-256 manifests. The main real-data comparison joins 648 preserved records with 108 additional AC-FW records on the same cases; it does not count them as new datasets. The certificate study replays 432 existing paths and reports 2160 dependent causal profiles. The initialization factorial also reuses earlier cases. Table values are generated from immutable CSVs; the claim ledger records each selection and its value.

The recorded environment uses Python 3.13.5, NumPy 2.5.2, SciPy 1.16.3, Matplotlib 3.10.9 and scikit-learn 1.8.0, with BLAS/OpenMP thread environment variables set to one. From the extracted repository root:

```
python -m pip install -r PART3/requirements.txt
python -m unittest discover -s PART3/Work -p "test_*.py"
python PART3/Work/validate_results.py
python PART3/Paper/prepare_article_evidence.py
```

The 65 tests cover derivatives, reference independence, actual oracle calls, certificates, accepted progress and curvature limits. Full trajectory audits run before trace sampling; saved validators also check provenance and budget accounting. A separate finite audit uses exact rational arithmetic for 8370 phase configurations, 57989 abstract curvature-state visits, phase budgets, gap inversion and atom-removal counts. It overapproximates possible curvature histories and does not assert that every state is realizable by an objective. Finite tests are not a formal proof or independent mathematical review.

Seven additional tests cover the exact differential envelope, its inverse and large-parameter limits, one-dimensional extremizers in 70-digit decimal arithmetic, and actual gaps on five earlier synthetic and regression objectives. A separate rational archive contains 24 AFW runs from uniform atom weights on simplex and box examples, six planted local states (12 maximal drops), and 180 quadratic trial checks for the positive-base counterexample. All 30 zero-base runs reach an exact zero objective and certificate. These are finite algebraic checks of the stated mechanisms, not evidence of a uniform empirical speedup. Both exact-audit archives replay from their own saved code.

For the simplex examples $t = c^\top x$, $c = (0, 0, 1, 2)$, and $f = t + \beta t^3$, $\beta \in \{0, 1/16, 1\}$. The Hessian norm is $6\beta t \|c\|^2 = 30\beta t \le 30\beta f$ on the simplex. For the box $0 \le t \le 1$, $|y| \le 1/2$, use $f = t + \beta t^3(1 + y^2)$. Its Hessian has nonnegative diagonal and determinant $12\beta^2 t^4(1 - 2y^2) \ge 0$; its norm is at most its trace, $6\beta t(1 + y^2) + 2\beta t^3 \le (19\beta/2)t \le 10\beta f$. Both families have a non-singleton exposed optimal face and atom margin $m = 1$. Replacing t^3 by $(\max\{t, 0\})^3$ supplies a convex C^2 extension to a neighborhood without changing any feasible value. The four

initial estimates are $1/4, 1, 4, 64$. Planted local states retain two optimal and two nonoptimal atoms and terminate by two exact maximal away steps. Two further rational paths in the tests start near that face with $M = 65536$, giving respectively seven and eight nonmaximal away updates before their last two drops. They exercise the separate initialization potential Q_0 in Theorem 4.2.

The broader legacy experiments include repeated Bike Sharing initial vertices: ten historical seeds give seven distinct starts. Later activity-sweep summaries explicitly remove those duplicates. Those historical seed-weighted summaries are not used as independent replications in the main count-data comparison, which enumerates all distinct vertices. Numerical references and floating-point audits do not provide directed-rounding guarantees.

Generative-tool disclosure. An OpenAI Codex assistant was used substantially for literature triage, proof development, implementation, experiment analysis, manuscript drafting and verification automation. The explicit derivations, source snapshots and executable checks are supplied for scrutiny. Automated checks and AI-assisted internal review are not independent mathematical peer review. Responsibility for the submitted scientific claims rests with the named authors.