
MATRIX ANALYSIS

Through Coordinate Geometry

G. V. V. Sharma



Copyright ©2022 by G. V. V. Sharma.

<https://creativecommons.org/licenses/by-sa/3.0/>

and

<https://www.gnu.org/licenses/fdl-1.3.en.html>

Contents

| | |
|--|-----------|
| Introduction | iii |
| 1 Quadrilaterals | 1 |
| 1.1 Properties | 1 |
| 2 This is Chapter Two Title | 13 |
| 2.1 This is First Level Heading | 13 |
| 2.1.1 This is Second Level Heading | 14 |
| A Vectors | 17 |
| A.1 2×1 vectors | 17 |
| A.2 3×1 vectors | 22 |
| B Matrices | 25 |
| B.1 Eigenvalues and Eigenvectors | 25 |
| B.2 Determinants | 26 |
| B.3 Rank of a Matrix | 27 |
| B.4 Inverse of a Matrix | 28 |
| B.5 Orthogonality | 28 |
| C Linear Forms | 31 |

| | |
|---|-----------|
| C.1 Two Dimensions | 31 |
| C.2 Three Dimensions | 35 |
| D Quadratic Forms | 45 |
| D.1 Conic equation | 45 |
| D.2 Standard Form | 50 |
| E Conic Parameters | 57 |
| E.1 Standard Form | 57 |
| E.2 Quadratic Form | 59 |
| F Conic Lines | 61 |
| F.1 Pair of Straight Lines | 61 |
| F.2 Intersection of Conics | 64 |
| F.3 Chords of a Conic | 65 |
| F.4 Tangent and Normal | 71 |

Introduction

This book links high school coordinate geometry to linear algebra and matrix analysis through solved problems.

Chapter 1

Quadrilaterals

1.1. Properties

1. The angles of quadrilateral are in the ratio 3:5:9:13. Find all the angles of the quadrilateral.
2. If diagonals of a parallelogram are equal then show that it is a rectangle.

Solution: See Fig. 1.1. From (A.1.18),

$$\mathbf{B} - \mathbf{A} = \mathbf{C} - \mathbf{D} \quad (1.1)$$

$$\implies \mathbf{B} - \mathbf{C} = \mathbf{A} - \mathbf{D} \quad (1.2)$$

Also, it is given that the diagonals of $ABCD$ are equal. Hence,

$$\|\mathbf{C} - \mathbf{A}\|^2 = \|\mathbf{D} - \mathbf{B}\|^2 \quad (1.3)$$

$$\implies \|(\mathbf{C} - \mathbf{B}) + (\mathbf{B} - \mathbf{A})\|^2 = \|(\mathbf{D} - \mathbf{C}) + (\mathbf{C} - \mathbf{B})\|^2 \quad (1.4)$$




Figure 1.1:

which can be expressed as

$$\begin{aligned} & \| \mathbf{C} - \mathbf{B} \|^2 + \| \mathbf{B} - \mathbf{A} \|^2 + 2(\mathbf{C} - \mathbf{B})^\top (\mathbf{B} - \mathbf{A}) \\ &= \| \mathbf{D} - \mathbf{C} \|^2 + \| \mathbf{C} - \mathbf{B} \|^2 + 2(\mathbf{D} - \mathbf{C})^\top (\mathbf{C} - \mathbf{B}) \end{aligned} \quad (1.5)$$

which, can be simplified to obtain

$$(\mathbf{C} - \mathbf{B})^\top (\mathbf{B} - \mathbf{A}) = (\mathbf{D} - \mathbf{C})^\top (\mathbf{C} - \mathbf{B}) \quad (1.6)$$

since

$$\|\mathbf{D} - \mathbf{C}\| = \|\mathbf{B} - \mathbf{A}\| \quad (1.7)$$

yielding

$$(\mathbf{A} - \mathbf{B})^\top (\mathbf{B} - \mathbf{C}) = \mathbf{0} \quad (1.8)$$

from (1.1).

3. Show that if the diagonals of a quadrilateral bisect each other at right angles, then it is a rhombus.

Solution: See Fig. 1.2. From the given information,

$$\frac{\mathbf{B} + \mathbf{D}}{2} = \frac{\mathbf{A} + \mathbf{C}}{2} \quad (1.9)$$

$$(\mathbf{B} - \mathbf{D})^\top (\mathbf{A} - \mathbf{C}) = 0 \quad (1.10)$$

From (1.9),

$$\mathbf{B} - \mathbf{A} = \mathbf{C} - \mathbf{D} \quad (1.11)$$

which, from (A.1.18), is the definition of a parallelogram. Further, substituting

$$\mathbf{B} - \mathbf{D} = (\mathbf{B} - \mathbf{A}) + (\mathbf{A} - \mathbf{D}) \quad (1.12)$$

$$\mathbf{A} - \mathbf{C} = (\mathbf{A} - \mathbf{B}) + (\mathbf{B} - \mathbf{C}) \quad (1.13)$$




Figure 1.2: Rhombus

in (1.10),

$$\begin{aligned}
 & [(\mathbf{B} - \mathbf{A}) + (\mathbf{A} - \mathbf{D})]^\top [(\mathbf{A} - \mathbf{B}) + (\mathbf{B} - \mathbf{C})] = 0 \\
 \implies & -\|\mathbf{B} - \mathbf{A}\|^2 + (\mathbf{B} - \mathbf{A})^\top (\mathbf{B} - \mathbf{C}) + \\
 & (\mathbf{A} - \mathbf{D})^\top (\mathbf{A} - \mathbf{B}) + (\mathbf{A} - \mathbf{D})^\top (\mathbf{B} - \mathbf{C}) = 0 \quad (1.14)
 \end{aligned}$$

From (1.11),

$$\mathbf{B} - \mathbf{C} = \mathbf{A} - \mathbf{D} \quad (1.15)$$

$$\implies (\mathbf{B} - \mathbf{A})^\top (\mathbf{B} - \mathbf{C}) + (\mathbf{A} - \mathbf{D})^\top (\mathbf{A} - \mathbf{B}) = \mathbf{0} \quad (1.16)$$

and

$$(\mathbf{A} - \mathbf{D})^\top (\mathbf{B} - \mathbf{C}) = \|\mathbf{B} - \mathbf{C}\|^2 \quad (1.17)$$

Substituting from

(1.16) and (1.17) in (1.14),

$$\|\mathbf{A} - \mathbf{B}\|^2 = \|\mathbf{B} - \mathbf{C}\|^2 \quad (1.18)$$

which means that the adjacent sides of the parallelogram are equal. Thus, the quadrilateral is a rhombus

4. Show that the diagonals of a square are equal and bisect each other at right angles.

Solution: This is obvious from Problems (2) and (3).

5.

6. Diagonal AC of a parallelogram ABCD bisects $\angle A$ in Fig (1.3). Show that

- (a) it bisects $\angle C$ also
- (b) $ABCD$ is a rhombus

Solution:




Figure 1.3:

(a) From (A.23),

$$\angle BAC = \angle DAC \quad (1.19)$$

$$\Rightarrow \frac{(\mathbf{A} - \mathbf{B})^T (\mathbf{A} - \mathbf{C})}{\|\mathbf{A} - \mathbf{B}\| \|\mathbf{A} - \mathbf{C}\|} = \frac{(\mathbf{A} - \mathbf{D})^T (\mathbf{A} - \mathbf{C})}{\|\mathbf{A} - \mathbf{D}\| \|\mathbf{A} - \mathbf{C}\|} \quad (1.20)$$

Also,

$$\cos \angle ACD = \frac{(\mathbf{C} - \mathbf{D})^T (\mathbf{C} - \mathbf{A})}{\|\mathbf{C} - \mathbf{D}\| \|\mathbf{C} - \mathbf{A}\|} \quad (1.21)$$

From Appendix A.1.18,

$$\mathbf{B} - \mathbf{A} = \mathbf{C} - \mathbf{D} \quad (1.22)$$

$$\Rightarrow \frac{(\mathbf{C} - \mathbf{D})^T(\mathbf{C} - \mathbf{A})}{\|\mathbf{C} - \mathbf{D}\| \|\mathbf{C} - \mathbf{A}\|} = \frac{(\mathbf{B} - \mathbf{A})^T(\mathbf{C} - \mathbf{A})}{\|\mathbf{B} - \mathbf{A}\| \|\mathbf{C} - \mathbf{A}\|} \quad (1.23)$$

upon substituting in (1.21). Thus, from (1.21) and (1.19),

$$\angle BAC = \angle DAC = \angle ACD \quad (1.24)$$

Similarly, it can be shown that

$$\angle ACD = \angle ACB \quad (1.25)$$

(b)

7. $ABCD$ is a rhombus. Show that the diagonal AC bisects angle A as well as angle C and diagonal BD bisects angle B as well as angle D .

Solution: For the rhombus in Fig. 1.4,

$$\|\mathbf{A} - \mathbf{B}\| = \|\mathbf{A} - \mathbf{D}\| \quad (1.26)$$

$$\mathbf{A} - \mathbf{B} = \mathbf{D} - \mathbf{C}$$

From (A.23),

$$\begin{aligned} \cos \angle BAC &= \frac{(\mathbf{A} - \mathbf{B})^T(\mathbf{A} - \mathbf{C})}{\|\mathbf{A} - \mathbf{B}\| \|\mathbf{A} - \mathbf{C}\|} \\ \cos \angle DAC &= \frac{(\mathbf{C} - \mathbf{D})^T(\mathbf{C} - \mathbf{A})}{\|\mathbf{C} - \mathbf{D}\| \|\mathbf{C} - \mathbf{A}\|} \end{aligned} \quad (1.27)$$




Figure 1.4:

From (1.26) and (1.27), we obtain

$$\cos \angle BAC = \cos \angle DAC \quad (1.28)$$

Thus, AC bisects $\angle A$. Similarly, the remaining results can be proved.

8.

9. In parallelogram $ABCD$, two points \mathbf{P} and \mathbf{Q} are taken on diagonal BD such that $DP = BQ$. Show that

$$(a) \quad \triangle APD \cong \triangle CQB$$

$$(b) \quad AP = CQ$$

$$(c) \quad \triangle AQB \cong \triangle CPD$$

$$(d) \quad AQ = CP$$

$$(e) \quad APCQ \text{ is a parallelogram}$$

Solution: See Fig. 1.5.

From (A.23) and the given information,

$$\mathbf{A} - \mathbf{B} = \mathbf{D} - \mathbf{C} \quad (1.29)$$

$$\implies \mathbf{A} - \mathbf{D} = \mathbf{B} - \mathbf{C} \quad (1.30)$$

$$\mathbf{B} - \mathbf{Q} = \mathbf{P} - \mathbf{D} \quad (1.31)$$




Figure 1.5:

From (1) and (3):

$$\begin{aligned}\mathbf{A} + \mathbf{C} &= \mathbf{B} + \mathbf{D} \\ \mathbf{P} + \mathbf{Q} &= \mathbf{B} + \mathbf{D}\end{aligned}\tag{1.32}$$

From (4)

$$\mathbf{A} + \mathbf{C} = \mathbf{P} + \mathbf{Q}\tag{1.33}$$

From (5)

$$\implies \mathbf{A} - \mathbf{Q} = \mathbf{P} - \mathbf{C}\tag{1.34}$$

$$\mathbf{A} - \mathbf{P} = \mathbf{Q} - \mathbf{C}\tag{1.35}$$

(a) From (2),(3) and (7)

$$\triangle APD \cong \triangle CQB\tag{1.36}$$

(b)

$$\text{Equation(7)} \implies AP = CQ\tag{1.37}$$

(c) From (1),(3) and (6)

$$\triangle AQB \cong \triangle CPD\tag{1.38}$$

(d)

$$\text{Equation(6)} \implies AQ = CP \quad (1.39)$$

(e) Equation (6) and (7) \implies Quadrilateral APCQ is a parallelogram.

Chapter 2

This is Chapter Two Title

2.1. This is First Level Heading

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus

luctus mauris.

2.1.1. This is Second Level Heading

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

2.1.1.1. This is Third Level Heading

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem ipsum dolor sit amet, consectetur adipiscing elit. In hac habitasse platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.

This is Fourth Level Heading. Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla,

wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur.

This is Fifth Level Heading. Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio.

Appendix A

Vectors

A.1. 2×1 vectors

A.1.1. Let

$$\mathbf{A} \equiv \vec{A} = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} \tag{A.1}$$

$$\equiv a_1 \vec{i} + a_2 \vec{j}, \tag{A.2}$$

$$\mathbf{B} = \begin{pmatrix} b_1 \\ b_2 \end{pmatrix}, \tag{A.3}$$

be 2×1 vectors. Then, the determinant of the 2×2 matrix

$$\mathbf{M} = \begin{pmatrix} \mathbf{A} & \mathbf{B} \end{pmatrix} \tag{A.4}$$

is defined as

$$\left| \mathbf{M} \right| = \left| \begin{array}{cc} \mathbf{A} & \mathbf{B} \end{array} \right| \quad (\text{A.5})$$

$$= \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix} = a_1 b_2 - a_2 b_1 \quad (\text{A.6})$$

A.1.2. The value of the cross product of two vectors is given by (A.5).

A.1.3. The area of the triangle with vertices $\mathbf{A}, \mathbf{B}, \mathbf{C}$ is given by the absolute value of

$$\frac{1}{2} \left| \begin{array}{cc} \mathbf{A} - \mathbf{B} & \mathbf{A} - \mathbf{C} \end{array} \right| \quad (\text{A.7})$$

A.1.4. The transpose of \mathbf{A} is defined as

$$\mathbf{A}^\top = \begin{pmatrix} a_1 & a_2 \end{pmatrix} \quad (\text{A.8})$$

A.1.5. The inner product or dot product is defined as

$$\mathbf{A}^\top \mathbf{B} \equiv \mathbf{A} \cdot \mathbf{B} \quad (\text{A.9})$$

$$= \begin{pmatrix} a_1 & a_2 \end{pmatrix} \begin{pmatrix} b_1 \\ b_2 \end{pmatrix} = a_1 b_1 + a_2 b_2 \quad (\text{A.10})$$

A.1.6. norm of \mathbf{A} is defined as

$$\|A\| \equiv \left| \vec{A} \right| \quad (\text{A.11})$$

$$= \sqrt{\mathbf{A}^\top \mathbf{A}} = \sqrt{a_1^2 + a_2^2} \quad (\text{A.12})$$

Thus,

$$\|\lambda \mathbf{A}\| \equiv \left| \lambda \vec{A} \right| \quad (\text{A.13})$$

$$= |\lambda| \|\mathbf{A}\| \quad (\text{A.14})$$

A.1.7. The distance between the points \mathbf{A} and \mathbf{B} is given by

$$\|\mathbf{A} - \mathbf{B}\| \quad (\text{A.15})$$

A.1.8. Let \mathbf{x} be equidistant from the points \mathbf{A} and \mathbf{B} . Then

$$(\mathbf{A} - \mathbf{B})^\top \mathbf{x} = \frac{\|\mathbf{A}\|^2 - \|\mathbf{B}\|^2}{2} \quad (\text{A.16})$$

Solution:

$$\|\mathbf{x} - \mathbf{A}\| = \|\mathbf{A} - \mathbf{B}\| \quad (\text{A.17})$$

$$\implies \|\mathbf{x} - \mathbf{A}\|^2 = \|\mathbf{x} - \mathbf{B}\|^2 \quad (\text{A.18})$$

which can be expressed as

$$\begin{aligned} (\mathbf{x} - \mathbf{A})^\top (\mathbf{x} - \mathbf{A}) &= (\mathbf{x} - \mathbf{B})^\top (\mathbf{x} - \mathbf{B}) \\ \implies \|\mathbf{x}\|^2 - 2\mathbf{x}^\top \mathbf{A} + \|\mathbf{A}\|^2 &= \|\mathbf{x}\|^2 - 2\mathbf{x}^\top \mathbf{B} + \|\mathbf{B}\|^2 \quad (\text{A.19}) \end{aligned}$$

which can be simplified to obtain (A.16).

A.1.9. If \mathbf{x} lies on the x -axis and is equidistant from the points \mathbf{A} and \mathbf{B} ,

$$\mathbf{x} = x\mathbf{e}_1 \quad (\text{A.20})$$

where

$$x = \frac{\|\mathbf{A}\|^2 - \|\mathbf{B}\|^2}{2(\mathbf{A} - \mathbf{B})^\top \mathbf{e}_1} \quad (\text{A.21})$$

Solution: From (A.16).

$$x(\mathbf{A} - \mathbf{B})^\top \mathbf{e}_1 = \frac{\|\mathbf{A}\|^2 - \|\mathbf{B}\|^2}{2} \quad (\text{A.22})$$

yielding (A.21).

A.1.10. The angle between two vectors is given by

$$\theta = \cos^{-1} \frac{\mathbf{A}^\top \mathbf{B}}{\|\mathbf{A}\| \|\mathbf{B}\|} \quad (\text{A.23})$$

A.1.11. If two vectors are orthogonal (perpendicular),

$$\mathbf{A}^\top \mathbf{B} = 0 \quad (\text{A.24})$$

A.1.12. The direction vector of the line joining two points \mathbf{A}, \mathbf{B} is given by

$$\mathbf{m} = \mathbf{A} - \mathbf{B} \quad (\text{A.25})$$

A.1.13. The unit vector in the direction of \mathbf{m} is defined as

$$\frac{\mathbf{m}}{\|\mathbf{m}\|} \quad (\text{A.26})$$

A.1.14. If the direction vector of a line is expressed as

$$\mathbf{m} = \begin{pmatrix} 1 \\ m \end{pmatrix}, \quad (\text{A.27})$$

the m is defined to be the slope of the line.

A.1.15. The normal vector to \mathbf{m} is defined by

$$\mathbf{m}^\top \mathbf{n} = 0 \quad (\text{A.28})$$

A.1.16. The point \mathbf{P} that divides the line segment AB in the ratio $k : 1$ is given by

$$\mathbf{P} = \frac{k\mathbf{B} + \mathbf{A}}{k + 1} \quad (\text{A.29})$$

A.1.17. The standard basis vectors are defined as

$$\mathbf{e}_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad (\text{A.30})$$

$$\mathbf{e}_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \quad (\text{A.31})$$

A.1.18. If $ABCD$ be a parallelogram,

$$\mathbf{B} - \mathbf{A} = \mathbf{C} - \mathbf{D} \quad (\text{A.32})$$

A.2. 3×1 vectors

A.2.1. Let

$$\mathbf{A} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \equiv a_1 \vec{i} + a_2 \vec{j} + a_3 \vec{j}, \quad (\text{A.33})$$

$$\mathbf{B} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}, \quad (\text{A.34})$$

and

$$\mathbf{A}_{ij} = \begin{pmatrix} a_i \\ a_j \end{pmatrix}, \quad (\text{A.35})$$

$$\mathbf{B}_{ij} = \begin{pmatrix} b_i \\ b_j \end{pmatrix}. \quad (\text{A.36})$$

A.2.2. The cross product or vector product of \mathbf{A}, \mathbf{B} is defined as

$$\mathbf{A} \times \mathbf{B} = \begin{pmatrix} \begin{vmatrix} \mathbf{A}_{23} & \mathbf{B}_{23} \\ \mathbf{A}_{31} & \mathbf{B}_{31} \\ \mathbf{A}_{12} & \mathbf{B}_{12} \end{vmatrix} \end{pmatrix} \quad (\text{A.37})$$

A.2.3. Verify that

$$\mathbf{A} \times \mathbf{B} = -\mathbf{B} \times \mathbf{A} \quad (\text{A.38})$$

A.2.4. The area of a triangle is given by

$$\frac{1}{2} \|\mathbf{A} \times \mathbf{B}\| \quad (\text{A.39})$$

Appendix B

Matrices

B.1. Eigenvalues and Eigenvectors

B.1.1. The eigenvalue λ and the eigenvector \mathbf{x} for a matrix \mathbf{A} are defined as,

$$\mathbf{A}\mathbf{x} = \lambda\mathbf{x} \tag{B.1}$$

B.1.2. The eigenvalues are calculated by solving the equation

$$f(\lambda) = \left| \lambda\mathbf{I} - \mathbf{A} \right| = 0 \tag{B.2}$$

The above equation is known as the characteristic equation.

B.1.3. According to the Cayley-Hamilton theorem,

$$f(\lambda) = 0 \implies f(\mathbf{A}) = 0 \tag{B.3}$$

B.1.4. The trace of a square matrix is defined to be the sum of the diagonal elements.

$$\text{tr}(\mathbf{A}) = \sum_{i=1}^N a_{ii}. \tag{B.4}$$

where a_{ii} is the i th diagonal element of the matrix \mathbf{A} .

B.1.5. The trace of a matrix is equal to the sum of the eigenvalues

$$\text{tr}(\mathbf{A}) = \sum_{i=1}^N \lambda_i \quad (\text{B.5})$$

B.2. Determinants

B.2.1. Let

$$\mathbf{A} = \begin{pmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{pmatrix}. \quad (\text{B.6})$$

be a 3×3 matrix. Then,

$$\begin{aligned} |\mathbf{A}| &= a_1 \begin{pmatrix} b_2 & c_2 \\ b_3 & c_3 \end{pmatrix} - a_2 \begin{pmatrix} b_1 & c_1 \\ b_3 & c_3 \end{pmatrix} \\ &\quad + a_3 \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix}. \end{aligned} \quad (\text{B.7})$$

B.2.2. Let $\lambda_1, \lambda_2, \dots, \lambda_n$ be the eigenvalues of a matrix \mathbf{A} . Then, the product of the eigenvalues is equal to the determinant of \mathbf{A} .

$$|\mathbf{A}| = \prod_{i=1}^n \lambda_i \quad (\text{B.8})$$

B.2.3.

$$\left| \mathbf{AB} \right| = \left| \mathbf{A} \right| \left| \mathbf{B} \right| \quad (\text{B.9})$$

B.2.4. If \mathbf{A} be an $n \times n$ matrix,

$$\left| k\mathbf{A} \right| = k^n \left| \mathbf{A} \right| \quad (\text{B.10})$$

B.3. Rank of a Matrix

B.3.1. The rank of a matrix is defined as the number of linearly independent rows. This is also known as the row rank.

B.3.2. Row rank = Column rank.

B.3.3. The rank of a matrix is obtained as the number of nonzero rows obtained after row reduction.

B.3.4. An $n \times n$ matrix is invertible if and only if its rank is n .

B.3.5. Points $\mathbf{A}, \mathbf{B}, \mathbf{C}$ are on a line if

$$\text{rank} \begin{pmatrix} \mathbf{A} \\ \mathbf{B} \\ \mathbf{C} \end{pmatrix} = 1 \quad (\text{B.11})$$

B.3.6. Points $\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}$ form a paralelogram if

$$\text{rank} \begin{pmatrix} \mathbf{A} \\ \mathbf{B} \\ \mathbf{C} \\ \mathbf{D} \end{pmatrix} = 1, \text{rank} \begin{pmatrix} \mathbf{A} \\ \mathbf{B} \\ \mathbf{C} \end{pmatrix} = 2 \quad (\text{B.12})$$

B.4. Inverse of a Matrix

B.4.1. For a 2×2 matrix

$$\mathbf{A} = \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix}, \quad (\text{B.13})$$

the inverse is given by

$$\mathbf{A}^{-1} = \frac{1}{|\mathbf{A}|} \begin{pmatrix} b_2 & -b_1 \\ -a_2 & a_1 \end{pmatrix}, \quad (\text{B.14})$$

B.4.2. For higher order matrices, the inverse should be calculated using row operations.

B.5. Orthogonality

B.5.1. The rotation matrix is defined as

$$\mathbf{R}_\theta = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}, \quad \theta \in [0, 2\pi] \quad (\text{B.15})$$

B.5.2. The rotation matrix is orthogonal

$$\mathbf{R}_\theta^\top \mathbf{R}_\theta = \mathbf{R}_\theta \mathbf{R}_\theta^\top = \mathbf{I} \quad (\text{B.16})$$

B.5.3.

$$\mathbf{m}^\top \mathbf{n} = 0 \implies \mathbf{n} = \mathbf{R}_{\frac{\pi}{2}} \mathbf{m} \quad (\text{B.17})$$

B.5.4.

$$\mathbf{n}^\top \mathbf{h} = 1 \implies \mathbf{n} = \frac{\mathbf{e}_1}{\mathbf{e}_1^\top \mathbf{h}} + \mu \mathbf{R}_{\frac{\pi}{2}} \mathbf{h}, \quad \mu \in \mathbb{R}. \quad (\text{B.18})$$

B.5.5. The affine transformation is given by

$$\mathbf{x} = \mathbf{P}\mathbf{y} + \mathbf{c} \quad (\text{Affine Transformation}) \quad (\text{B.19})$$

where \mathbf{P} is invertible.

B.5.6. The eigenvalue decomposition of a symmetric matrix \mathbf{V} is given by

$$\mathbf{P}^\top \mathbf{V} \mathbf{P} = \mathbf{D}. \quad (\text{Eigenvalue Decomposition}) \quad (\text{B.20})$$

$$\mathbf{D} = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}, \quad (\text{B.21})$$

$$\mathbf{P} = \begin{pmatrix} \mathbf{p}_1 & \mathbf{p}_2 \end{pmatrix}, \quad \mathbf{P}^\top = \mathbf{P}^{-1}, \quad (\text{B.22})$$

Appendix C

Linear Forms

C.1. Two Dimensions

C.1.1. The equation of a line is given by

$$\mathbf{n}^\top \mathbf{x} = c \tag{C.1}$$

where \mathbf{n} is the normal vector of the line.

C.1.2. The equation of a line with normal vector \mathbf{n} and passing through a point \mathbf{A} is given by

$$\mathbf{n}^\top (\mathbf{x} - \mathbf{A}) = 0 \tag{C.2}$$

C.1.3. The equation of a line L is also given by

$$\mathbf{n}^\top \mathbf{x} = \begin{cases} 0 & \mathbf{0} \in L \\ 1 & \text{otherwise} \end{cases} \tag{C.3}$$

C.1.4. The parametric equation of a line is given by

$$\mathbf{x} = \mathbf{A} + \lambda \mathbf{m} \quad (\text{C.4})$$

where \mathbf{m} is the direction vector of the line and \mathbf{A} is any point on the line.

C.1.5. Let \mathbf{A} and \mathbf{B} be two points on a straight line and let $\mathbf{P} = \begin{pmatrix} p_1 \\ p_2 \end{pmatrix}$ be any point on it.

If p_2 is known, then

$$\mathbf{P} = \mathbf{A} + \frac{p_2 - \mathbf{e}_2^\top \mathbf{A}}{\mathbf{e}_2^\top (\mathbf{B} - \mathbf{A})} (\mathbf{B} - \mathbf{A}) \quad (\text{C.5})$$

Solution: The equation of the line can be expressed in parametric form as

$$\mathbf{x} = \mathbf{A} + \lambda (\mathbf{B} - \mathbf{A}) \quad (\text{C.6})$$

$$\implies \mathbf{P} = \mathbf{A} + \lambda (\mathbf{B} - \mathbf{A}) \quad (\text{C.7})$$

$$\implies \mathbf{e}_2^\top \mathbf{P} = \mathbf{e}_2^\top \mathbf{A} + \lambda \mathbf{e}_2^\top (\mathbf{B} - \mathbf{A}) \quad (\text{C.8})$$

$$\implies p_2 = \mathbf{e}_2^\top \mathbf{A} + \lambda \mathbf{e}_2^\top (\mathbf{B} - \mathbf{A}) \quad (\text{C.9})$$

$$\text{or, } \lambda = \frac{p_2 - \mathbf{e}_2^\top \mathbf{A}}{\mathbf{e}_2^\top (\mathbf{B} - \mathbf{A})} \quad (\text{C.10})$$

yielding (C.5).

C.1.6. The distance from a point \mathbf{P} to the line in (C.1) is given by

$$d = \frac{|\mathbf{n}^\top \mathbf{P} - c|}{\|\mathbf{n}\|} \quad (\text{C.11})$$

Solution: Without loss of generality, let \mathbf{A} be the foot of the perpendicular from \mathbf{P}

to the line in (C.4). The equation of the normal to (C.1) can then be expressed as

$$\mathbf{x} = \mathbf{A} + \lambda \mathbf{n} \quad (\text{C.12})$$

$$\implies \mathbf{P} - \mathbf{A} = \lambda \mathbf{n} \quad (\text{C.13})$$

$\because \mathbf{P}$ lies on (C.12). From the above, the desired distance can be expressed as

$$d = \|\mathbf{P} - \mathbf{A}\| = |\lambda| \|\mathbf{n}\| \quad (\text{C.14})$$

From (C.13),

$$\mathbf{n}^\top (\mathbf{P} - \mathbf{A}) = \lambda \mathbf{n}^\top \mathbf{n} = \lambda \|\mathbf{n}\|^2 \quad (\text{C.15})$$

$$\implies |\lambda| = \frac{|\mathbf{n}^\top (\mathbf{P} - \mathbf{A})|}{\|\mathbf{n}\|^2} \quad (\text{C.16})$$

Substituting the above in (C.14) and using the fact that

$$\mathbf{n}^\top \mathbf{A} = c \quad (\text{C.17})$$

from (C.1), yields (C.11)

C.1.7. The distance from the origin to the line in (C.1) is given by

$$d = \frac{|c|}{\|\mathbf{n}\|} \quad (\text{C.18})$$

C.1.8. The distance between the parallel lines

$$\begin{aligned}\mathbf{n}^\top \mathbf{x} &= c_1 \\ \mathbf{n}^\top \mathbf{x} &= c_2\end{aligned}\tag{C.19}$$

is given by

$$d = \frac{|c_1 - c_2|}{\|\mathbf{n}\|}\tag{C.20}$$

C.1.9. The equation of the line perpendicular to (C.1) and passing through the point \mathbf{P} is given by

$$\mathbf{m}^\top (\mathbf{x} - \mathbf{P}) = 0\tag{C.21}$$

C.1.10. The foot of the perpendicular from \mathbf{P} to the line in (C.1) is given by

$$\begin{pmatrix} \mathbf{m} & \mathbf{n} \end{pmatrix}^\top \mathbf{x} = \begin{pmatrix} \mathbf{m}^\top \mathbf{P} \\ c \end{pmatrix}\tag{C.22}$$

Solution: From (C.1) and (C.2) the foot of the perpendicular satisfies the equations

$$\mathbf{n}^\top \mathbf{x} = c\tag{C.23}$$

$$\mathbf{m}^\top (\mathbf{x} - \mathbf{P}) = 0\tag{C.24}$$

where \mathbf{m} is the direction vector of the given line. Combining the above into a matrix equation results in (C.22).

C.1.11. The equations of the angle bisectors of the lines

$$\mathbf{n}_1^\top \mathbf{x} = c_1 \quad (\text{C.25})$$

$$\mathbf{n}_2^\top \mathbf{x} = c_2 \quad (\text{C.26})$$

are given by

$$\frac{\mathbf{n}_1^\top \mathbf{x} - c_1}{\|\mathbf{n}_1\|} = \pm \frac{\mathbf{n}_2^\top \mathbf{x} - c_2}{\|\mathbf{n}_2\|} \quad (\text{C.27})$$

Proof. Any point on the angle bisector is equidistant from the lines. □

C.2. Three Dimensions

C.2.1. Points $\mathbf{A}, \mathbf{B}, \mathbf{C}$ are on a line if

$$\text{rank} \begin{pmatrix} \mathbf{A} \\ \mathbf{B} \\ \mathbf{C} \end{pmatrix} = 1 \quad (\text{C.28})$$

C.2.2. Points $\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}$ form a parallelogram if

$$\text{rank} \begin{pmatrix} \mathbf{A} \\ \mathbf{B} \\ \mathbf{C} \\ \mathbf{D} \end{pmatrix} = 1, \text{rank} \begin{pmatrix} \mathbf{A} \\ \mathbf{B} \\ \mathbf{C} \end{pmatrix} = 2 \quad (\text{C.29})$$

C.2.3. The equation of a line is given by (C.4)

C.2.4. The equation of a plane is given by (C.1)

C.2.5. The distance from the origin to the line in (C.1) is given by (C.18)

C.2.6. The distance from a point \mathbf{P} to the line in (C.4) is given by

$$d = \|\mathbf{A} - \mathbf{P}\|^2 - \frac{\{\mathbf{m}^\top (\mathbf{A} - \mathbf{P})\}^2}{\|\mathbf{m}\|^2} \quad (\text{C.30})$$

Solution:

$$d(\lambda) = \|\mathbf{A} + \lambda\mathbf{m} - \mathbf{P}\| \quad (\text{C.31})$$

$$\implies d^2(\lambda) = \|\mathbf{A} + \lambda\mathbf{m} - \mathbf{P}\|^2 \quad (\text{C.32})$$

which can be simplified to obtain

$$d^2(\lambda) = \lambda^2 \|\mathbf{m}\|^2 + 2\lambda\mathbf{m}^\top (\mathbf{A} - \mathbf{P}) + \|\mathbf{A} - \mathbf{P}\|^2 \quad (\text{C.33})$$

which is of the form

$$d^2(\lambda) = a\lambda^2 + 2b\lambda + c \quad (\text{C.34})$$

$$= a \left\{ \left(\lambda + \frac{b}{a} \right)^2 + \left[\frac{c}{a} - \left(\frac{b}{a} \right)^2 \right] \right\} \quad (\text{C.35})$$

with

$$a = \|\mathbf{m}\|^2, b = \mathbf{m}^\top (\mathbf{A} - \mathbf{P}), c = \|\mathbf{A} - \mathbf{P}\|^2 \quad (\text{C.36})$$

which can be expressed as From the above, $d^2(\lambda)$ is smallest when upon substituting

from (C.36)

$$\lambda + \frac{b}{2a} = 0 \implies \lambda = -\frac{b}{2a} \quad (\text{C.37})$$

$$= -\frac{\mathbf{m}^\top (\mathbf{A} - \mathbf{P})}{\|\mathbf{m}\|^2} \quad (\text{C.38})$$

and consequently,

$$d_{\min}(\lambda) = a \left(\frac{c}{a} - \left(\frac{b}{a} \right)^2 \right) \quad (\text{C.39})$$

$$= c - \frac{b^2}{a} \quad (\text{C.40})$$

yielding (C.30) after substituting from (C.36).

C.2.7. The distance between the parallel planes (C.19) is given by (C.20).

C.2.8. The plane

$$\mathbf{n}^\top \mathbf{x} = c \quad (\text{C.41})$$

contains the line

$$\mathbf{x} = \mathbf{A} + \lambda \mathbf{m} \quad (\text{C.42})$$

if

$$\mathbf{m}^\top \mathbf{n} = 0 \quad (\text{C.43})$$

Solution: Any point on the line (C.42) should also satisfy (C.41). Hence,

$$\mathbf{n}^\top (\mathbf{A} + \lambda \mathbf{m}) = \mathbf{n}^\top \mathbf{A} = c \quad (\text{C.44})$$

which can be simplified to obtain (C.43)

C.2.9. The foot of the perpendicular from a point \mathbf{P} to the plane

$$\mathbf{n}^\top \mathbf{x} = c \quad (\text{C.45})$$

is given by

Solution: The equation of the line perpendicular to the given plane and passing through \mathbf{P} is

$$\mathbf{x} = \mathbf{P} + \lambda \mathbf{n} \quad (\text{C.46})$$

From (C.58), the intersection of the above line with the given plane is

$$\mathbf{x} = \mathbf{P} + \frac{c - \mathbf{n}^\top \mathbf{P}}{\|\mathbf{n}\|^2} \mathbf{n} \quad (\text{C.47})$$

C.2.10. The image of a point \mathbf{P} with respect to the plane

$$\mathbf{n}^\top \mathbf{x} = c \quad (\text{C.48})$$

is given by

$$\mathbf{R} = \mathbf{P} + 2 \frac{c - \mathbf{n}^\top \mathbf{P}}{\|\mathbf{n}\|^2} \mathbf{n} \quad (\text{C.49})$$

Solution: Let \mathbf{R} be the desired image. Then, substituting the expression for the foot of the perpendicular from \mathbf{P} to the given plane using (C.47),

$$\frac{\mathbf{P} + \mathbf{R}}{2} = \mathbf{P} + \frac{c - \mathbf{n}^\top \mathbf{P}}{\|\mathbf{n}\|^2} \quad (\text{C.50})$$

C.2.11. Let a plane pass through the points \mathbf{A}, \mathbf{B} and be perpendicular to the plane

$$\mathbf{n}^\top \mathbf{x} = c \quad (\text{C.51})$$

Then the equation of this plane is given by

$$\mathbf{p}^\top \mathbf{x} = 1 \quad (\text{C.52})$$

where

$$\mathbf{p} = \begin{pmatrix} \mathbf{A} & \mathbf{B} & \mathbf{n} \end{pmatrix}^{-\top} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \quad (\text{C.53})$$

Solution: From the given information,

$$\mathbf{p}^\top \mathbf{A} = d \quad (\text{C.54})$$

$$\mathbf{p}^\top \mathbf{B} = d \quad (\text{C.55})$$

$$\mathbf{p}^\top \mathbf{n} = 0 \quad (\text{C.56})$$

\therefore the normal vectors to the two planes will also be perpendicular. The system of

equations in (C.56) can be expressed as the matrix equation

$$\begin{pmatrix} \mathbf{A} & \mathbf{B} & \mathbf{n} \end{pmatrix}^\top \mathbf{p} = d \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \quad (\text{C.57})$$

which yields (C.53) upon normalising with d .

C.2.12. The intersection of the line represented by (C.4) with the plane represented by (C.1) is given by

$$\mathbf{x} = \mathbf{A} + \frac{c - \mathbf{n}^\top \mathbf{A}}{\mathbf{n}^\top \mathbf{m}} \mathbf{m} \quad (\text{C.58})$$

Solution: From (C.4) and (C.1),

$$\mathbf{x} = \mathbf{A} + \lambda \mathbf{m} \quad (\text{C.59})$$

$$\mathbf{n}^\top \mathbf{x} = c \quad (\text{C.60})$$

$$\implies \mathbf{n}^\top (\mathbf{A} + \lambda \mathbf{m}) = c \quad (\text{C.61})$$

which can be simplified to obtain

$$\mathbf{n}^\top \mathbf{A} + \lambda \mathbf{n}^\top \mathbf{m} = c \quad (\text{C.62})$$

$$\implies \lambda = \frac{c - \mathbf{n}^\top \mathbf{A}}{\mathbf{n}^\top \mathbf{m}} \quad (\text{C.63})$$

Substituting the above in (C.61) yields (C.58).

C.2.13. The foot of the perpendicular from the point \mathbf{P} to the line represented by (C.4) is

given by

$$\mathbf{x} = \mathbf{A} + \frac{\mathbf{m}^\top (\mathbf{P} - \mathbf{A})}{\|\mathbf{m}\|^2} \mathbf{m} \quad (\text{C.64})$$

Solution: Let the equation of the line be

$$\mathbf{x} = \mathbf{A} + \lambda \mathbf{m} \quad (\text{C.65})$$

The equation of the plane perpendicular to the given line passing through \mathbf{P} is given by

$$\mathbf{m}^\top (\mathbf{x} - \mathbf{P}) = 0 \quad (\text{C.66})$$

$$\implies \mathbf{m}^\top \mathbf{x} = \mathbf{m}^\top \mathbf{P} \quad (\text{C.67})$$

The desired foot of the perpendicular is the intersection of (C.65) with (C.66) which can be obtained from (C.58) as (C.64)

C.2.14. The foot of the perpendicular from a point \mathbf{P} to a plane is \mathbf{Q} . The equation of the plane is given by

$$(\mathbf{P} - \mathbf{Q})^\top (\mathbf{x} - \mathbf{Q}) = 0 \quad (\text{C.68})$$

Solution: The normal vector to the plane is given by

$$\mathbf{n} = \mathbf{P} - \mathbf{Q} \quad (\text{C.69})$$

Hence, the equation of the plane is (C.68).

C.2.15. Let $\mathbf{A}, \mathbf{B}, \mathbf{C}$ be points on a plane. The equation of the plane is then given by

$$\begin{pmatrix} \mathbf{A} & \mathbf{B} & \mathbf{C} \end{pmatrix}^\top \mathbf{n} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \quad (\text{C.70})$$

Solution: Let the equation of the plane be

$$\mathbf{n}^\top \mathbf{x} = 1 \quad (\text{C.71})$$

Then

$$\mathbf{n}^\top \mathbf{A} = 1 \quad (\text{C.72})$$

$$\mathbf{n}^\top \mathbf{B} = 1 \quad (\text{C.73})$$

$$\mathbf{n}^\top \mathbf{C} = 1 \quad (\text{C.74})$$

which can be combined to obtain (C.70).

C.2.16. (Parallelogram Law) Let $\mathbf{A}, \mathbf{B}, \mathbf{D}$ be three vertices of a parallelogram. Then the vertex \mathbf{C} is given by

$$\mathbf{C} = \mathbf{B} + \mathbf{D} - \mathbf{A} \quad (\text{C.75})$$

Solution: Shifting \mathbf{A} to the origin, we obtain a parallelogram with corresponding vertices

$$\mathbf{0}, \mathbf{B} - \mathbf{A}, \mathbf{D} - \mathbf{A} \quad (\text{C.76})$$

The fourth vertex of this parallelogram is then obtained as

$$(\mathbf{B} - \mathbf{A}) + (\mathbf{D} - \mathbf{A}) = \mathbf{D} + \mathbf{B} - 2\mathbf{A} \quad (\text{C.77})$$

Shifting the origin to \mathbf{A} , the fourth vertex is obtained as

$$\mathbf{C} = \mathbf{D} + \mathbf{B} - 2\mathbf{A} + \mathbf{A} \quad (\text{C.78})$$

$$= \mathbf{D} + \mathbf{B} - \mathbf{A} \quad (\text{C.79})$$

C.2.17. (Affine Transformation) Let \mathbf{A}, \mathbf{C} , be opposite vertices of a square. The other two points can be obtained as

$$\mathbf{B} = \frac{\|\mathbf{A} - \mathbf{C}\|}{\sqrt{2}} \mathbf{P} \mathbf{e}_1 + \mathbf{A} \quad (\text{C.80})$$

$$\mathbf{D} = \frac{\|\mathbf{A} - \mathbf{C}\|}{\sqrt{2}} \mathbf{P} \mathbf{e}_2 + \mathbf{A} \quad (\text{C.81})$$

where

$$\mathbf{P} = \begin{pmatrix} \cos\left(\theta - \frac{\pi}{4}\right) & \sin\left(\theta - \frac{\pi}{4}\right) \\ \sin\left(\theta - \frac{\pi}{4}\right) & \cos\left(\theta - \frac{\pi}{4}\right) \end{pmatrix} \quad (\text{C.82})$$

and

$$\cos \theta = \frac{(\mathbf{C} - \mathbf{A})^\top \mathbf{e}_1}{\|\mathbf{A} - \mathbf{C}\| \|\mathbf{e}_1\|} \quad (\text{C.83})$$

Appendix D

Quadratic Forms

D.1. Conic equation

D.1.1. Let \mathbf{q} be a point such that the ratio of its distance from a fixed point \mathbf{F} and the distance (d) from a fixed line

$$L : \mathbf{n}^\top \mathbf{x} = c \tag{D.1}$$

is constant, given by

$$\frac{\|\mathbf{q} - \mathbf{F}\|}{d} = e \tag{D.2}$$

The locus of \mathbf{q} is known as a conic section. The line L is known as the directrix and the point \mathbf{F} is the focus. e is defined to be the eccentricity of the conic.

(a) For $e = 1$, the conic is a parabola

(b) For $e < 1$, the conic is an ellipse

(c) For $e > 1$, the conic is a hyperbola

D.1.2. The equation of a conic with directrix $\mathbf{n}^\top \mathbf{x} = c$, eccentricity e and focus \mathbf{F} is given by

$$\mathbf{x}^\top \mathbf{V} \mathbf{x} + 2\mathbf{u}^\top \mathbf{x} + f = 0 \quad (\text{D.3})$$

where

$$\mathbf{V} = \|\mathbf{n}\|^2 \mathbf{I} - e^2 \mathbf{n} \mathbf{n}^\top, \quad (\text{D.4})$$

$$\mathbf{u} = ce^2 \mathbf{n} - \|\mathbf{n}\|^2 \mathbf{F}, \quad (\text{D.5})$$

$$f = \|\mathbf{n}\|^2 \|\mathbf{F}\|^2 - c^2 e^2 \quad (\text{D.6})$$

Proof. Using Definition D.1.1 and Lemma C.11, for any point \mathbf{x} on the conic,

$$\|\mathbf{x} - \mathbf{F}\|^2 = e^2 \frac{(\mathbf{n}^\top \mathbf{x} - c)^2}{\|\mathbf{n}\|^2} \quad (\text{D.7})$$

$$\implies \|\mathbf{n}\|^2 (\mathbf{x} - \mathbf{F})^\top (\mathbf{x} - \mathbf{F}) = e^2 (\mathbf{n}^\top \mathbf{x} - c)^2 \quad (\text{D.8})$$

$$\implies \|\mathbf{n}\|^2 (\mathbf{x}^\top \mathbf{x} - 2\mathbf{F}^\top \mathbf{x} + \|\mathbf{F}\|^2) = e^2 \left(c^2 + (\mathbf{n}^\top \mathbf{x})^2 - 2c\mathbf{n}^\top \mathbf{x} \right) \quad (\text{D.9})$$

$$= e^2 \left(c^2 + (\mathbf{x}^\top \mathbf{n} \mathbf{n}^\top \mathbf{x}) - 2c\mathbf{n}^\top \mathbf{x} \right) \quad (\text{D.10})$$

which can be expressed as (D.3) after simplification.

□

D.1.3. The eccentricity, directrices and foci of (D.3) are given by

$$e = \sqrt{1 - \frac{\lambda_1}{\lambda_2}} \quad (\text{D.11})$$

$$\mathbf{n} = \sqrt{\lambda_2} \mathbf{p}_1, \quad c = \begin{cases} \frac{e \mathbf{u}^\top \mathbf{n} \pm \sqrt{e^2 (\mathbf{u}^\top \mathbf{n})^2 - \lambda_2 (e^2 - 1) (\|\mathbf{u}\|^2 - \lambda_2 f)}}{\lambda_2 e (e^2 - 1)} & e \neq 1 \\ \frac{\|\mathbf{u}\|^2 - \lambda_2 f}{2 \mathbf{u}^\top \mathbf{n}} & e = 1 \end{cases} \quad (\text{D.12})$$

$$\mathbf{F} = \frac{ce^2 \mathbf{n} - \mathbf{u}}{\lambda_2} \quad (\text{D.13})$$

Proof. From (D.4), using the fact that \mathbf{V} is symmetric with $\mathbf{V} = \mathbf{V}^\top$,

$$\mathbf{V}^\top \mathbf{V} = \left(\|\mathbf{n}\|^2 \mathbf{I} - e^2 \mathbf{n} \mathbf{n}^\top \right)^\top \left(\|\mathbf{n}\|^2 \mathbf{I} - e^2 \mathbf{n} \mathbf{n}^\top \right) \quad (\text{D.14})$$

$$\implies \mathbf{V}^2 = \|\mathbf{n}\|^4 \mathbf{I} + e^4 \mathbf{n} \mathbf{n}^\top \mathbf{n} \mathbf{n}^\top - 2e^2 \|\mathbf{n}\|^2 \mathbf{n} \mathbf{n}^\top \quad (\text{D.15})$$

$$= \|\mathbf{n}\|^4 \mathbf{I} + e^4 \|\mathbf{n}\|^2 \mathbf{n} \mathbf{n}^\top - 2e^2 \|\mathbf{n}\|^2 \mathbf{n} \mathbf{n}^\top \quad (\text{D.16})$$

$$= \|\mathbf{n}\|^4 \mathbf{I} + e^2 (e^2 - 2) \|\mathbf{n}\|^2 \mathbf{n} \mathbf{n}^\top \quad (\text{D.17})$$

$$= \|\mathbf{n}\|^4 \mathbf{I} + (e^2 - 2) \|\mathbf{n}\|^2 \left(\|\mathbf{n}\|^2 \mathbf{I} - \mathbf{V} \right) \quad (\text{D.18})$$

which can be expressed as

$$\mathbf{V}^2 + (e^2 - 2) \|\mathbf{n}\|^2 \mathbf{V} - (e^2 - 1) \|\mathbf{n}\|^4 \mathbf{I} = 0 \quad (\text{D.19})$$

Using the Cayley-Hamilton theorem, (D.19) results in the characteristic equation,

$$\lambda^2 - (2 - e^2) \|\mathbf{n}\|^2 \lambda + (1 - e^2) \|\mathbf{n}\|^4 = 0 \quad (\text{D.20})$$

which can be expressed as

$$\left(\frac{\lambda}{\|\mathbf{n}\|^2}\right)^2 - (2 - e^2) \left(\frac{\lambda}{\|\mathbf{n}\|^2}\right) + (1 - e^2) = 0 \quad (\text{D.21})$$

$$\implies \frac{\lambda}{\|\mathbf{n}\|^2} = 1 - e^2, 1 \quad (\text{D.22})$$

$$\text{or, } \lambda_2 = \|\mathbf{n}\|^2, \lambda_1 = (1 - e^2) \lambda_2 \quad (\text{D.23})$$

From (D.23), the eccentricity of (D.3) is given by (D.11). Multiplying both sides of (D.4) by \mathbf{n} ,

$$\mathbf{V}\mathbf{n} = \|\mathbf{n}\|^2 \mathbf{n} - e^2 \mathbf{n} \mathbf{n}^\top \mathbf{n} \quad (\text{D.24})$$

$$= \|\mathbf{n}\|^2 (1 - e^2) \mathbf{n} \quad (\text{D.25})$$

$$= \lambda_1 \mathbf{n} \quad (\text{D.26})$$

$$(\text{D.27})$$

from (D.23). Thus, λ_1 is the corresponding eigenvalue for \mathbf{n} . From (B.22) and (D.27), this implies that

$$\mathbf{p}_1 = \frac{\mathbf{n}}{\|\mathbf{n}\|} \quad (\text{D.28})$$

$$\text{or, } \mathbf{n} = \|\mathbf{n}\| \mathbf{p}_1 = \sqrt{\lambda_2} \mathbf{p}_1 \quad (\text{D.29})$$

from (D.23) . From (D.5) and (D.23),

$$\mathbf{F} = \frac{ce^2 \mathbf{n} - \mathbf{u}}{\lambda_2} \quad (\text{D.30})$$

$$\implies \|\mathbf{F}\|^2 = \frac{(ce^2 \mathbf{n} - \mathbf{u})^\top (ce^2 \mathbf{n} - \mathbf{u})}{\lambda_2^2} \quad (\text{D.31})$$

$$\implies \lambda_2^2 \|\mathbf{F}\|^2 = c^2 e^4 \lambda_2 - 2ce^2 \mathbf{u}^\top \mathbf{n} + \|\mathbf{u}\|^2 \quad (\text{D.32})$$

Also, (D.6) can be expressed as

$$\lambda_2 \|\mathbf{F}\|^2 = f + c^2 e^2 \quad (\text{D.33})$$

From (D.32) and (D.33),

$$c^2 e^4 \lambda_2 - 2ce^2 \mathbf{u}^\top \mathbf{n} + \|\mathbf{u}\|^2 = \lambda_2 (f + c^2 e^2) \quad (\text{D.34})$$

$$\implies \lambda_2 e^2 (e^2 - 1) c^2 - 2ce^2 \mathbf{u}^\top \mathbf{n} + \|\mathbf{u}\|^2 - \lambda_2 f = 0 \quad (\text{D.35})$$

yielding (D.13). □

D.1.4. (D.3) represents

(a) a parabola for $\left| \mathbf{V} \right| = 0$,

(b) ellipse for $\left| \mathbf{V} \right| > 0$ and

(c) hyperbola for $\left| \mathbf{V} \right| < 0$.

Proof. From (D.11),

$$\frac{\lambda_1}{\lambda_2} = 1 - e^2 \quad (\text{D.36})$$

Also,

$$\left| \mathbf{V} \right| = \lambda_1 \lambda_2 \quad (\text{D.37})$$

yielding Table D.2

□

| Eccentricity | Conic | Eigenvalue | Determinant | | |
|--------------|-----------|--------------------------------|-------------|--------------|-------|
| $e = 1$ | Parabola | $\lambda_1 = 0$ | | \mathbf{V} | $= 0$ |
| $e < 1$ | Ellipse | $\lambda_1 > 0, \lambda_2 > 0$ | | \mathbf{V} | > 0 |
| $e > 1$ | Hyperbola | $\lambda_1 < 0, \lambda_2 > 0$ | | \mathbf{V} | < 0 |

Table D.2:

D.2. Standard Form

D.2.1. Using the affine transformation in (B.19), the conic in (D.3) can be expressed in standard form as

$$\mathbf{y}^\top \left(\frac{\mathbf{D}}{f_0} \right) \mathbf{y} = 1 \quad |\mathbf{V}| \neq 0 \quad (\text{D.38})$$

$$\mathbf{y}^\top \mathbf{D} \mathbf{y} = -\eta \mathbf{e}_1^\top \mathbf{y} \quad |\mathbf{V}| = 0 \quad (\text{D.39})$$

where

$$f_0 = \mathbf{u}^\top \mathbf{V}^{-1} \mathbf{u} - f \neq 0 \quad (\text{D.40})$$

$$\eta = 2\mathbf{u}^\top \mathbf{p}_1 \quad (\text{D.41})$$

$$\mathbf{e}_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (\text{D.42})$$

Proof. Using (B.19) (D.3) can be expressed as

$$(\mathbf{P}\mathbf{y} + \mathbf{c})^\top \mathbf{V} (\mathbf{P}\mathbf{y} + \mathbf{c}) + 2\mathbf{u}^\top (\mathbf{P}\mathbf{y} + \mathbf{c}) + f = 0, \quad (\text{D.43})$$

yielding

$$\mathbf{y}^\top \mathbf{P}^\top \mathbf{V} \mathbf{P} \mathbf{y} + 2(\mathbf{V}\mathbf{c} + \mathbf{u})^\top \mathbf{P} \mathbf{y} + \mathbf{c}^\top \mathbf{V} \mathbf{c} + 2\mathbf{u}^\top \mathbf{c} + f = 0 \quad (\text{D.44})$$

From (D.44) and (B.20),

$$\mathbf{y}^\top \mathbf{D} \mathbf{y} + 2(\mathbf{V}\mathbf{c} + \mathbf{u})^\top \mathbf{P} \mathbf{y} + \mathbf{c}^\top (\mathbf{V}\mathbf{c} + \mathbf{u}) + \mathbf{u}^\top \mathbf{c} + f = 0 \quad (\text{D.45})$$

When \mathbf{V}^{-1} exists, choosing

$$\mathbf{V}\mathbf{c} + \mathbf{u} = \mathbf{0}, \quad \text{or, } \mathbf{c} = -\mathbf{V}^{-1}\mathbf{u}, \quad (\text{D.46})$$

and substituting (D.46) in (D.45) yields (D.38). When $|\mathbf{V}| = 0, \lambda_1 = 0$ and

$$\mathbf{V}\mathbf{p}_1 = \mathbf{0}, \mathbf{V}\mathbf{p}_2 = \lambda_2 \mathbf{p}_2. \quad (\text{D.47})$$

where $\mathbf{p}_1, \mathbf{p}_2$ are the eigenvectors of \mathbf{V} such that (B.20)

$$\mathbf{P} = \begin{pmatrix} \mathbf{p}_1 & \mathbf{p}_2 \end{pmatrix}, \quad (\text{D.48})$$

Substituting (D.48) in (D.45),

$$\mathbf{y}^\top \mathbf{D} \mathbf{y} + 2 \left(\mathbf{c}^\top \mathbf{V} + \mathbf{u}^\top \right) \begin{pmatrix} \mathbf{p}_1 & \mathbf{p}_2 \end{pmatrix} \mathbf{y} + \mathbf{c}^\top (\mathbf{V} \mathbf{c} + \mathbf{u}) + \mathbf{u}^\top \mathbf{c} + f = 0 \quad (\text{D.49})$$

$$\implies \mathbf{y}^\top \mathbf{D} \mathbf{y} + 2 \left((\mathbf{c}^\top \mathbf{V} + \mathbf{u}^\top) \mathbf{p}_1 (\mathbf{c}^\top \mathbf{V} + \mathbf{u}^\top) \mathbf{p}_2 \right) \mathbf{y} + \mathbf{c}^\top (\mathbf{V} \mathbf{c} + \mathbf{u}) + \mathbf{u}^\top \mathbf{c} + f = 0 \quad (\text{D.50})$$

$$\implies \mathbf{y}^\top \mathbf{D} \mathbf{y} + 2 \left(\mathbf{u}^\top \mathbf{p}_1 \quad (\lambda_2 \mathbf{c}^\top + \mathbf{u}^\top) \mathbf{p}_2 \right) \mathbf{y} + \mathbf{c}^\top (\mathbf{V} \mathbf{c} + \mathbf{u}) + \mathbf{u}^\top \mathbf{c} + f = 0 \quad (\text{D.51})$$

upon substituting from (D.47) yielding

$$\lambda_2 y_2^2 + 2 \left(\mathbf{u}^\top \mathbf{p}_1 \right) y_1 + 2 y_2 (\lambda_2 \mathbf{c} + \mathbf{u})^\top \mathbf{p}_2 + \mathbf{c}^\top (\mathbf{V} \mathbf{c} + \mathbf{u}) + \mathbf{u}^\top \mathbf{c} + f = 0 \quad (\text{D.52})$$

Thus, (D.52) can be expressed as (D.39) by choosing

$$\eta = 2 \mathbf{u}^\top \mathbf{p}_1 \quad (\text{D.53})$$

and \mathbf{c} in (D.45) such that

$$2 \mathbf{P}^\top (\mathbf{V} \mathbf{c} + \mathbf{u}) = \eta \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (\text{D.54})$$

$$\mathbf{c}^\top (\mathbf{V} \mathbf{c} + \mathbf{u}) + \mathbf{u}^\top \mathbf{c} + f = 0 \quad (\text{D.55})$$

$\therefore \mathbf{P}^\top \mathbf{P} = \mathbf{I}$, multiplying (D.54) by \mathbf{P} yields

$$(\mathbf{V}\mathbf{c} + \mathbf{u}) = \frac{\eta}{2}\mathbf{p}_1, \quad (\text{D.56})$$

which, upon substituting in (D.55) results in

$$\frac{\eta}{2}\mathbf{c}^\top \mathbf{p}_1 + \mathbf{u}^\top \mathbf{c} + f = 0 \quad (\text{D.57})$$

(D.56) and (D.57) can be clubbed together to obtain (E.7). \square

D.2.2. For the standard conic,

$$\mathbf{P} = \mathbf{I} \quad (\text{D.58})$$

$$\mathbf{u} = \begin{cases} 0 & e \neq 1 \\ \frac{\eta}{2}\mathbf{e}_1 & e = 1 \end{cases} \quad (\text{D.59})$$

$$\lambda_1 \begin{cases} = 0 & e = 1 \\ \neq 0 & e \neq 1 \end{cases} \quad (\text{D.60})$$

where

$$\mathbf{I} = \begin{pmatrix} \mathbf{e}_1 & \mathbf{e}_2 \end{pmatrix} \quad (\text{D.61})$$

is the identity matrix.

D.2.3.

(a) The directrices for the standard conic are given by

$$\mathbf{e}_1^\top \mathbf{y} = \pm \frac{1}{e} \sqrt{\frac{|f_0|}{\lambda_2 (1 - e^2)}} \quad e \neq 1 \quad (\text{D.62})$$

$$\mathbf{e}_1^\top \mathbf{y} = \frac{\eta}{2\lambda_2} \quad e = 1 \quad (\text{D.63})$$

(b) The foci of the standard ellipse and hyperbola are given by

$$\mathbf{F} = \begin{cases} \pm e \sqrt{\frac{|f_0|}{\lambda_2 (1 - e^2)}} \mathbf{e}_1 & e \neq 1 \\ -\frac{\eta}{4\lambda_2} \mathbf{e}_1 & e = 1 \end{cases} \quad (\text{D.64})$$

Proof. (a) For the standard hyperbola/ellipse in (D.38), from (D.58), (D.12) and (D.59),

$$\mathbf{n} = \sqrt{\frac{\lambda_2}{f_0}} \mathbf{e}_1 \quad (\text{D.65})$$

$$c = \pm \frac{\sqrt{-\frac{\lambda_2}{f_0} (e^2 - 1) \left(\frac{\lambda_2}{f_0}\right)}}{\frac{\lambda_2}{f_0} e (e^2 - 1)} \quad (\text{D.66})$$

$$= \pm \frac{1}{e \sqrt{1 - e^2}} \quad (\text{D.67})$$

yielding (D.62) upon substituting from (D.11) and simplifying. For the standard parabola in (D.39), from (D.58), (D.12) and (D.59), noting that $f = 0$,

$$\mathbf{n} = \sqrt{\lambda_2} \mathbf{e}_1 \quad (\text{D.68})$$

$$c = \frac{\left\| \frac{\eta}{2} \mathbf{e}_1 \right\|^2}{2 \left(\frac{\eta}{2} \right) (\mathbf{e}_1)^\top \mathbf{n}} \quad (\text{D.69})$$

$$(\text{D.70})$$

$$= \frac{\eta}{4\sqrt{\lambda_2}} \quad (\text{D.71})$$

yielding (D.63).

- (b) For the standard ellipse/hyperbola, substituting from (D.67), (D.65), (D.59) and (D.11) in (D.13),

$$\mathbf{F} = \pm \frac{\left(\frac{1}{e\sqrt{1-e^2}} \right) (e^2) \sqrt{\frac{\lambda_2}{f_0}} \mathbf{e}_1}{\frac{\lambda_2}{f_0}} \quad (\text{D.72})$$

yielding (D.64) after simplification. For the standard parabola, substituting from (D.71), (D.68), (D.59) and (D.11) in (D.13),

$$\mathbf{F} = \frac{\left(\frac{\eta}{4\sqrt{\lambda_2}} \right) \sqrt{\lambda_2} \mathbf{e}_1 - \frac{\eta}{2} \mathbf{e}_1}{\lambda_2} \quad (\text{D.73})$$

$$(\text{D.74})$$

yielding (D.64) after simplification.

□

Appendix E

Conic Parameters

E.1. Standard Form

E.1.1. The center of the standard ellipse/hyperbola, defined to be the mid point of the line joining the foci, is the origin.

E.1.2. The principal (major) axis of the standard ellipse/hyperbola, defined to be the line joining the two foci is the x -axis.

Proof. From (D.64), it is obvious that the line joining the foci passes through the origin. Also, the direction vector of this line is \mathbf{e}_1 . Thus, the principal axis is the x -axis. \square

E.1.3. The minor axis of the standard ellipse/hyperbola, defined to be the line orthogonal to the x -axis is the y -axis.

E.1.4. The axis of symmetry of the standard parabola, defined to be the line perpendicular to the directrix and passing through the focus, is the x -axis.

Proof. From (D.68) and (D.64), the axis of the parabola can be expressed using (C.2)

as

$$\mathbf{e}_2^\top \left(\mathbf{y} + \frac{\eta}{4\lambda_2} \mathbf{e}_1 \right) = 0 \quad (\text{E.1})$$

$$\implies \mathbf{e}_2^\top \mathbf{y} = 0, \quad (\text{E.2})$$

which is the equation of the x -axis. \square

E.1.5. The point where the parabola intersects its axis of symmetry is called the vertex. For the standard parabola, the vertex is the origin.

Proof. (E.2) can be expressed as

$$\mathbf{y} = \alpha \mathbf{e}_1, \quad (\text{E.3})$$

using (C.2). Substituting (E.3) in (D.39),

$$\alpha^2 \mathbf{e}_1^\top \mathbf{D} \mathbf{e}_1 = -\eta \alpha \mathbf{e}_1^\top \mathbf{e}_1 \quad (\text{E.4})$$

$$\implies \alpha = 0, \text{ or, } \mathbf{y} = \mathbf{0}. \quad (\text{E.5})$$

\square

E.1.6. The focal length of the standard parabola, , defined to be the distance between the vertex and the focus, measured along the axis of symmetry, is $\left| \frac{\eta}{4\lambda_2} \right|$

E.2. Quadratic Form

E.2.1. The center/vertex of a conic section are given by

$$\mathbf{c} = -\mathbf{V}^{-1}\mathbf{u} \quad \left| \mathbf{V} \right| \neq 0 \quad (\text{E.6})$$

$$\begin{pmatrix} \mathbf{u}^\top + \frac{\eta}{2}\mathbf{p}_1^\top \\ \mathbf{v} \end{pmatrix} \mathbf{c} = \begin{pmatrix} -f \\ \frac{\eta}{2}\mathbf{p}_1 - \mathbf{u} \end{pmatrix} \quad \left| \mathbf{V} \right| = 0 \quad (\text{E.7})$$

Proof. In (B.19), substituting $\mathbf{y} = \mathbf{0}$, the center/vertex for the quadratic form is obtained as

$$\mathbf{x} = \mathbf{c}, \quad (\text{E.8})$$

where \mathbf{c} is derived as (E.6) and (E.7) in Appendix D.2.1. □

E.2.2. The equation of the minor and major axes for the ellipse/hyperbola are respectively given by

$$\mathbf{p}_i^\top (\mathbf{x} - \mathbf{c}) = 0, i = 1, 2 \quad (\text{E.9})$$

The axis of symmetry for the parabola is also given by (E.9).

Proof. From (E.1.2), the major/symmetry axis for the hyperbola/ellipse/parabola can be expressed using (B.19) as

$$\mathbf{e}_2^\top \mathbf{P}^\top (\mathbf{x} - \mathbf{c}) = 0 \quad (\text{E.10})$$

$$\implies (\mathbf{P}\mathbf{e}_2)^\top (\mathbf{x} - \mathbf{c}) = 0 \quad (\text{E.11})$$

yielding (E.9), and the proof for the minor axis is similar.

□

Appendix F

Conic Lines

F.1. Pair of Straight Lines

F.1.1. The asymptotes of the hyperbola in (D.38), defined to be the lines that do not intersect the hyperbola, are given by

$$\left(\sqrt{|\lambda_1|} \pm \sqrt{|\lambda_2|} \right) \mathbf{y} = 0 \quad (\text{F.1})$$

Proof. From (D.38), it is obvious that the pair of lines represented by

$$\mathbf{y}^\top \mathbf{D} \mathbf{y} = 0 \quad (\text{F.2})$$

do not intersect the conic

$$\mathbf{y}^\top \mathbf{D} \mathbf{y} = f_0 \quad (\text{F.3})$$

Thus, (F.2) represents the asymptotes of the hyperbola in (D.38) and can be expressed as

$$\lambda_1 y_1^2 + \lambda_2 y_1^2 = 0, \quad (\text{F.4})$$

which can then be simplified to obtain (F.1).

□

F.1.2. (D.3) represents a pair of straight lines if

$$\mathbf{u}^\top \mathbf{V}^{-1} \mathbf{u} - f = 0 \quad (\text{F.5})$$

F.1.3. (D.3) represents a pair of straight lines if the matrix

$$\begin{pmatrix} \mathbf{V} & \mathbf{u} \\ \mathbf{u} & f \end{pmatrix} \quad (\text{F.6})$$

is singular.

Proof. Let

$$\begin{pmatrix} \mathbf{V} & \mathbf{u} \\ \mathbf{u} & f \end{pmatrix} \mathbf{x} = \mathbf{0} \quad (\text{F.7})$$

Expressing

$$\mathbf{x} = \begin{pmatrix} \mathbf{y} \\ y_3 \end{pmatrix}, \quad (\text{F.8})$$

$$\begin{pmatrix} \mathbf{V} & \mathbf{u} \\ \mathbf{u}^\top & f \end{pmatrix} \begin{pmatrix} \mathbf{y} \\ y_3 \end{pmatrix} = \mathbf{0} \quad (\text{F.9})$$

$$\implies \mathbf{V}\mathbf{y} + y_3\mathbf{u} = \mathbf{0} \quad \text{and} \quad (\text{F.10})$$

$$\mathbf{u}^\top \mathbf{y} + f y_3 = 0 \quad (\text{F.11})$$

From (F.10) we obtain,

$$\mathbf{y}^\top \mathbf{V}\mathbf{y} + y_3 \mathbf{y}^\top \mathbf{u} = 0 \quad (\text{F.12})$$

$$\implies \mathbf{y}^\top \mathbf{V}\mathbf{y} + y_3 \mathbf{u}^\top \mathbf{y} = 0 \quad (\text{F.13})$$

yielding (F.5) upon substituting from (F.11). □

F.1.4. Using the affine transformation, (F.1) can be expressed as the lines

$$\left(\sqrt{|\lambda_1|} \quad \pm \sqrt{|\lambda_2|} \right) \mathbf{P}^\top (\mathbf{x} - \mathbf{c}) = 0 \quad (\text{F.14})$$

F.1.5. The angle between the asymptotes can be expressed as

$$\cos \theta = \frac{|\lambda_1| - |\lambda_2|}{|\lambda_1| + |\lambda_2|} \quad (\text{F.15})$$

Proof. The normal vectors of the lines in (F.14) are

$$\begin{aligned} \mathbf{n}_1 &= \mathbf{P} \begin{pmatrix} \sqrt{|\lambda_1|} \\ \sqrt{|\lambda_2|} \end{pmatrix} \\ \mathbf{n}_2 &= \mathbf{P} \begin{pmatrix} \sqrt{|\lambda_1|} \\ -\sqrt{|\lambda_2|} \end{pmatrix} \end{aligned} \quad (\text{F.16})$$

The angle between the asymptotes is given by

$$\cos \theta = \frac{\mathbf{n}_1^\top \mathbf{n}_2}{\|\mathbf{n}_1\| \|\mathbf{n}_2\|} \quad (\text{F.17})$$

The orthogonal matrix \mathbf{P} preserves the norm, i.e.

$$\|\mathbf{n}_1\| = \left\| \mathbf{P} \begin{pmatrix} \sqrt{|\lambda_1|} \\ \sqrt{|\lambda_2|} \end{pmatrix} \right\| = \left\| \begin{pmatrix} \sqrt{|\lambda_1|} \\ \sqrt{|\lambda_2|} \end{pmatrix} \right\| \quad (\text{F.18})$$

$$= \sqrt{|\lambda_1| + |\lambda_2|} = \|\mathbf{n}_2\| \quad (\text{F.19})$$

It is easy to verify that

$$\mathbf{n}_1^\top \mathbf{n}_2 = |\lambda_1| - |\lambda_2| \quad (\text{F.20})$$

Thus, the angle between the asymptotes is obtained from (F.17) as (F.15). \square

F.2. Intersection of Conics

F.2.1. Let

$$\mathbf{x}^\top \mathbf{V}_i \mathbf{x} + 2\mathbf{u}_i^\top \mathbf{x} + f_i = 0, \quad i = 1, 2 \quad (\text{F.21})$$

be the equation of two conics. The locus of their intersection is a pair of straight lines if

$$\begin{vmatrix} \mathbf{V}_1 + \mu \mathbf{V}_2 & \mathbf{u}_1 + \mu \mathbf{u}_2 \\ (\mathbf{u}_1 + \mu \mathbf{u}_2)^\top & f \end{vmatrix} = 0, \quad \left| \mathbf{V}_1 + \mu \mathbf{V}_2 \right| < 0 \quad (\text{F.22})$$

Proof. The intersection of the conics in (F.21) is given by the curve

$$\mathbf{x}^\top (\mathbf{V}_1 + \mu \mathbf{V}_2) \mathbf{x} + 2 (\mathbf{u}_1 + \mu \mathbf{u}_2)^\top \mathbf{x} + f_1 + \mu f_2 = 0, \quad (\text{F.23})$$

which, from Theorem F.1.3 represents a pair of straight lines if (F.22) is satisfied. \square

F.2.2. The points of intersection of the conics in (F.21) are the points of the intersection of the lines in (F.23).

F.3. Chords of a Conic

F.3.1. The points of intersection of the line

$$L : \quad \mathbf{x} = \mathbf{q} + \mu \mathbf{m} \quad \mu \in \mathbb{R} \quad (\text{F.24})$$

with the conic section in (D.3) are given by

$$\mathbf{x}_i = \mathbf{q} + \mu_i \mathbf{m} \quad (\text{F.25})$$

where

$$\mu_i = \frac{1}{\mathbf{m}^\top \mathbf{V} \mathbf{m}} \left(-\mathbf{m}^\top (\mathbf{V} \mathbf{q} + \mathbf{u}) \pm \sqrt{[\mathbf{m}^\top (\mathbf{V} \mathbf{q} + \mathbf{u})]^2 - (\mathbf{q}^\top \mathbf{V} \mathbf{q} + 2\mathbf{u}^\top \mathbf{q} + f)(\mathbf{m}^\top \mathbf{V} \mathbf{m})} \right) \quad (\text{F.26})$$

Proof. Substituting (F.24) in (D.3),

$$(\mathbf{q} + \mu \mathbf{m})^\top \mathbf{V} (\mathbf{q} + \mu \mathbf{m}) + 2\mathbf{u}^\top (\mathbf{q} + \mu \mathbf{m}) + f = 0 \quad (\text{F.27})$$

$$\implies \mu^2 \mathbf{m}^\top \mathbf{V} \mathbf{m} + 2\mu \mathbf{m}^\top (\mathbf{V} \mathbf{q} + \mathbf{u}) + \mathbf{q}^\top \mathbf{V} \mathbf{q} + 2\mathbf{u}^\top \mathbf{q} + f = 0 \quad (\text{F.28})$$

Solving the above quadratic in (F.28) yields (F.26). \square

F.3.2. If L in (F.24) touches (D.3) at exactly one point \mathbf{q} ,

$$\mathbf{m}^\top (\mathbf{V} \mathbf{q} + \mathbf{u}) = 0 \quad (\text{F.29})$$

Proof. In this case, (F.28) has exactly one root. Hence, in (F.26)

$$\left[\mathbf{m}^\top (\mathbf{V} \mathbf{q} + \mathbf{u}) \right]^2 - \left(\mathbf{m}^\top \mathbf{V} \mathbf{m} \right) \left(\mathbf{q}^\top \mathbf{V} \mathbf{q} + 2\mathbf{u}^\top \mathbf{q} + f \right) = 0 \quad (\text{F.30})$$

$\because \mathbf{q}$ is the point of contact, \mathbf{q} satisfies (D.3) and

$$\mathbf{q}^\top \mathbf{V} \mathbf{q} + 2\mathbf{u}^\top \mathbf{q} + f = 0 \quad (\text{F.31})$$

Substituting (F.31) in (F.30) and simplifying, we obtain (F.29). \square

F.3.3. The length of the chord in (F.24) is given by

$$\frac{2\sqrt{[\mathbf{m}^\top (\mathbf{V}\mathbf{q} + \mathbf{u})]^2 - (\mathbf{q}^\top \mathbf{V}\mathbf{q} + 2\mathbf{u}^\top \mathbf{q} + f)(\mathbf{m}^\top \mathbf{V}\mathbf{m})}}{\mathbf{m}^\top \mathbf{V}\mathbf{m}} \|\mathbf{m}\| \quad (\text{F.32})$$

Proof. The distance between the points in (F.25) is given by

$$\|\mathbf{x}_1 - \mathbf{x}_2\| = |\mu_1 - \mu_2| \|\mathbf{m}\| \quad (\text{F.33})$$

Substituting μ_i from (F.26) in (F.33) yields (F.32). \square

F.3.4. The affine transform for the conic section, preserves the norm. This implies that the length of any chord of a conic is invariant to translation and/or rotation.

Proof. Let

$$\mathbf{x}_i = \mathbf{P}\mathbf{y}_i + \mathbf{c} \quad (\text{F.34})$$

be any two points on the conic. Then the distance between the points is given by

$$\|\mathbf{x}_1 - \mathbf{x}_2\| = \|\mathbf{P}(\mathbf{y}_1 - \mathbf{y}_2)\| \quad (\text{F.35})$$

which can be expressed as

$$\|\mathbf{x}_1 - \mathbf{x}_2\|^2 = (\mathbf{y}_1 - \mathbf{y}_2)^\top \mathbf{P}^\top \mathbf{P} (\mathbf{y}_1 - \mathbf{y}_2) \quad (\text{F.36})$$

$$= \|\mathbf{y}_1 - \mathbf{y}_2\|^2 \quad (\text{F.37})$$

since

$$\mathbf{P}^\top \mathbf{P} = \mathbf{I} \quad (\text{F.38})$$

□

F.3.5. For the standard hyperbola/ellipse, the length of the major axis is

$$2\sqrt{\left|\frac{f_0}{\lambda_1}\right|} \quad (\text{F.39})$$

and the minor axis is

$$2\sqrt{\left|\frac{f_0}{\lambda_2}\right|} \quad (\text{F.40})$$

Proof. Since the major axis passes through the origin,

$$\mathbf{q} = \mathbf{0} \quad (\text{F.41})$$

Further, from Corollary (E.1.2),

$$\mathbf{m} = \mathbf{e}_2, \quad (\text{F.42})$$

and from (D.38),

$$\mathbf{V} = \frac{\mathbf{D}}{f_0}, \mathbf{u} = 0, f = -1 \quad (\text{F.43})$$

Substituting the above in (F.32),

$$\frac{2\sqrt{\mathbf{e}_1^\top \frac{\mathbf{D}}{f_0} \mathbf{e}_1}}{\mathbf{e}_1^\top \frac{\mathbf{D}}{f_0} \mathbf{e}_1} \|\mathbf{e}_1\| \quad (\text{F.44})$$

yielding (F.39). Similarly, for the minor axis, the only different parameter is

$$\mathbf{m} = \mathbf{e}_2, \quad (\text{F.45})$$

Substituting the above in (F.32),

$$\frac{2\sqrt{\mathbf{e}_2^\top \frac{\mathbf{D}}{f_0} \mathbf{e}_2}}{\mathbf{e}_2^\top \frac{\mathbf{D}}{f_0} \mathbf{e}_2} \|\mathbf{e}_2\| \quad (\text{F.46})$$

yielding (F.40).

□

F.3.6. The latus rectum of a conic section is the chord that passes through the focus and is perpendicular to the major axis. The length of the latus rectum for a conic is given by

$$l = \begin{cases} 2\frac{\sqrt{|f_0\lambda_1|}}{\lambda_2} & e \neq 1 \\ \frac{\eta}{\lambda_2} & e = 1 \end{cases} \quad (\text{F.47})$$

Proof. The latus rectum is perpendicular to the major axis for the standard conic. Hence, from Corollary (E.1.2),

$$\mathbf{m} = \mathbf{e}_2, \quad (\text{F.48})$$

Since it passes through the focus, from (D.64)

$$\mathbf{q} = \mathbf{F} = \pm e \sqrt{\frac{f_0}{\lambda_2(1-e^2)}} \mathbf{e}_1 \quad (\text{F.49})$$

for the standard hyperbola/ellipse. Also, from (D.38),

$$\mathbf{V} = \frac{\mathbf{D}}{f_0}, \mathbf{u} = 0, f = -1 \quad (\text{F.50})$$

Substituting the above in (F.32),

$$\frac{2\sqrt{\left[\mathbf{e}_2^\top \left(\frac{\mathbf{D}}{f_0} e \sqrt{\frac{f_0}{\lambda_2(1-e^2)}} \mathbf{e}_1\right)\right]^2 - \left(e \sqrt{\frac{f_0}{\lambda_2(1-e^2)}} \mathbf{e}_1^\top \frac{\mathbf{D}}{f_0} e \sqrt{\frac{f_0}{\lambda_2(1-e^2)}} \mathbf{e}_1 - 1\right) \left(\mathbf{e}_2^\top \frac{\mathbf{D}}{f_0} \mathbf{e}_2\right)}}{\mathbf{e}_2^\top \frac{\mathbf{D}}{f_0} \mathbf{e}_2} \|\mathbf{e}_2\| \quad (\text{F.51})$$

Since

$$\mathbf{e}_2^\top \mathbf{D} \mathbf{e}_1 = 0, \mathbf{e}_1^\top \mathbf{D} \mathbf{e}_1 = \lambda_1, \mathbf{e}_1^\top \mathbf{e}_1 = 1, \|\mathbf{e}_2\| = 1, \mathbf{e}_2^\top \mathbf{D} \mathbf{e}_2 = \lambda_2, \quad (\text{F.52})$$

(F.51) can be expressed as

$$\frac{2\sqrt{\left(1 - \frac{\lambda_1 e^2}{\lambda_2(1-e^2)}\right) \left(\frac{\lambda_2}{f_0}\right)}}{\frac{\lambda_2}{f_0}} \quad (\text{F.53})$$

$$= 2 \frac{\sqrt{f_0 \lambda_1}}{\lambda_2} \quad \left(\because e^2 = 1 - \frac{\lambda_1}{\lambda_2}\right) \quad (\text{F.54})$$

For the standard parabola, the parameters in (F.32) are

$$\mathbf{q} = \mathbf{F} = -\frac{\eta}{4\lambda_2} \mathbf{e}_1, \mathbf{m} = \mathbf{e}_1, \mathbf{V} = \mathbf{D}, \mathbf{u} = \frac{\eta}{2} \mathbf{e}_1^\top, f = 0 \quad (\text{F.55})$$

Substituting the above in (F.32), the length of the latus rectum can be expressed as

$$\frac{2\sqrt{\left[\mathbf{e}_2^\top \left(\mathbf{D} \left(-\frac{\eta}{4\lambda_2} \mathbf{e}_1\right) + \frac{\eta}{2} \mathbf{e}_1\right)\right]^2 - \left(\left(-\frac{\eta}{4\lambda_2} \mathbf{e}_1\right)^\top \mathbf{D} \left(-\frac{\eta}{4\lambda_2} \mathbf{e}_1\right) + 2\frac{\eta}{2} \mathbf{e}_1^\top \left(-\frac{\eta}{4\lambda_2} \mathbf{e}_1\right)\right) (\mathbf{e}_2^\top \mathbf{D} \mathbf{e}_2)}}{\mathbf{e}_2^\top \mathbf{D} \mathbf{e}_2}} \|\mathbf{e}_2\| \quad (\text{F.56})$$

Since

$$\mathbf{e}_2^\top \mathbf{D} \mathbf{e}_1 = 0, \mathbf{e}_2^\top \mathbf{e}_2 = 0, \mathbf{e}_1^\top \mathbf{D} \mathbf{e}_1 = 0, \mathbf{e}_1^\top \mathbf{e}_1 = 1, \|\mathbf{e}_1\| = 1, \mathbf{e}_2^\top \mathbf{D} \mathbf{e}_2 = \lambda_2, \quad (\text{F.57})$$

(F.56) can be expressed as

$$2 \frac{\sqrt{\frac{\eta^2}{4\lambda_2} \lambda_2}}{\lambda_2} = \frac{\eta}{\lambda_2} \quad (\text{F.58})$$

□

F.4. Tangent and Normal

F.4.1. Given the point of contact \mathbf{q} , the equation of a tangent to (D.3) is

$$(\mathbf{V} \mathbf{q} + \mathbf{u})^\top \mathbf{x} + \mathbf{u}^\top \mathbf{q} + f = 0 \quad (\text{F.59})$$

Proof. The normal vector is obtained from (F.29) and (A.28) as

$$\kappa \mathbf{n} = \mathbf{V} \mathbf{q} + \mathbf{u}, \kappa \in \mathbb{R} \quad (\text{F.60})$$

From (F.60) and (C.2), the equation of the tangent is

$$(\mathbf{V}\mathbf{q} + \mathbf{u})^\top (\mathbf{x} - \mathbf{q}) = 0 \quad (\text{F.61})$$

$$\implies (\mathbf{V}\mathbf{q} + \mathbf{u})^\top \mathbf{x} - \mathbf{q}^\top \mathbf{V}\mathbf{q} - \mathbf{u}^\top \mathbf{q} = 0 \quad (\text{F.62})$$

which, upon substituting from (F.31) and simplifying yields (F.59) \square

F.4.2. If \mathbf{V}^{-1} exists, given the normal vector \mathbf{n} , the tangent points of contact to (D.3) are given by

$$\begin{aligned} \mathbf{q}_i &= \mathbf{V}^{-1} (\kappa_i \mathbf{n} - \mathbf{u}), i = 1, 2 \\ \text{where } \kappa_i &= \pm \sqrt{\frac{f_0}{\mathbf{n}^\top \mathbf{V}^{-1} \mathbf{n}}} \end{aligned} \quad (\text{F.63})$$

Proof. From (F.60),

$$\mathbf{q} = \mathbf{V}^{-1} (\kappa \mathbf{n} - \mathbf{u}), \quad \kappa \in \mathbb{R} \quad (\text{F.64})$$

Substituting (F.64) in (F.31),

$$(\kappa \mathbf{n} - \mathbf{u})^\top \mathbf{V}^{-1} (\kappa \mathbf{n} - \mathbf{u}) + 2\mathbf{u}^\top \mathbf{V}^{-1} (\kappa \mathbf{n} - \mathbf{u}) + f = 0 \quad (\text{F.65})$$

$$\implies \kappa^2 \mathbf{n}^\top \mathbf{V}^{-1} \mathbf{n} - \mathbf{u}^\top \mathbf{V}^{-1} \mathbf{u} + f = 0 \quad (\text{F.66})$$

$$\text{or, } \kappa = \pm \sqrt{\frac{f_0}{\mathbf{n}^\top \mathbf{V}^{-1} \mathbf{n}}} \quad (\text{F.67})$$

Substituting (F.67) in (F.64) yields (F.63). \square

F.4.3. If \mathbf{V} is not invertible, given the normal vector \mathbf{n} , the point of contact to (D.3) is given

by the matrix equation

$$\begin{pmatrix} (\mathbf{u} + \kappa \mathbf{n})^\top \\ \mathbf{V} \end{pmatrix} \mathbf{q} = \begin{pmatrix} -f \\ \kappa \mathbf{n} - \mathbf{u} \end{pmatrix} \quad (\text{F.68})$$

$$\text{where } \kappa = \frac{\mathbf{p}_1^\top \mathbf{u}}{\mathbf{p}_1^\top \mathbf{n}}, \quad \mathbf{V} \mathbf{p}_1 = 0 \quad (\text{F.69})$$

Proof. If \mathbf{V} is non-invertible, it has a zero eigenvalue. If the corresponding eigenvector is \mathbf{p}_1 , then,

$$\mathbf{V} \mathbf{p}_1 = 0 \quad (\text{F.70})$$

From (F.60),

$$\kappa \mathbf{n} = \mathbf{V} \mathbf{q} + \mathbf{u}, \quad \kappa \in \mathbb{R} \quad (\text{F.71})$$

$$\implies \kappa \mathbf{p}_1^\top \mathbf{n} = \mathbf{p}_1^\top \mathbf{V} \mathbf{q} + \mathbf{p}_1^\top \mathbf{u} \quad (\text{F.72})$$

$$\text{or, } \kappa \mathbf{p}_1^\top \mathbf{n} = \mathbf{p}_1^\top \mathbf{u}, \quad \because \mathbf{p}_1^\top \mathbf{V} = 0, \quad (\text{from (F.70)}) \quad (\text{F.73})$$

yielding κ in (F.69). From (F.71),

$$\kappa \mathbf{q}^\top \mathbf{n} = \mathbf{q}^\top \mathbf{V} \mathbf{q} + \mathbf{q}^\top \mathbf{u} \quad (\text{F.74})$$

$$\implies \kappa \mathbf{q}^\top \mathbf{n} = -f - \mathbf{q}^\top \mathbf{u} \quad \text{from (F.31)}, \quad (\text{F.75})$$

$$\text{or, } (\kappa \mathbf{n} + \mathbf{u})^\top \mathbf{q} = -f \quad (\text{F.76})$$

(F.71) can be expressed as

$$\mathbf{V} \mathbf{q} = \kappa \mathbf{n} - \mathbf{u}. \quad (\text{F.77})$$

(F.76) and (F.77) clubbed together result in (F.68). \square

F.4.4. The normal vectors of the tangents to the conic in (D.3) satisfy

$$\mathbf{n}^\top \mathbf{V}^{-1} \mathbf{n} - f_0 = 0 \quad (\text{F.78})$$

Proof. From (F.29), the normal vector to the tangent at \mathbf{q} can be expressed as

$$\mathbf{n} = \mathbf{V}\mathbf{q} + \mathbf{u} \quad (\text{F.79})$$

$$\implies \mathbf{q} = \mathbf{V}^{-1}(\mathbf{n} - \mathbf{u}) \quad (\text{F.80})$$

which upon substituting in (D.3) yields

$$(\mathbf{n} - \mathbf{u})^\top \mathbf{V}^{-1} \mathbf{V} \mathbf{V}^{-1} (\mathbf{n} - \mathbf{u}) + 2\mathbf{u}^\top \mathbf{V}^{-1} (\mathbf{n} - \mathbf{u}) + f = 0 \quad (\text{F.81})$$

which can be simplified to obtain (F.78). \square

F.4.5. The normal vectors of the tangents to the conic in (D.3) from a point \mathbf{h} are given by

Proof. Let the equation of the tangent be

$$\mathbf{n}^\top \mathbf{x} = c \quad (\text{F.82})$$

If \mathbf{q} be the point of contact, since \mathbf{h}, \mathbf{q} lie on (F.82),

$$\mathbf{n}^\top \mathbf{q} = \mathbf{n}^\top \mathbf{h} = c \quad (\text{F.83})$$

From (F.79), \square

F.4.6. The normal vectors of the tangents to the conic in (D.3) from a point \mathbf{h} are given by

$$\begin{aligned}\mathbf{n}_1 &= \mathbf{P} \begin{pmatrix} \sqrt{|\lambda_1|} \\ \sqrt{|\lambda_2|} \end{pmatrix} \\ \mathbf{n}_2 &= \mathbf{P} \begin{pmatrix} \sqrt{|\lambda_1|} \\ -\sqrt{|\lambda_2|} \end{pmatrix}\end{aligned}\tag{F.84}$$

where λ_i, \mathbf{P} are the eigenparameters of

$$\Sigma = (\mathbf{V}\mathbf{h} + \mathbf{u})(\mathbf{V}\mathbf{h} + \mathbf{u})^\top - \mathbf{V}(\mathbf{h}^\top \mathbf{V}\mathbf{h} + 2\mathbf{u}^\top \mathbf{h} + f).\tag{F.85}$$

Proof. From (F.26), and (F.30)

$$\left[\mathbf{m}^\top (\mathbf{V}\mathbf{h} + \mathbf{u})\right]^2 - (\mathbf{m}^\top \mathbf{V}\mathbf{m}) (\mathbf{h}^\top \mathbf{V}\mathbf{h} + 2\mathbf{u}^\top \mathbf{h} + f) = 0\tag{F.86}$$

$$\implies \mathbf{m}^\top \left[(\mathbf{V}\mathbf{h} + \mathbf{u})(\mathbf{V}\mathbf{h} + \mathbf{u})^\top - \mathbf{V}(\mathbf{h}^\top \mathbf{V}\mathbf{h} + 2\mathbf{u}^\top \mathbf{h} + f)\right] \mathbf{m} = 0\tag{F.87}$$

yielding (F.85). Consequently, from (F.16), (F.84) can be obtained. \square

