

A person is standing on a rocky shore at night, looking out at the ocean. The moon is bright in the dark sky, and its light reflects on the water. The person is silhouetted against the moonlight.

# Planned Field Campaigns

Jennifer Hutchings

Oregon State University

College of Earth Ocean and Atmospheric Sciences

[jhutchings@coas.oregonstate.edu](mailto:jhutchings@coas.oregonstate.edu)

Presenting Several Team Efforts




# Summary

- Needs: Gaps in knowledge / Model validation
- MOSAiC
- SIDEX
- Remote Sensing / Airborne Campaigns
- What's Missing?

The following summarizes my knowledge of field plans.  
We need your insight to design field campaigns that support you.



- 
- Models generally poor at reproducing sea ice deformation.
  - Hence may misrepresent ice thickness redistribution, wind stress transfer to the ocean, distribution of leads and ridges.
  - Lead and ridge spatial patterns not parameterized.



# Gaps in knowledge

- Interaction with coastal boundaries.
- Scaling properties of sea ice deformation below ~10km.
- How transitions in scaling properties relate to changes in mechanical properties of ice.
- Mechanisms of floe cracking.
- Stress propagation through mixtures of ice type.



# Gaps in knowledge

- Interaction with coastal boundaries. **NSF/NASA**
- Scaling properties of sea ice deformation below ~10km. **MOSAiC**
- How transitions in scaling properties relate to changes in mechanical properties of ice. **Ongoing**
- Mechanisms of floe cracking. **MOSAiC/SIDEx**
- Stress propagation through mixtures of ice type. **SIDEx + Modeling (proposed, unfunded)**
- There are more gaps e.g. anisotropy in ice structure, stress weakening by ridging, time scales of healing, stress propagation in mixed ice.




# Model Validation


Thoughts on validating Robert's model.

- Survey ice thickness, snow and area incorporated into ridges.
- Requires (1) high fidelity deformation and thickness measurements, (2) coordinated mapping and surveying.
- Ridge height statistics and shapes.
- How to estimate density on large scales? Snow is a large uncertainty.






JD 90 06:14 2007



JD 91 06:57 2007






JD 91 23:38 2007







# Ice – Coast Interaction








# SEDNA

- late March - end May 2007
- Coordinated strain-rate, thickness and stress measurement
- Campaign focus on stress / strain-rate relationship.







# GPS buoy array






# Spatial Scaling






# Space-Time Coupling






# Space-Time Coupling






# Space-Time Coupling






# Space-Time Coupling




# Space-Time Coupling






# Space-Time Coupling




# Space-Time Coupling

## Homogeneous Deformation






# Spectral Analysis to Estimate Fractal Dimension (Frequency Scaling)



- At largest scale (140km): Deformation is a red noise ( $1/f^2$ ) process.
- At smaller scales: becoming increasingly pink.

# Spectral Analysis to Estimate Fractal Dimension (Frequency Scaling)



Whitening rate is enhanced as L reduces

Whitening



Red Noise

- At largest scale (140km): Deformation is a red noise ( $1/f^2$ ) process.
- At smaller scales: becoming increasingly pink.



# Current Models

- Hurst exponent reflects the balance between external forcing (winds/currents) and internal energy dissipation by the ice pack.

$$\frac{\partial m \mathbf{U}}{\partial t} + \nabla \cdot (m \mathbf{U} \mathbf{U}) = \tau_a + \tau_w - m f \hat{\mathbf{k}} \times \mathbf{U} - m g \nabla H + \nabla \cdot \boldsymbol{\sigma}$$



Surface forcing

Body forcing

Dissipative term

Analogy to turbulence: In current models  $\text{Div}(\boldsymbol{\sigma})$  assumes homogenous deformation.



# Spectral Analysis to Estimate Fractal Dimension (Frequency Scaling)



- Rampal found a similar result.
- IABP buoy position error > SEDNA buoy position error
- Pierre: is your transition due to signal:noise?
- See Hutchings et al. (2012), erratum (2018), for error propagation.



# Spectral Analysis to Estimate Fractal Dimension (Frequency Scaling)



- No data resolving deformation at the scale of individual lead spacing.
- Higher precision GPS deployment in the field could resolve this question

# MOSAiC

- Distributed Network
- Scaling up concepts. Experiment built around up-scaling ocean-air fluxes.
- Sea ice team is highly collaborative. 8 onboard running many experiments.
- Hence limited opportunity to run labor intensive experiments
- Good opportunity: (i) second year / first year ice (new Arctic); (ii) full annual cycle






Conditions expected during MOSAiC deployment

# Projected drift tracks

## Drift Statistics

Start point: 85°N/105°E



Thomas Krumpen



# Where would you place the Central MOSAiC Camp? The remote sites?

MODIS

Sep 29, 2017

~79° N, 140° E

15 km

open water

new ice

multi-year ice

PS

Central  
Observatory

40 km

15 km

2 - 5 km

GPS Positions

Super-site: Atmo + Ice + Ocean  
(revisited regularly)  
Medium-site: Atmo + Ice + Ocean  
(revisited opportunistically / only selected sites)  
ML: lead site, variable position



# MOSAiC Distributed Network

new ice


multi-year ice



# High Fidelity Deformation Monitoring

- RADAR
- GNSS (partially covered by collaboration)
- Seismic
- SAR drift + deformation
- Laser Strain Array
- Attempting repeat worldview imagery
- Repeat aerial surveys.
- AUV not funded (no long-range underwater surveying)


# Tracking Seasonal Transitions



*[Toyota et, al 2006] Two regimes, with differing fractal scaling (gradient on graph) for floe size exist within the MIZ, and vary with distance from the ice edge. These may be attributed to different processes controlling floe size. Note that the observed data can also be fit to a function that emerges from competition between three stochastic processes controlling floe size (Herman 2010).*



# MOSAiC Summer Deformation



GNSS, accelerometers:


- high resolution horizontal and vertical motion
- Investigate modes of mechanical failure

Phil Hwang and collaborators

# SIDEx

- Chris Polashenski, Andy Mahoney, Jenny Hutchings + others.
- Small scale: floe cracking to floe-floe interaction.
- Mapping stress and strain fields.
- Identifying stress concentrators.
- Dissociating thermal and mechanical stresses.
- Identifying mechanisms and modes of failure
- 3 campaigns.





Compiled by Chris Polashenski



# 2019


- Landfast ice experiment in a lagoon
- Thermal cracking
- Related deformation
- and Stress field

Seismic, acoustic, high-resolution strain monitoring, point stress measurements, stress from linear elastic model.

# 2020

- Test equipment and methods at MOSAiC.

# 2021





# Remote Sensing

- OIB. ATM inappropriate for measuring ridges.
- Steriophotography and shadow methods
- Worldview (~50cm resolution satellite imagery, single band)
- SAR cover greatly improved
- New missions: Tandem-X, SWOT

# What's Missing?

- A few details in field campaigns still unfunded.
- A coordinated modeling project that utilizes field observations to build process understanding across ice aggregates. (Will recent MURI call fill this gap?)
- Coordinated effort to validate new redistribution and rheological models.
- Workshops/Collaborations to collate observational knowledge.
- Models that accurately simulate deformation and associated episodic ice motion.
- Models that can be applied to any scale and adjust to transitions in physical processes controlling deformation.



# Thoughts for developing rheological models ..... $\text{Div}(\sigma)$

- Differing mechanisms for sea ice deformation for different spatial scales, seasons and ice types.
- This could explain transitions in scaling behavior across different ice zones and seasons.
- Is a homogenous (elliptical) rheological model ever valid? Perhaps for large regions of the MIZ? Perhaps for scale between cracks (floe scale)? Within damage zones?



# Thoughts

- Observation evidence points to several physical mechanisms of sea ice deformation on various spatial and time scales.
- Perhaps variability in scaling behavior can be used to identify dynamically distinct sea ice zones.
- i.e. while one process may be scale invariant, it does not universally apply to the entire pack at all times.

# Thoughts

- To date we know that scaling laws vary with ice properties (strength), but do not have a clear understanding of how the dissipative processes in deformation change as the behaviour transitions seasonally and spatially.

# Summary

- Needs: Gaps in knowledge / Model validation
- MOSAiC
- SIDEX
- Remote Sensing / Airborne Campaigns
- What's Missing?

Let's continue this conversation over coffee/dinner/email.  
We need your insight to design field campaigns that support models.

Thanks: NSF, ONR, CMI (MMS), NASA, MOSAiC collaborators