

## Peter Herreshoff

Gunn High School Analysis H

## A Geometric Approach To Matrices Copyright © 2004 by Peter Herreshoff

Peter Herreshoff c/o Gunn High School 780 Arastradero Road Palo Alto, CA, 94306 USA

pherreshoff@pausd.org

Thanks to Josh Zucker who wrote the first draft of several chapters (particularly The Geometry of Complex Numbers, Vitamin i, Matrix Multiplication, and Inverses part I).

Also, thanks to my fellow teachers at Gunn High School Dave Deggeller and Danny Hahn for resurrecting the use of this pamphlet in their 11<sup>th</sup> grade Analysis classes and to their Analysis classes of 2013-14 who took on the project of retyping the text after the loss of the soft copy in a computer theft.

In addition, thanks to Gautam Mittal for editing the text for the 2017-18 academic year.

Above all, thanks to my student assistant Maya Sankar for her work on editing the text, standardizing the text in LaTeX, and incorporating an answer key.




Figure 1: A grid of posts with three strings.




Figure 2: A grid of posts with three strings.

## 1 It's a Snap

We begin with a problem that ties together ideas from geometry, complex numbers, matrices, combinatorics, and group theory. You likely studied geometry in 9<sup>th</sup> grade and complex numbers in 10<sup>th</sup> grade, so you should have a basis to start your investigation. The other three words may be unfamiliar at this point. You will be coming back to this problem in a series of investigations over the coming weeks.

Consider a rectangle of posts with n rows and 3 columns. An elastic string is anchored to one post in the topmost row and one post in the bottommost row. As the string descends top to bottom, it loops taut around one post in each row. Two other elastic strings are anchored and looped in the same way, with the condition that each post has exactly one string on it; no more, no less. An example where n=3 is depicted in Figure 1.

Figure 2 shows the six ways that two consecutive rows can be connected in this way — convince yourself these are the only six. I have labeled them with letters  $I,\,A,\,B,\,C,\,D,$  and E. We will call each of these six ways an *element* of our group. Each element has 6 posts (points) and 3 strings (segments).