

Critical points and approximation

Sketch a function

Sketch the graph of a function (any function you like, no need to specify a functional form) that is:¹

- Continuous on $[0, 3]$ and has the following properties: an absolute minimum at 0, an absolute maximum at 3, a local maximum at 1 and a local minimum at 2.
- Do the same for another function with the following properties: 2 is a **critical number** (i.e. $f'(x) = 0$ or $f'(x)$ is undefined), but there is no local minimum and no local maximum.

Find critical values

Find the critical values of these functions:²

- $f(x) = 5x^{3/2} - 4x$
- $s(t) = 3t^4 + 4t^3 - 6t^2$
- $f(r) = \frac{r}{r^2 + 1}$
- $h(x) = x \log(x)$

Find absolute minimum/maximum values

Find the absolute minimum and absolute maximum values of the functions on the given interval:³

- $f(x) = 3x^2 - 12x + 5, [0, 3]$
- $f(t) = t\sqrt{4 - t^2}, [-1, 4]$
- $s(x) = x - \ln(x), [1/2, 2]$
- $h(p) = 1 - e^{-p}, [0, 1000]$

A function with no local minima/maxima

Demonstrate that the function $f(x) = x^5 + x^3 + x + 1$ has no local maximum and no local minimum.⁴

Approximate root-finding

Show that the equation

$$x^7 - 6x + 4 = 0$$

¹Grimmer HW3.1

²Grimmer HW3.2

³Grimmer HW3.3

⁴Grimmer HW3.4

has a root between 0 and 1.⁵

- a. Find an initial approximation by ignoring the term x^7 .
- b. Use Newton's method to find the root correct to 3 decimal places.

Apply the mean value theorem

Does a continuous, differentiable function exist on $[0, 2]$ such that $f(0) = -1$, $f(2) = 4$, and $f'(x) \leq 2 \forall x$? Use the mean value theorem to explain your answer.⁶

⁵Pemberton and Rau 10.1.3

⁶Grimmer HW3.5