
at formulation
(extraction-period encoding, $x_{j,t}$)
by formulation
(cumulative encoding, $y_{j,t}$)
Validity penalty (one block j)
every pair of periods
consecutive periods only
$x_{j,1}$ $x_{j,2}$ $x_{j,3}$ $x_{j,4}$ $x_{j,T}$
$T(T-1)/2$ couplings per block
t
$y_{j,1}$ $y_{j,2}$ $y_{j,3}$ $y_{j,4}$ $y_{j,T}$
$T-1$ couplings per block
t
Precedence penalty (one relation $k \in \Lambda(j)$)
$x_{j,t}$ coupled to $x_{k,\tau}$ for all $\tau \le t$
$x_{k,1}$ $x_{k,2}$ $x_{k,3}$ $x_{k,4}$ $x_{k,T}$
$x_{j,1}$ $x_{j,2}$ $x_{j,3}$ $x_{j,4}$ $x_{j,T}$
$T(T+1)/2$ couplings per relation
$y_{j,t}$ coupled to $y_{k,t}$ only
$y_{k,1}$ $y_{k,2}$ $y_{k,3}$ $y_{k,4}$ $y_{k,T}$
$y_{j,1}$ $y_{j,2}$ $y_{j,3}$ $y_{j,4}$ $y_{j,T}$
T couplings per relation
block j (successor)
block $k \in \Lambda(j)$ (predecessor)
quadratic coupling