

The Generation of Rank 2 Drinfeld Modular Forms by Eisenstein Series: A Computational Approach

Liam Baker

February 16, 2021

Abstract

A computational method is presented for determining whether or not the space of Drinfeld modular forms for a principal congruence subgroup $\Gamma(N)$ is generated by Eisenstein series, which is currently an open question. This method boils down to expressing the dimension of the weight 2 space generated by Eisenstein series as the rank of a matrix of coefficients of products of Eisenstein series. As a result, the question is answered in the case of $q = 2$ and N quadratic or cubic and the case of $q = 3$ and N quadratic.

1 Introduction and Outline

Drinfeld modules are analogous to elliptic curves in the function field setting, and were first defined by [Drinfeld], who called them *elliptic modules* and used them to prove [Langlands]. Later, [Goss] defined what he called *Drinfeld modular forms* of rank 2 analogously with classical modular forms (in this paper, we will call Drinfeld modular forms simply modular forms, and explicitly mention when their classical counterparts are named). These functions form a graded algebra graded by their *weight*. Goss also imposed a condition of ‘boundedness at infinity’ which allowed him to show the finite-dimensionality of the spaces of D-modular forms of any given weight. [Gekeler] defined Eisenstein series as cases of modular forms, and showed that Eisenstein series of higher weight are generated by Eisenstein series of weight 1. The question then naturally arose of whether or not the space of all modular forms is generated as an algebra by the Eisenstein series of weight 1; this was answered in the affirmative by [Cornellissen] for linear principal congruence subgroups, but the question for general N is still open. The most significant progress thus far has been by [someone], who showed that the space of all modular forms is generated by the Eisenstein series of weight 1, possibly together with some cusp forms of weight 2 (these are modular forms which are zero at all the cusps). It thus suffices to establish whether or not the Eisenstein series of weight 1 generates the space of modular forms of weight 2, and in this paper we give a method for doing so for any specific characteristic q and principal congruence subgroup $\Gamma(N)$, as well as an implementation of this method in SageMath.

Let q be a prime power, and let \mathbb{F}_q denote the finite field of cardinality q .

2 Conclusion

Due to computational constraints we are only able to run the code for some small nonlinear values of N , but we hope that this progress will inspire others to improve on our method or to revisit this interesting question with fresh eyes.