INTRODUCTION

GENRES OF MUSIC:

i.CLASSICAL:

Classical music generally refers to the art music of the Western world, considered to be distinct from Western folk music or popular music traditions. It is sometimes distinguished as western classical music , as the term "classical music" can also be applied to non-western art music. Classical music is often characterized by formality and complexity in its musical forms and harmonic organization,particularly with the use of polyphony Since at least the ninth century it has been primarily a written tradition, spawning a sophisticated notational system, as well as accompanying literature in analytical, critical, historiographical, musicological and philosophical practices. A foundational component of Western culture, classical music is frequently seen from the perspective of individual or groups of composers, whose compositions, personalities and beliefs have fundamentally shaped its history.

ii. HIP-HOP:

Hip-hop or hip hop (formerly known as disco rap) is a genre of popular music that emerged in the early 1970s in New York City. The genre is characterized by stylized rhythmic sounds—often built around funk grooves, electronic drum beats, and rapping, a percussive vocal delivery of rhymed poetic speech as consciousness-raising expression. The music developed as part of the broader hip-hop culture; while often used to refer solely to rapping and rap music, "hip-hop" more properly denotes the practice(s) of the entire subculture.] The term *hip-hop music* is sometimes used synonymously with the term *rap music*, though rapping may not be the focus of hip-hop music. The genre also centers DJing, turntablism, scratching, beatboxing, and instrumental tracks.

iii. POP:

Pop music *is a genre of popular music that originated in its modern form during the mid-1950s in the United States and the United Kingdom. During the 1950s and 1960s, pop music encompassed rock and roll and the youth-oriented styles it influenced. Rock and pop music remained roughly synonymous until the late 1960s, after which pop became associated with music that was more commercial, ephemeral, and accessible.*

Identifying factors of pop music usually include repeated choruses and hooks, short to medium-length songs written in a basic format (often the verse–chorus structure), and rhythms or tempos that can be easily danced to. Much of pop music also borrows elements from other styles such as rock, hip hop, urban, dance, Latin, and country.

Iv. ROCK:

ROCK is a broad genre of popular music that originated in the United States as "rock and roll" in the late 1940s and early 1950s, developing into a range of different styles from the mid-1960s, primarily in the United States and the United Kingdom. It has its roots in rock and roll, a style that drew directly from the black musical genres of blues, rhythm and blues, and country music. Rock also drew strongly from genres such as electric blues and folk, and incorporated influences from jazz and other musical styles. For instrumentation, rock is typically centered on the electric guitar, usually as part of a rock group with electric bass guitar, drums, and one or more singers. Usually, rocKsong-based music with a 4 time signature and using a verse–chorus form, but the genre has become extremely diverse. Like pop music, lyrics often stress romantic love but also address a wide variety of other themes that are frequently social or political. Rock was the most popular genre of music in the U.S. and much of the Western world from the 1950s until its decline in the 2010s

V. COUNTRY:

Country (also called country and western) is a music genre originating in the southern regions of the United States, both the American South and the Southwest. First produced in the 1920s, country music is primarily focused on singing stories about working-class and blue-collar American life.

Country music is known for its ballads and dance tunes (i.e., "honky-tonk music") with simple form, folk lyrics, and harmonies generally accompanied by instruments such as banjos, fiddles, harmonicas, and many types of guitar (including acoustic, electric, steel, and resonator guitars). Though it is primarily rooted in various forms of American folk music, such as old-time music and Appalachian music, many other traditions, including Mexican, Irish, and Hawaiian music, have had a formative influence on the genre.Blues modes from blues music have been used extensively throughout its history as well.

VI. FOLK:

Folk music is a music genre that includes traditional folk music and the contemporary genre that evolved from the former during the 20th-century folk revival. Some types of folk music may be called world music. Traditional folk music has been defined in several ways: as music transmitted orally, music with unknown composers, music that is played on traditional instruments, music about cultural or national identity, music that changes between generations (folk process), music associated with a people's folklore, or music performed by custom over a long period of time. It has been contrasted with commercial and classical styles. The term originated in the 19th century, but folk music extends beyond that.

VII. MELODY:

A melody , also tune, voice, or line, is a linear succession of musical tones that the listener perceives as a single entity. In its most literal sense, a melody is a combination of pitch and rhythm, while more figuratively, the term can include other musical elements such as tonal color. It is the foreground to the background accompaniment. A line or part need not be a foreground melody.

Melodies often consist of one or more musical phrases or motifs, and are usually repeated throughout a composition in various forms. Melodies may also be described by

their melodic motion or the pitches or the intervals between pitches (predominantly conjunct or disjunct or with further restrictions), pitch range, tension and release, continuity and coherence, cadence, and shape**.**

VIII. REGGAE:

Reggae is a music genre that originated in Jamaica during the late 1960s. The term also denotes the modern popular music of Jamaica and its diaspora. A 1968 single by Toots and the Maytals, "Do the Reggay", was the first popular song to use the word *reggae*, effectively naming the genre and introducing it to a global audience.Reggae is rooted in traditional Jamaican Kumina, Pukkumina, Revival Zion, Nyabinghi, and burru drumming. Jamaican reggae music evolved out of the earlier genres mento, ska and rocksteady. Reggae usually relates news, social gossip, and political commentary. It is recognizable from the counterpoint between the bass and drum downbeat and the offbeat rhythm section. The immediate origins of reggae were in ska and rocksteady; from the latter, reggae took over the use of the bass as a percussion instrument.

RESEARCH METHODOLOGY

DATA COLLECTION:

Using questionnaires, primary data was gathered from students who are currently enrolled, and analysis was done using those data.

PRIMARY DATA COLLECTION:

Primary data collection is the process of gathering data directly from a first-hand source. In other words, it's data that's collected by the organization that expects to use it. Methods include surveys, interviews, observation, and focus groups. Primary sources provide an understanding that experts rely on, even as they support their conclusions with more extensive studies. The goal of primary data collection is to collect data that is as accurate and complete as possible. There are two types of primary data collection:

- *online*
- *offline*

There are three main types of primary data collection: qualitative, quantitative, and mixed mode. Each has its own advantages and disadvantages.

Qualitative primary data collection is the most important type for research because it allows you to collect rich information that can be used to improve your products or services. However, it's difficult to interpret and use this data in a way that's useful for business decisions.

Quantitative primary data collection is good for measuring how people are using your product or service, but it doesn't allow you to understand their thoughts and feelings very well.

Mixed mode primary data collection combines elements of both qualitative and quantitative methods; this makes it easier to get valuable insights without sacrificing accuracy.

Qualitative primary data collection

Qualitative primary data collection is a form of research that involves collecting data in an unstructured manner. This type of data collection can be helpful in gaining a more in-depth understanding of a topic or phenomenon. It can be useful when you want to explore an issue from many different perspectives. It can also be helpful when you want to gather information about people's opinions and feelings about a topic or phenomenon. Additionally, qualitative primary data collection can be helpful when you want to explore how people think about a particular issue or problem.

Qualitative primary data collection is a type of data collection that uses interviews, focus groups, and surveys to collect information from people. It's often used in research projects to gather feedback on products or services. This kind of data can be useful for understanding how people use products or how they think about problems. Qualitative primary data can also help you design better products or services.

Quantitative primary data collection

Quantitative primary data collection is the process of collecting data that can be measured. There are two types of quantitative primary data collection: online surveys and observation studies. Quantitative primary data collection has several advantages, including the ability to measure how people behave in natural settings, the ability to track changes over time, and the potential for large-scale studies. It has its own set of drawbacks, including low response rates and difficulty getting accurate results.

Quantitative primary data collection refers to any form of research where you collect information that can be measured (usually using a numeric scale). There are two main types – Online Surveys & Observational Studies – both with their own set of benefits and drawbacks. With Online Surveys in particular there's no need for respondents to leave their homes or face any awkward questions – they're able to take part from anywhere with an internet connection! But while response rates may be high due to this accessibility, it's often difficult getting accurate results as many people choose not answer surveys because they find them intrusive or inconvenient rather than ignorant/uninterested in the topic at hand! Lastly whilst observational studies have been around for a long time, they often suffer from low response rates as people are less likely to volunteer for them or feel comfortable about being observed.

SAMPLING METHODOLOGY

Methods of sampling from a population

It is normally impractical to study an entire population, such as when conducting a questionnaire survey. Sampling is a technique that allows researchers to infer information about a population based on results from a subset of the population rather than investigating every individual. Reducing the number of participants in a study lowers costs and workload while also making it easier to obtain high-quality data, but this must be balanced against having a large enough sample size with enough power to detect a true association.

If a sample is to be used, it is critical that the individuals chosen are representative of the entire population, regardless of the method used. This may entail specifically targeting difficult-to-reach groups.

There are several sampling techniques available, which can be divided into two categories: probability sampling and non-probability sampling. Probability (random) sampling begins with a complete sampling frame of all eligible individuals from which to draw your sample. This way, all eligible individuals have a chance of being chosen for the sample, and the results of your study will be more generalizable. Probability sampling methods are typically more time-consuming and costlier than non-probability sampling methods. Non-probability (non-random) sampling does not begin with a complete sampling frame, so some people have no chance of being chosen.

SIMPLE RANDOM SAMPLING

Each person is picked in this scenario purely at random, and everyone in the population has an equal likelihood of being chosen. Giving each person in a population a number and selecting which ones to include from a table of random numbers is one method of obtaining a random sample. 1 Use groups of three digits from the random number table to select your sample, for instance, if your sampling frame consists of 1000 people labelled 0 to 999. Select the person designated "94" if the first three numbers in the random number table were 094, and so on.

Simple random sampling lowers selection bias while allowing the sample error to be quantified, as is the case with all probability sampling techniques. The fact that it is the most user-friendly probability sampling technique is a distinct benefit. Simple random selection has the drawback of maybe not selecting enough people who have your feature of interest, especially if it is not common. Also, it could be challenging to set up a thorough sampling frame and difficult to get in touch with them, particularly if you need to use multiple communication methods (email, phone, and mail) and your sample units are dispersed throughout a large geographic area.

ANALYTICS METHODS

DESCRIPTIVE METHOD

Descriptive analytics is the process of identifying trends and relationships in current and historical data. It is sometimes referred to as the most basic form of data analysis because it describes trends and relationships but does not delve deeper.

Descriptive analytics is inexpensive and likely used on a daily basis in your organisation. Basic statistical software, such as Microsoft Excel, or data visualisation tools, such as Google Charts and Tableau, can aid in the parsing of data, the identification of trends and relationships between variables, and the visual display of information.

BAR DIAGRAM

A bar chart or bar graph is a chart or graph that presents categorical data with rectangular bars with heights or lengths proportional to the values that they represent. The bars can be plotted vertically or horizontally. A vertical bar chart is sometimes called a column chart.

A bar graph shows comparisons among discrete categories. One axis of the chart shows the specific categories being compared, and the other axis represents a measured value. Some bar graphs present bars clustered in groups of more than one, showing the values of more than one measured variable.

PIE CHART

A pie chart (or a circle chart) is a circular statistical graphic, which is divided into slices to illustrate numerical proportion. In a pie chart, the arc length of each slice (and consequently its central angle and area) is proportional to the quantity it represents. While it is named for its resemblance to a pie which has been sliced, there are variations on the way it can be presented. The earliest known pie chart is generally credited to William Playfair's Statistical Breviary of 1801.

Pie charts are very widely used in the business world and the mass media. However, they have been criticized, and many experts recommend avoiding them, as research has shown it is difficult to compare different sections of a given pie chart, or to compare data across different pie charts.

MULTIPLE BAR DIAGRAM

A multiple bar graph is nothing but a usual bar graph except for the fact that each category has two or more bars – one for each subdivision. For instance, if we plot the population of a country over a period of five years we can draw two bars for each year – one showing the number of females and the other bar showing the male population. The diagram below shows an example of a multiple bar graph.

INFERENCIAL METHOD

Non-parametric statistics rely on either a distribution that is free of parameters or a distribution that is predetermined.

Non-parametric tests are the experiments that don't need a sample population to make assumptions; they are what define non-parametric statistics. Due to the fact that non-parametric tests don't rely on information from any particular parametric collection of probability distributions, they are often referred to as distribution-free tests.

In other words, a statistical method known as non-parametric statistics is one in which a certain set of data is not required to fit into a normal distribution. Ordinal data, which doesn't rely on numbers but rather a ranking or order, is typically employed in non-parametric statistics. Tests, conclusions, models, and descriptive statistics all need to be considered.

So, a statistical method known as non-parametric statistics is one in which data are not obtained from a predetermined model that is based on a limited set of parameters. Non parametric statistics has a whole separate set of topics than the normal distribution model, factorial design, and regression modelling.

Non-parametric models are much simpler to use than parametric models, but they do not provide the same level of precision as other statistical models. In investigations where a net change in input has little to no impact on the output, non-parametric statistics is typically selected. like the outcomes are probably going to remain the same even if the numerical data changes.

CHI-SQUARE INDEPENDENCE OF ATTRIBUTES

A specific version of Pearson's chi-square test is the chi-square (X2) test of independence. Chi-square tests by Pearson are nonparametric assessments of categorical variables. They are employed to ascertain whether your data differ considerably from your expectations.

To ascertain if two categorical variables are connected, you can perform a chi-square test of independence, also referred to as a chi-square test of association. When two variables are connected, the likelihood that one will have a particular value depends on the value of the other variable.

The observed frequencies, or the number of observations in each combined group, are used in the chi-square test of independence calculations.

In order to determine if two variables are unrelated, the test compares observed frequencies to those you would anticipate. The observed and expected frequencies will be comparable if the variables are unrelated.

KRUSKAL WALLIS TEST

Kruskal-Wallis test is a nonparametric test (also called a distribution-free test), the data used do not have to be normally distributed, in contrast to analysis of variance. The only requirement is that the data be ordinal scale.

In the Kruskal-Wallis test, ordinal variables are sufficient, since non-parametric tests do not use the differences of the values, but the ranks (which value is larger, which is smaller). Therefore, the Kruskal-Wallis test is also often called "Kruskal-Wallis one-way ANOVA by ranks test".

Assumptions for the Kruskal-Wallis test

To compute a Kruskal-Wallis test, only several independent random samples with at least ordinally scaled characteristics must be available. The variables do not have to satisfy a distribution curve.

MANN-WHITNEY U -TEST

The Mann-Whitney U-Test can be used to test whether there is a difference between two samples (groups), and the data need not be normally distributed.

The Mann-Whitney U test is thus the non-parametric counterpart to the t-test for independent samples; it is subject to less stringent assumptions than the t-test. Therefore, the Mann-Whitney U test is always used when the requirement of normal distribution for the t-test is not met.

Assumptions Mann-Whitney U Tests

To be able to calculate a Mann-Whitney U test, only two independent random samples with at least ordinally scaled characteristics must be available. The variables do not have to satisfy any distribution curve.

PIE – Chart between the Stream

POP	30
HIP-HOP	16
CLASSICAL	9
ROCK	8
COUNTRY	2
FOLK	1
MELODY	1
REGGAE	1



Interpretation:

The Pie-Chart Graphical representation of the table shows that out of 68 people, 44.12% of people listen to pop music, 23.53% of people listen to hip-hop music , 13.24% of people listen to classical music,11.76% of people listen to rock music, 2.94% of people listen to country music, 1.47% of people listen to folk music, 1.47% of people listen to melody music, 1.47 of people listen to reggae music. So, we conclude that most people listen to pop music.

Bar Diagram between the streaming service

SPOTIFY	*43*
YOUTUBE MUSIC	*16*
AMAZON MUSIC	*2*
APPLE MUSIC	*2*
INSTAGRAM	*1*
JIO MUSIC	*1*
MASS TAMILIAN	*1*
OFFLINE MUSIC	*1*
WYNK	*1*



INTERPRETATION*:*

This bar chart shows which streaming service people often used. 43 people uses spotify to listen music, 16 people uses youtube music to listen music, 2 people uses amazon music and apple music to listen music, 1 people use instagram,masstamilian,offline music and wynk. So, we conclude that most people uses spotify to listen music.

Multiple Bar Diagram between

	CLASSICAL	COUNTRY	FOLK	HIP-HOP	MELODY	POP	REGGAE	ROCK
10-20	4	2	1	10	0	20	0	4
21-30	2	0	0	6	1	9	1	2
31-40	0	0	0	0	0	1	0	0
41-50	2	0	0	0	0	0	0	1
51-60	1	0	0	0	0	0	0	1
TOTAL	9	2	1	16	1	30	1	8



INTERPRETATION:

Multiple Bar Diagram of how much time they spend listening to music

	MALE	*FEMALE*
MORE THAN 1 HOUR	*16*	*6*
30 MINUTES	*10*	*8*
1 HOUR	*11*	*5*
LESS THAN 30 MINUTES	*11*	*1*

.



INTERPRETATION:

This histogram is about how many minutes does the people listen to music.22 people(6 female,16 male) listen more than 1 hour,18 people(8 female,10 male) listen exactly 30 minutes, 16 people(5 female, 11 male) listen exactly 1 hour and 12 people(1 female, 11 male) listen less than 30 minutes. So, we conclude that most of the people according to the data listen music for more than 1 hour.

CHI – SQUARE TEST FOR INDEPENDENCE OF ATTRIBUTES

The following data are for a sample of 69 people who were classified with respect to gender and the streaming service they listen to music. Test whether there is any similarity between these two variables.

	CLASSICAL	COUNTRY	FOLK	HIP-HOP	MELODY	POP	REGGAE	ROCK	GRAND TOTAL
MALE	5	1		4		7		3	20
FEMALE	4	1	1	12	1	23	1	5	48

Procedure:

Null Hypothesis

H0: There is no significance difference between two variables.

Alternative Hypothesis

H1: There is significance difference between two variables.

Level of Significance =5% ~ 0.05

Test Statistic

Chi square=$\sum[(O-E)^2/E] \sim (r-1)(c-1)$

where,

r = No. of Rows,c = No. of Columns

The expected frequency E = $R \times C/N$

R = Each Row Total

C = Each Column Total

N = Grand Total

CALCULATION:

Observed value(O)	Expected value(E)	(O-E)	(O-E)^2	[(O-E)^2]/E
5	2.6471	2.3529	5.5361	2.0914
1	0.5882	0.4118	0.1696	0.2883
4	4.7059	-0.7059	0.4983	0.1059
7	8.8235	-1.8235	3.3252	0.3769
3	2.3529	0.6471	0.4188	0.1779
4	6.3529	-2.3529	5.5361	0.8714
1	1.4117	-0.4117	0.1695	0.1201
1	0.7059	0.2941	0.0865	0.1225
12	11.2941	0.7059	0.4983	0.0441
1	0.7059	0.2941	0.0865	0.1225
23	21.1765	1.8235	3.3252	0.157
1	0.7059	0.2941	0.0865	0.1225
5	5.6471	-0.6471	0.4187	0.0741
0	0.2941	-0.2941	0.0865	0.2941
0	0.2941	-0.2941	0.0865	0.2941
0	0.2941	-0.2941	0.0865	0.2941
			Chi square value	5.5569

Table Value:

Use 5% Level of significance at (2-1) (8-1), 7 d.f is 14.067

Inference:

Here, Calculation value (5.5569) is less than Table value (14.067) so, we reject the Alternative Hypothesis H1 and we accept the Null Hypothesis H0. Hence, we conclude There is no significance difference between two variables.

Test – 1 ***Using R-Software***

Hypothesis Setting

Null Hypothesis

H0 :There is no significance difference between two variables.

Alternative Hypothesis

H1:There is significance difference between two variables

. Level of Significance 5%

Data1 <- matrix(c(5,1,0,4,0,7,0,3,4,1,1,12,1,23,1,5)) nrow = 2, ncol = 8, byrow = TRUE , dimnames =list(c("Male", "Female"), c("CLASSICAL", "COUNTRY", "FOLK", "HIP-HOP","MELODY","POP","REGGAE","ROCK")))

Data

print(Data1)

	CLASSICAL	COUNTRY	FOLK	HIP-HOP	MELODY	POP	REGGAE	ROCK
MALE	5	1	0	4	0	7	0	3
FEMALE	4	1	1	12	1	23	1	5

Chi-Square Test for above data

chisq.test(Data1)

Pearson's Chi-squared test

data: Data1

X-squared =5.5569, df = 7, p-value = 0.2545

Table Value for (r-1)(c-1) d.f. LOS 5% qchisq(.95, 7)

[1] 14.067

Inference

Here, Calculation value (5.5569) is less than Table value (14.067) so, we reject the Alternative Hypothesis H1 and we accept the Null Hypothesis H0. Hence, we conclude There is no significance difference between two variables.

KRUSKAL WALLIS H TEST

The following data are for a sample of 68 people who were classified with pop,hip-hop,classical. Test whether there is any similarity between these three variables.

POP	*HIP-HOP*	*CLASSICAL*
20	*10*	*2*
9	*6*	*2*
1	*0*	*0*
0	*0*	*1*
0	*0*	*1*

Procedure

Null Hypothesis

H0 : There is no significance difference between the given variables.

Alternative Hypothesis

H1: There is significance difference between the given variables.

Level of Significance α*=5% ~ 0.05*

Test Statistic

$H=12\ N(N+1)\sum Ti\ 2\ Ni\ -3(N+1)\sim\ (K-1)$

Where,

K = No. of groups used for comparison

N = Total size of the sample

ni = ith group's sample size

Ri = Total of the ranks related to i-th group

CALCULATION*:*

	15	14	10.5
	13	12	10.5
	8	3.5	3.5
	3.5	3.5	8
	3.5	3.5	8
R	43	36.5	40.5
n	5	5	5
R^2/n	369.8	266.45	328.05
N	15		
k	3		

H= 15 /15(15+1)∑28.5^2/4 +24^2/4 +25.5^2/4 −3(15+1)

H=1.2556

Table Value

Use 5% Level of significance at (K-1) (3-1), 2 d.f is 5.9914

Inference

Here, Calculation value (1.2556) is less than Table value (5.9914) so, we reject the Alternative Hypothesis H1 and we accept the Null Hypothesis H0. Hence, we conclude There is no significance difference between two variables.

Kruskal-Wallis Test

Hypothesis

Null Hypothesis

H0: There is no significance difference between the given variables.

Alternative Hypothesis

H1: There is significance difference between the given variables.

Level of Significance =5%

```
## Data
x <- c(20,9,1,0,0)
 y <- c(10,6,0,0,0)
z <- c(2,2,0,1,1)
data4 <- c(x,y,z)
## Kruskal - wallis Test
g <- factor(rep(1:3, c(5,5,5)), labels = c("POP", "HIP-HOP", "CLASSICAL"))
kruskal.test(data4, g)
##
## Kruskal-Wallis rank sum test
##
 ## data: data4 and g
## Kruskal-Wallis chi-squared =1.2556, df = 2, p-value = 0.1856
## Table value
qchisq(.95, 2)
 ## [1] 5.991465
```

Inference

Here, Calculation value (1.2556) is less than Table value (5.9914) so, we reject the Alternative Hypothesis H1 and we accept the Null Hypothesis H0. Hence, we conclude There is no significance difference between two variables.

MANN – WHITNEY U TEST

From the following sample, test the hypothesis given below. Test whether the people who listens music in spotify is more than the people who listens music in youtube music

SPOTIFY	YOUTUBE MUSIC
29	6
12	4
1	0
0	2
0	2

Procedure

Null Hypothesis

H0 : There is no significance difference.

Alternative Hypothesis

H1: There is significance difference.

Level of Significance α=5% ~ 0.05

Test Statistic

$U=n1n2(n2(n2+1)/2)\sum Ri$

where, U = Mann-Whitney U test n1 = Sample size one n2 = Sample size two Ri = Rank total of the sample size

CALCULATION:

SPOTIFY	RANK	YOUTUBE MUSIC	RANK
29	10	6	8
12	9	4	7
1	4	0	2
0	2	2	5.5
0	2	2	5.5
	27		28

n1 = 5 ; n2 = 5

R1 = 27 R2 = 28

U1= 27−5(5+1)/2=12 *U2= 28−5(5+1)/2=13*

U=min(U1 ,U2)
U=MIN(12,13)
U=12

Table Value
Use 5% Level of significance at n1, n2 d.f is 4
Inference
Here, Calculation Value (12) is greater than Table Value (4) so, we accept the Alternative Hypothesis H1 and reject the Null hypothesis H0. Hence, we concluded that the currently pursuing students are having the Knowledge about their course.

Mann-Whitney U -Test

Hypothesis

Null Hypothesis

H0: There is no significance difference

Alternative Hypothesis

H1: There is significance difference

Level of Significance alpha = 5%

spotify <- c(29,12,1,0,0) youtube music <- c(6,4,0,2,2)

Data

matrix(c(spotify,youtube music), nrow = 2, ncol = 5, byrow = TRUE, dimname= list(c ("Neutral", "Agree")))

spotify 29 12,1 0 0

youtube music 6 4 0 2 2

Mann-Whitney U - Test

wilcoxin.test(spotify, youtube music)

Wilcoxon rank sum test with continuity correction

##

data: spotify and youtube music

W = 12, p-value = 0.5614 ## alternative hypothesis: true location shift is not equal to 0

Inference

Here, Calculation Value (12) is greater than Table Value (4) so, we accept the Alternative Hypothesis H1 and reject the Null hypothesis H0. Hence, we concluded that the currently pursuing students are having Knowledge about their course.

CONCLUSION:

Our Project "Types of music enjoyed by the people", We have analyzed the interest of the people are

Taking which type of Music.

- *The people prefer to listen the pop music mostly*
- *people like to listen in Spotify app mostly*
- *According to our Data most people listen music more than 1 hour .*
- *Both Male & Female listeners prefer pop music compared to other music genres.*
- *Music is mostly listened at a certain age group (10-20).*