1-1.9-8

EE24BTECH11010 - BALAJI B

Question:

The distance between the points $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} a - b \\ a + b \end{pmatrix}$ is

Answer:

Variable	Value	Description
P	$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$	First Endpoint
Q	$\begin{pmatrix} a-b\\a+b \end{pmatrix}$	Second Endpoint

TABLE 0: Input parameters

$$\therefore \mathbf{P} - \mathbf{Q} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} - \begin{pmatrix} a - b \\ a + b \end{pmatrix} = \begin{pmatrix} b - a \\ -a - b \end{pmatrix} \tag{0.1}$$

1

$$(\mathbf{P} - \mathbf{Q})^{\top} (\mathbf{P} - \mathbf{Q}) = \begin{pmatrix} b - a & -a - b \end{pmatrix} \begin{pmatrix} b - a \\ -a - b \end{pmatrix}$$
(0.2)

$$(\mathbf{P} - \mathbf{Q})^{\mathsf{T}} (\mathbf{P} - \mathbf{Q}) = (b - a)^2 + (a + b)^2$$
 (0.3)

Thus the desired distance is

$$d = ||P - Q|| = \sqrt{(P - Q)^{\top}(P - Q)}$$
 (0.4)

$$d = ||P - Q|| = \sqrt{(b - a)^2 + (a + b)^2}$$
(0.5)

Assuming the points a = 2 and b = 3

$$\mathbf{P} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \text{ and } \mathbf{Q} = \begin{pmatrix} -1 \\ 5 \end{pmatrix} \tag{0.6}$$

$$d = ||P - Q|| = \sqrt{(3-2)^2 + (2+3)^2}$$
 (0.7)

$$d = \sqrt{26} \tag{0.8}$$




Fig. 0.1: Plot of the line passing through P and Q