

- [150] M. Girod, T.N.T. Phan, L. Charles, Tuning block copolymer structural information by adjusting salt concentration in liquid chromatography at critical conditions coupled with electrospray tandem mass spectrometry, *Rapid Commun. Mass Spectrom.* 23 (10) (2009) 1476–1482.
- [151] J. Han, B.H. Jeon, C.Y. Ryu, J.J. Semler, Y.K. Jhon, J. Genzer, Discriminating among co-monomer sequence distributions in random copolymers using interaction chromatography, *Macromol. Rapid Commun.* 30 (18) (2009) 1543–1548.
- [152] W. Hiller, P. Sinha, H. Pasch, Online HPLC-NMR of PS-b-PMMA and blends of PS and PMMA, 2-LCCC-NMR at critical conditions of PMMA, *Macromol. Chem. Phys.* 210 (8) (2009) 605–613.
- [153] K. Im, H.-w. Park, S. Lee, T. Chang, Two-dimensional liquid chromatography analysis of synthetic polymers using fast size exclusion chromatography at high column temperature, *J. Chromatogr. A* 1216 (21) (2009) 4606–4610.
- [154] S. Julka, H. Cortes, R. Harfmann, B. Bell, A. Schweizer-Theobaldt, M. Pursch, L. Mondello, S. Maynard, D. West, Quantitative characterization of solid epoxy resins using comprehensive two dimensional liquid chromatography coupled with electrospray ionization-time of flight mass spectrometry, *Anal. Chem.* 81 (11) (2009) 4271–4279.
- [155] M.I. Malik, H. Ahmed, B. Trathnigg, Liquid chromatography under critical conditions: Practical applications in the analysis of amphiphilic polymers, *Anal. Bioanal. Chem.* 393 (6–7) (2009) 1797–1804.
- [156] M.I. Malik, B. Trathnigg, R. Saf, Characterization of ethylene oxide-propylene oxide block copolymers by combination of different chromatographic techniques and matrix-assisted laser desorption ionization time-of-flight mass spectroscopy, *J. Chromatogr. A* 1216 (38) (2009) 6627–6635.
- [157] M.I. Malik, B. Trathnigg, C.O. Kappe, Amphiphilic polymers based on higher alkylene oxides, *J. Chromatogr. A* 1216 (7) (2009) 1167–1173.
- [158] M.I. Malik, B. Trathnigg, C. Oliver Kappe, Microwave-assisted polymerization of higher alkylene oxides, *Eur. Polym. J.* 45 (3) (2009) 899–910.
- [159] M.I. Malik, B. Trathnigg, Full separation of oligomers in block copolymers of ethylene oxide and propylene oxide, *J. Sep. Sci.* 32 (11) (2009) 1771–1781.
- [160] A. Takano, Y. Kushida, Y. Ohta, K. Masuoka, Y. Matsushita, The second virial coefficients of highly-purified ring polystyrenes in cyclohexane, *Polymer* 50 (5) (2009) 1300–1303.
- [161] S. Abrar, B. Trathnigg, Characterization of polyoxyethylenes according to the number of hydroxy end groups by hydrophilic interaction chromatography at critical conditions for polyethylene glycol, *Anal. Bioanal. Chem.* 400 (8) (2010) 2531–2537.
- [162] H. Ahmed, B. Trathnigg, C.O. Kappe, R. Saf, Synthesis of poly(ϵ -caprolactone) diols and EO-CL block copolymers and their characterization by liquid chromatography and MALDI-TOF-MS, *Eur. Polym. J.* 46 (3) (2010) 494–505.
- [163] H. Ahmed, B. Trathnigg, Characterization of PEG in monomethyl ethers of PEG by 2-D LC, *J. Sep. Sci.* 33 (10) (2010) 1448–1453.
- [164] M. Al Samman, W. Radke, A. Khalyavina, A. Lederer, Retention behavior of linear, branched, and hyperbranched polyesters in interaction liquid chromatography, *Macromol.* 43 (7) (2010) 3215–3220.
- [165] W.K. Elhrari, P.E. Mallon, Two-dimensional chromatographic analysis of polystyrene-block-poly(methyl methacrylate) copolymers synthesized by selective oxidation of polystyrene-9-borabicyclo[3.3.1]nonane, *Macromol. Symp.* 298 (1) (2010) 174–181.
- [166] N. Fandrich, J. Falkenhagen, S.M. Weidner, B. Staal, A.F. Thünemann, A. Laschewsky, Characterization of new amphiphilic block copolymers of N-vinylpyrrolidone and vinyl acetate, 2-chromatographic separation and analysis by MALDI-TOF and FT-IR coupling, *Macromol. Chem. Phys.* 211 (15) (2010) 1678–1688.
- [167] W. Hiller, H. Pasch, P. Sinha, T. Wagner, J. Thiel, M. Wagner, K. Müllen, Coupling of NMR and liquid chromatography at critical conditions: A new tool for the block length and microstructure analysis of block copolymers, *Macromol.* 43 (11) (2010) 4853–4863.
- [168] M.I. Malik, B. Trathnigg, K. Bartl, R. Saf, Characterization of polyoxyalkylene block copolymers by combination of different chromatographic techniques and MALDI-TOF-MS, *Anal. Chim. Acta* 658 (2) (2010) 217–224.
- [169] C. Schmid, J. Falkenhagen, C. Barner-Kowollik, An efficient avenue to poly(styrene)-block-poly(ϵ -caprolactone) polymers via switching from RAFT to hydroxyl functionality: Synthesis and characterization, *J. Polym. Sci. Part A: Polym. Chem.* 49 (1) (2010) 1–10.
- [170] P. Sinha, W. Hiller, H. Pasch, Characterisation of blends of polyisoprene and polystyrene by on-line hyphenation of HPLC and 1H-NMR: LC-CC-NMR at critical conditions of both homopolymers, *J. Sep. Sci.* 33 (22) (2010) 3494–3500.
- [171] B. Trathnigg, M.I. Malik, N. Pircher, S. Hayden, Liquid chromatography at critical conditions in ternary mobile phases: Gradient elution along the critical line, *J. Sep. Sci.* 33 (14) (2010) 2052–2059.
- [172] B. Trathnigg, H. Ahmed, Separation of all oligomers in polyethylene glycols and their monomethyl ethers by one-dimensional liquid chromatography, *Anal. Bioanal. Chem.* 399 (4) (2010) 1535–1545.
- [173] B. Trathnigg, S. Abrar, Characterization of complex copolymers by two-dimensional liquid chromatography, *Procedia Chem.* 2 (1) (2010) 130–139.
- [174] M. van Hulst, A. van der Horst, W.T. Kok, P.J. Schoenmakers, Comprehensive 2-D chromatography of random and block methacrylate copolymers, *J. Sep. Sci.* 33 (10) (2010) 1414–1420.
- [175] E.H.H. Wong, M.H. Stenzel, T. Junkers, C. Barner-Kowollik, Spin capturing with ‘clickable’ nitrones: Generation of miktoarmed star polymers, *Macromol.* 43 (8) (2010) 3785–3793.
- [176] A. Baumgaertel, C. Weber, N. Fritz, G. Festag, E. Altuntaş, K. Kempe, R. Hoogenboom, U.S. Schubert, Characterization of poly(2-oxazoline) homo- and copolymers by liquid chromatography at critical conditions, *J. Chromatogr. A* 1218 (46) (2011) 8370–8378.
- [177] W. Hiller, P. Sinha, M. Hehn, H. Pasch, T. Hofe, Separation analysis of polyisoprenes regarding microstructure by online LCCC-NMR and SEC-NMR, *Macromol.* 44 (6) (2011) 1311–1318.
- [178] A.J. Inglis, C. Barner-Kowollik, Visualizing the efficiency of rapid modular block copolymer construction, *Polym. Chem.* 2 (1) (2011) 126–136.
- [179] M.I. Malik, P. Sinha, G.M. Bayley, P.E. Mallon, H. Pasch, Characterization of polydimethylsiloxane-block-polystyrene (PDMS-b-PS) copolymers by liquid chromatography at critical conditions, *Macromol. Chem. Phys.* 212 (12) (2011) 1221–1228.
- [180] C. Schmid, S. Weidner, J. Falkenhagen, C. Barner-Kowollik, In-depth LCCC-(GELC)-SEC characterization of ABA block copolymers generated by a mechanistic switch from RAFT to ROP, *Macromol.* 45 (1) (2011) 87–99.
- [181] A. Takano, Y. Ohta, K. Masuoka, K. Matsubara, T. Nakano, A. Hieno, M. Itakura, K. Takahashi, S. Kinugasa, D. Kawaguchi, Y. Takahashi, Y. Matsushita, Radii of gyration of ring-shaped polystyrenes with high purity in dilute solutions, *Macromol.* 45 (1) (2011) 369–373.
- [182] H. Barqawi, E. Ostas, B. Liu, J.-F. Carpentier, W.H. Binder, Multidimensional characterization of α,ω -telechelic poly(ϵ -caprolactone)s via online coupling of 2D chromatographic methods (LC/SEC) and ESI-TOF/MALDI-TOF-MS, *Macromol.* 45 (24) (2012) 9779–9790.
- [183] M.I. Malik, G.W. Harding, M.E. Grabowsky, H. Pasch, Two-dimensional liquid chromatography of polystyrene-polyethylene oxide block copolymers, *J. Chromatogr. A* 1244 (2012) 77–87.
- [184] M.I. Malik, G.W. Harding, H. Pasch, Two-dimensional liquid chromatography of PDMS-PS block copolymers, *Anal. Bioanal. Chem.* 403 (2) (2012) 601–611.
- [185] Y. Ohta, M. Nakamura, Y. Matsushita, A. Takano, Synthesis, separation and characterization of knotted ring polymers, *Polymer* 53 (2) (2012) 466–470.
- [186] M. Rollet, D. Glé, T.N.T. Phan, Y. Guilleaume, D. Bertin, D. Gimes, Characterization of functional poly(ethylene oxide)s and their corresponding polystyrene block copolymers by liquid chromatography under critical conditions in organic solvents, *Macromol.* 45 (17) (2012) 7171–7178.
- [187] C. Schmid, J. Falkenhagen, T.F. Beskers, L.-T.T. Nguyen, M. Wilhelm, F.E. Du Prez, C. Barner-Kowollik, Multi-block polyurethanes via RAFT end-group switching and their characterization by advanced hyphenated techniques, *Macromol.* 45 (16) (2012) 6353–6362.
- [188] P. Sinha, G.W. Harding, K. Maiko, W. Hiller, H. Pasch, Comprehensive two-dimensional liquid chromatography for the separation of protonated and deuterated polystyrene, *J. Chromatogr. A* 1265 (2012) 95–104.
- [189] P. Sinha, W. Hiller, V. Bellas, H. Pasch, Analysis of polystyrene-b-polyisoprene copolymers by coupling of liquid chromatography at critical conditions to NMR at critical conditions of polystyrene and polyisoprene, *J. Sep. Sci.* 35 (14) (2012) 1731–1740.
- [190] P. Sinha, M. Grabowsky, M.I. Malik, G.W. Harding, H. Pasch, Characterization of polystyrene-block-polyethylene oxide diblock copolymers and blends of homopolymers by liquid chromatography at critical conditions (LCCC), *Macromol. Symp.* 313–314 (1) (2012) 162–169.
- [191] S.M. Weidner, J. Falkenhagen, I. Bressler, Copolymer composition determined by LC-MALDI-TOF MS coupling and ‘MassChrom2D’ data analysis, *Macromol. Chem. Phys.* 213 (22) (2012) 2404–2411.
- [192] A.-M. Zorn, C. Barner-Kowollik, Transformation of macromonomers into ring-opening polymerization macroinitiators: A detailed initiation efficiency study, *J. Polym. Sci. Part A: Polym. Chem.* 50 (12) (2012) 2366–2377.
- [193] H. Barqawi, M. Schulz, A. Olubummo, V. Saurland, W.H. Binder, 2D-LC/SEC-(MALDI-TOF)-MS characterization of symmetric and nonsymmetric biocompatible PEOm-PIB-PEOn block copolymers, *Macromol.* 46 (19) (2013) 7638–7649.
- [194] Y. Doi, Y. Ohta, M. Nakamura, A. Takano, Y. Takahashi, Y. Matsushita, Precise synthesis and characterization of tadpole-shaped polystyrenes with high purity, *Macromol.* 46 (3) (2013) 1075–1081.
- [195] D.J. Keddie, C. Guerrero-Sanchez, G. Moad, The reactivity of N-vinylcarbazole in RAFT polymerization: trithiocarbonates deliver optimal control for the synthesis of homopolymers and block copolymers, *Polym. Chem.* 4 (12) (2013) 3591.
- [196] M.I. Malik, H. Ahmed, B. Trathnigg, Separation of telechelic oligomers according to architecture by liquid chromatography, *J. Chromatogr. A* 1314 (2013) 180–187.
- [197] D. Mekap, T. Macko, R. Brüll, R. Cong, A. deGroot, A. Parrott, P. Cools, W. Yau, Liquid chromatography at critical conditions of polyethylene, *Polymer* 54 (21) (2013) 5518–5524.