

- [198] M. Mlynek, W. Radke, Critical chromatography in ternary solvents, *J. Chromatogr. A* 1284 (2013) 112–117.
- [199] A.S. Kasegaonkar, H. Barqawi, W.H. Binder, Synthesis and crystallization of star-shaped photocleavable poly(ϵ -caprolactone)s, *J. Polym. Sci. Part A: Polym. Chem.* 53 (5) (2014) 642–649.
- [200] M.I. Malik, T. Mahboob, S. Ahmed, Characterization of poly(2-vinylpyridine)-block-poly(methyl methacrylate) copolymers and blends of their homopolymers by liquid chromatography at critical conditions, *Anal. Bioanal. Chem.* 406 (25) (2014) 6311–6317.
- [201] M.P.F. Pepels, P. Soulié, R. Peters, R. Duchateau, Theoretical and experimental approach to accurately predict the complex molecular weight distribution in the polymerization of stainless cyclic esters, *Macromol.* 47 (16) (2014) 5542–5550.
- [202] P. Sinha, W. Hiller, H. Pasch, HPLC-1H-NMR characterization of polystyrene-block-polyisoprene copolymers: LCCC-1H-NMR using a single mobile phase, *Macromol. Symp.* 337 (1) (2014) 44–50.
- [203] S. Banerjee, P.N. Shah, Y. Jeong, T. Chang, K. Seethamraju, R. Faust, Structural characterization of telechelic polyisobutylene diol, *J. Chromatogr. A* 1376 (2015) 98–104.
- [204] J.L. Fischer, C.A. Lutomski, T.J. El-Baba, B.N. Siriwardena-Mahanama, S.M. Weidner, J. Falkenhagen, M.J. Allen, S. Trimpin, Matrix-assisted ionization-ion mobility spectrometry-mass spectrometry: Selective analysis of a europium-PEG complex in a crude mixture, *J. Am. Soc. Mass Spectrom.* 26 (12) (2015) 2086–2095.
- [205] M. Hehn, P. Sinha, H. Pasch, W. Hiller, Onflow liquid chromatography at critical conditions coupled to 1H and 2H nuclear magnetic resonance as powerful tools for the separation of poly(methylmethacrylate) according to isotopic composition, *J. Chromatogr. A* 1387 (2015) 69–74.
- [206] T.G. Wright, H. Pasch, A simple route to deuterated polystyrene block copolymers by reverse iodine transfer polymerisation, *Polym. Chem.* 6 (17) (2015) 3236–3244.
- [207] C. Appiah, K.R. Siefertmann, M. Jorewitz, H. Barqawi, W.H. Binder, Synthesis and characterization of new photoswitchable azobenzene-containing poly(ϵ -caprolactones), *RSC Adv.* 6 (8) (2016) 6358–6367.
- [208] S. Bhati, T. Macko, R. Brüll, Estimation of critical conditions of polymers based on monitoring the polymer recovery, *J. Chromatogr. A* 1451 (2016) 91–96.
- [209] L. Gao, J. Oh, T. Chang, D. Chen, X. Li, X. Yang, Y. Tu, X. Zhu, C.Y. Li, A nearly quantitative synthetic approach towards monocyclic polystyrenes and the solvent, concentration and molecular weight effect on cyclic yield, *Polymer* 101 (2016) 379–387.
- [210] M.I. Malik, S. Lee, T. Chang, Comprehensive two-dimensional liquid chromatographic analysis of poloxamers, *J. Chromatogr. A* 1442 (2016) 33–41.
- [211] M.I. Malik, M. Irfan, A. Khan, S. Rahim, R. Abdul-Karim, J. Hashim, Alkylene oxide polymerizations: identification of side reactions and by-products, *J. Polym. Res.* 23 (12) (2016).
- [212] Y.-Z. Wei, R.-X. Zhuo, X.-L. Jiang, Separation of polyethylene glycols and amino-terminated polyethylene glycols by high-performance liquid chromatography under near critical conditions, *J. Chromatogr. A* 1447 (2016) 122–128.
- [213] Y.-Z. Wei, Y.-F. Chu, E. Uliyanchenko, P.J. Schoenmakers, R.-X. Zhuo, X.-L. Jiang, Separation and characterization of benzaldehyde-functional polyethylene glycols by liquid chromatography under critical conditions, *Polym. Chem.* 7 (48) (2016) 7506–7513.
- [214] R. Abdul-Karim, S.G. Musharraf, M.I. Malik, Synthesis and characterization of novel biodegradable di- and tri-block copolymers based on ethylene carbonate polymer as hydrophobic segment, *J. Polym. Sci. Part A: Polym. Chem.* 55 (11) (2017) 1887–1893.
- [215] J. Ahn, T. Chang, X. Wang, Z. Limpouchová, K. Procházka, Influence of the chain architecture and the presence of end-groups or branching units chemically different from repeating structural units on the critical adsorption point in liquid chromatography, *Macromol.* 50 (21) (2017) 8720–8730.
- [216] N. Apel, E. Uliyanchenko, S. Moyses, S. Rommens, C. Wold, T. Macko, K. Rode, R. Brüll, Selective chromatographic separation of polycarbonate according to hydroxyl end-groups using a porous graphitic carbon column, *J. Chromatogr. A* 1488 (2017) 77–84.
- [217] Y. Jeong, Y. Jin, T. Chang, F. Uhlik, J. Roovers, Intrinsic viscosity of cyclic polystyrene, *Macromol.* 50 (19) (2017) 7770–7776.
- [218] S. Lee, H. Lee, T. Chang, A. Hirao, Synthesis and characterization of an exact polystyrene-graft-polyisoprene: A failure of size exclusion chromatography analysis, *Macromol.* 50 (7) (2017) 2768–2776.
- [219] M.K. Tufail, R. Abdul-Karim, S. Rahim, S.G. Musharraf, M.I. Malik, Analysis of individual block length of amphiphilic di- and tri-block copolymers containing poly(ethylene oxide) and poly(methyl methacrylate), *RSC Adv.* 7 (66) (2017) 41693–41704.
- [220] S. Abrar, B. Trathnigg, S. Javed, S. Kiran, T. Gulzar, Characterization of tween® surfactants by MALDI TOF-MS and high performance liquid chromatography in a ternary mobile phase, *Tenside Surfactants Deterg.* 55 (6) (2018) 447–454.
- [221] L. Gao, J. Oh, Y. Tu, T. Chang, Preparation of low molecular weight cyclic polystyrenes with high purity via liquid chromatography at the critical condition, *Polymer* 135 (2018) 279–284.
- [222] S. Lee, H. Choi, T. Chang, B. Staal, Two-dimensional liquid chromatography analysis of polystyrene/polybutadiene block copolymers, *Anal. Chem.* 90 (10) (2018) 6259–6266.
- [223] R. Abdul-Karim, S.G. Musharraf, M.I. Malik, Poly(propylene ether carbonate)-based di- and tri-block copolymers: Synthesis and chromatographic characterization, *Macromol. Res.* 27 (9) (2019) 911–918.
- [224] R. Khatoun, S. Rahim, R. Abdul-Karim, S.G. Musharraf, M.I. Malik, Characterization of polystyrene-block-poly(2-vinyl pyridine) copolymers and blends of their homopolymers by liquid chromatography at critical conditions, *Macromol.* 52 (20) (2019) 7688–7695.
- [225] M.I. Malik, Critical parameters of liquid chromatography at critical conditions in context of poloxamers: Pore diameter, mobile phase composition, temperature and gradients, *J. Chromatogr. A* 1609 (2020) 460440.
- [226] T. Ahmad, S.S. Durrani, F.J. Iftikhar, S.G. Musharraf, M.I. Malik, Concurrent ring-opening and atom transfer radical polymerization for synthesis of block copolymers, and their comprehensive chromatographic characterization, *Eur. Polym. J.* 142 (2021) 110161.
- [227] K. Molnar, C.A. Helfer, G. Kaszas, E. Krisch, D. Chen, G.B. McKenna, J.A. Kornfield, J.E. Puskas, Liquid chromatography at critical conditions (LCCC): Capabilities and limitations for polymer analysis, *J. Mol. Liq.* 322 (2021) 114956.
- [228] S.K. Panda, T.A. Al-Qunaysi, M. Al Taha, D.J. Kearney, Size exclusion chromatography of petroleum samples using amino-bonded silica phase, *Fuel* 302 (2021) 120914.
- [229] J. Yang, L. Chen, M. Zhu, M.W. Ishaq, S. Chen, L. Li, Investigation of the multimer cyclization effect during click step-growth polymerization of AB-type macromonomers, *Macromol.* 55 (15) (2022) 6830–6840.
- [230] B. Zdovc, H. Li, J. Zhao, D. Pahovnik, E. Žagar, Influence of microstructure on the elution behavior of gradient copolymers in different modes of liquid interaction chromatography, *Anal. Chem.* 94 (22) (2022) 7844–7852.
- [231] S. Deshmukh, T. Macko, J.-H. Arndt, F. Malz, R. Bernardo, S. Niessen, G. van Doremale, R. Brüll, Critical conditions for liquid chromatography of statistical polyolefins: Evaluation of diene distribution in EPDM terpolymers, *Anal. Chim. Acta* 1246 (2023) 340856.
- [232] M.I. Malik, Novel epoxy-terminated macromonomers and their polymerization for synthesis of bottle-brush type amphiphilic block copolymers, *RSC Adv.* 13 (40) (2023) 28288–28298.