

- [53] I. Souvignet, S.V. Olesik, Liquid chromatography at the critical condition using enhanced-fluidity liquid mobile phases, *Anal. Chem.* 69 (1) (1997) 66–71.
- [54] O. Wachsen, K. Reichert, R. Krüger, H. Much, G. Schulz, Thermal decomposition of biodegradable polyesters—III. Studies on the mechanisms of thermal degradation of oligo-l-lactide using SEC, LACCC and MALDI-TOF-MS, *Polym. Degrad. Stab.* 55 (2) (1997) 225–231.
- [55] D. Berek, M. Janco, G.R. Meira, Liquid chromatography of macromolecules at the critical adsorption point. II. Role of column packing: Bare silica gel, *J. Polym. Sci. Part A: Polym. Chem.* 36 (9) (1998) 1363–1371.
- [56] D. Berek, Liquid chromatography of macromolecules under limiting conditions of desorption. 1. Principles of the method, *Macromol.* 31 (24) (1998) 8517–8521.
- [57] D. Braun, E. Esser, H. Rasch, Chromatographic investigations of macromolecules in the critical range of liquid chromatography. XII. Analysis of block copolymers of styrene and butadiene, *Int. J. Polym. Anal. Charact.* 4 (6) (1998) 501–516.
- [58] A. Gorbunov, L. Ya Solov'yova, A. Skvortsov, General mechanism of the chromatography of macromolecules: Experimental test of the universal nature of adsorption effects, *Polymer* 39 (3) (1998) 697–702.
- [59] R. Murgasova, I. Capek, E. Lathova, D. Berek, S. Florian, Application of a new chromatographic method, LC CAP, in the characterization of polystyrene-graft-poly(ethylene oxide) copolymers, *Eur. Polym. J.* 34 (5–6) (1998) 659–663.
- [60] H. Pasch, K. Rode, Chromatographic investigations of macromolecules in the critical range of liquid chromatography: 11. Polymer blend separation using a reversed stationary phase, *Polymer* 39 (25) (1998) 6377–6383.
- [61] A. Skvortsov, A. Gorbunov, D. Berek, B. Trathnigg, Liquid chromatography of macromolecules at the critical adsorption point: behaviour of a polymer chain inside pores, *Polymer* 39 (2) (1998) 423–429.
- [62] H. Yun, S.V. Olesik, E.H. Marti, Improvements in polymer characterization by size-exclusion chromatography and liquid chromatography at the critical condition by using enhanced-fluidity liquid mobile phases with packed capillary columns, *Anal. Chem.* 70 (15) (1998) 3298–3303.
- [63] J. Adrian, D. Braun, K. Rode, H. Pasch, New analytical methods for epoxy resins, 1. Liquid chromatography at the critical point of adsorption and MALDI-TOF mass spectrometry, *Die Angew. Makromol. Chem.* 267 (1) (1999) 73–81.
- [64] J. Adrian, D. Braun, H. Pasch, New analytical methods for epoxy resins, 2. Two-dimensional chromatography, *Die Angew. Makromol. Chem.* 267 (1) (1999) 82–88.
- [65] K. Baran, S. Laugier, H. Cramail, Investigation of molecular dimensions of polystyrene as a function of the solvent composition: application to liquid chromatography at the exclusion-adsorption transition point, *Macromol. Chem. Phys.* 200 (9) (1999) 2074–2079.
- [66] Y. Brun, The mechanism of copolymer retention in interactive polymer chromatography. II. Gradient separation, *J. Liq. Chromatogr. Relat. Technol.* 22 (20) (1999) 3067–3090.
- [67] T. Chang, H.C. Lee, W. Lee, S. Park, C. Ko, Polymer characterization by temperature gradient interaction chromatography, *Macromol. Chem. Phys.* 200 (10) (1999) 2188–2204.
- [68] K.E. Esser, D. Braun, H. Pasch, Chromatographic investigations of macromolecules in the critical range of liquid chromatography, XIII. Separation of blends of styrene-butadiene rubber and butyl rubber, *Die Angew. Makromol. Chem.* 271 (1) (1999) 61–67.
- [69] R.-P. Krüger, H. Much, G. Schulz, E. Rikowski, Zur charakterisierung von si-polymeren durch flüssigchromatographische verfahren in kopplung mit MALDI-TOF-MS, *Monatsh. Chem. - Chem. Mon.* 130 (1) (1999) 163–174.
- [70] H. Lee, W. Lee, T. Chang, S. Choi, D. Lee, H. Ji, W.K. Nonidez, J.W. Mays, Characterization of poly(ethylene oxide)-block-poly(l-lactide) by HPLC and MALDI-TOF mass spectrometry, *Macromol.* 32 (12) (1999) 4143–4146.
- [71] J. Adrian, E. Esser, G. Hellmann, H. Pasch, Two-dimensional chromatography of complex polymers part 1. Analysis of a graft copolymer by two-dimensional chromatography with on-line FTIR detection, *Polymer* 41 (7) (2000) 2439–2449.
- [72] J. Falkenhagen, H. Much, W. Stauff, A.H.E. Müller, Characterization of block copolymers by liquid adsorption chromatography at critical conditions. 1. Diblock copolymers, *Macromol.* 33 (10) (2000) 3687–3693.
- [73] J. Falkenhagen, J.F. Friedrich, G. Schulz, R.-P. Krüger, H. Much, S. Weidner, Liquid adsorption Chromatography near Critical conditions of adsorption coupled with matrix-assisted laser desorption/ionization mass spectrometry, *Int. J. Polym. Anal. Charact.* 5 (4–6) (2000) 549–562.
- [74] M. Janco, T. Hirano, T. Kitayama, K. Hatada, D. Berek, Discrimination of poly(ethyl methacrylate)s according to their molar mass and tacticity by coupling size exclusion chromatography and liquid chromatography at the critical adsorption point, *Macromol.* 33 (5) (2000) 1710–1715.
- [75] T. Kitayama, M. Janco, K. Ute, R. Niimi, K. Hatada, D. Berek, Analysis of poly(ethyl methacrylate)s by on-line hyphenation of liquid chromatography at the critical adsorption point and nuclear magnetic resonance spectroscopy, *Anal. Chem.* 72 (7) (2000) 1518–1522.
- [76] H.C. Lee, H. Lee, W. Lee, T. Chang, J. Roovers, Fractionation of cyclic polystyrene from linear precursor by HPLC at the chromatographic critical condition, *Macromol.* 33 (22) (2000) 8119–8121.
- [77] B. Lepoittevin, M.-A. Dourges, M. Masure, P. Hemery, K. Baran, H. Cramail, Synthesis and characterization of ring-shaped polystyrenes, *Macromol.* 33 (22) (2000) 8218–8224.
- [78] K. Baran, S. Laugier, H. Cramail, Liquid chromatography of polymers at the exclusion — Adsorption transition point: Physicochemical interpretation, *Int. J. Polym. Anal. Charact.* 6 (1–2) (2001) 123–145.
- [79] K. Baran, S. Laugier, H. Cramail, Fractionation of functional polystyrenes, poly(ethylene oxide)s and poly(styrene)-b-poly(ethylene oxide) by liquid chromatography at the exclusion-adsorption transition point, *J. Chromatogr. B: Biomed. Sci. Appl.* 753 (1) (2001) 139–149.
- [80] I. Bruheim, P. Molander, M. Theodorsen, E. Ommundsen, E. Lundanes, T. Greibrokk, Temperature-programmed packed-capillary liquid chromatographic separation of polystyrenes, *Chromatogr.* 53 (S1) (2001) S266–S270.
- [81] D. Cho, S. Park, K. Kwon, T. Chang, J. Roovers, Structural characterization of ring polystyrene by liquid chromatography at the critical condition and MALDI-TOF mass spectrometry, *Macromol.* 34 (21) (2001) 7570–7572.
- [82] C. Keil, E. Esser, H. Pasch, Matrix-assisted laser desorption/ionization mass spectrometry of synthetic polymers, 5. Analysis of poly(propylene oxide)s by coupled liquid chromatography at the critical point of adsorption and MALDI-TOF mass spectrometry, *Macromol. Mater. Eng.* 286 (3) (2001) 161–167.
- [83] W. Lee, S. Park, T. Chang, Liquid chromatography at the critical condition for polyisoprene using a single solvent, *Anal. Chem.* 73 (16) (2001) 3884–3889.
- [84] W. Lee, D. Cho, T. Chang, K.J. Hanley, T.P. Lodge, Characterization of polystyrene-b-polyisoprene diblock copolymers by liquid chromatography at the chromatographic critical condition, *Macromol.* 34 (7) (2001) 2353–2358.
- [85] H. Lee, T. Chang, D. Lee, M.S. Shim, H. Ji, W.K. Nonidez, J.W. Mays, Characterization of poly(l-lactide)-block-poly(ethylene oxide)-block-poly(l-lactide) triblock copolymer by liquid chromatography at the critical condition and by MALDI-TOF mass spectrometry, *Anal. Chem.* 73 (8) (2001) 1726–1732.
- [86] B. Lepoittevin, X. Perrot, M. Masure, P. Hemery, New route to synthesis of cyclic polystyrenes using controlled free radical polymerization, *Macromol.* 34 (3) (2001) 425–429.
- [87] B. Lepoittevin, P. Hemery, Synthesis of multicyclic and grafted polystyrenes, *J. Polym. Sci. Part A: Polym. Chem.* 39 (16) (2001) 2723–2730.
- [88] H. Pasch, E. Esser, C. Kloninger, H. Iatrou, N. Hadjichristidis, Chromatographic investigations of macromolecules in the critical range of liquid chromatography, 14. Analysis of miktoarm star (μ -star) polymers, *Macromol. Chem. Phys.* 202 (8) (2001) 1424–1429.
- [89] A. Siewing, J. Schierholz, D. Braun, G. Hellmann, H. Pasch, Two-dimensional chromatography of complex polymers, 2. Analysis of the grafting reaction of methyl methacrylate onto EPDM, *Macromol. Chem. Phys.* 202 (14) (2001) 2890–2894.
- [90] T. Macko, D. Hunkeler, D. Berek, Liquid chromatography of synthetic polymers under critical conditions. the case of single eluents and the role of θ conditions, *Macromol.* 35 (5) (2002) 1797–1804.
- [91] S. Phillips, S.V. Olesik, Fundamental studies of liquid chromatography at the critical condition using enhanced-fluidity liquids, *Anal. Chem.* 74 (4) (2002) 799–808.
- [92] T. Biela, A. Duda, K. Rode, H. Pasch, Characterization of star-shaped poly(l-lactide)s by liquid chromatography at critical conditions, *Polymer* 44 (6) (2003) 1851–1860.
- [93] S.M. Graef, A.J.P. Van Zyl, R.D. Sanderson, B. Klumperman, H. Pasch, Use of gradient, critical, and two-dimensional chromatography in the analysis of styrene- and methyl methacrylate-grafted epoxidized natural rubber, *J. Appl. Polym. Sci.* 88 (10) (2003) 2530–2538.
- [94] I. Park, S. Park, D. Cho, T. Chang, E. Kim, K. Lee, Y.J. Kim, Effect of block copolymer chain architecture on chromatographic retention, *Macromol.* 36 (22) (2003) 8539–8543.
- [95] S.L. Phillips, L. Ding, M. Stegemiller, S.V. Olesik, Development of water-based liquid chromatography at the critical condition, *Anal. Chem.* 75 (20) (2003) 5539–5543.
- [96] S.L. Phillips, S.V. Olesik, Initial characterization of humic acids using liquid chromatography at the critical condition followed by size-exclusion chromatography and electrospray ionization mass spectrometry, *Anal. Chem.* 75 (20) (2003) 5544–5553.
- [97] C. Rappel, B. Trathnigg, A. Gorbunov, Liquid chromatography of polyethylene glycol mono- and diesters: functional macromolecules or block copolymers? *J. Chromatogr. A* 984 (1) (2003) 29–43.
- [98] K. Im, S. Park, D. Cho, T. Chang, K. Lee, N. Choi, HPLC and MALDI-TOF MS analysis of highly branched polystyrene: Resolution enhancement by branching, *Anal. Chem.* 76 (9) (2004) 2638–2642.
- [99] H.J. Philipsen, Determination of chemical composition distributions in synthetic polymers, *J. Chromatogr. A* 1037 (1–2) (2004) 329–350.
- [100] X. Jiang, P.J. Schoenmakers, X. Lou, V. Lima, J.L. van Dongen, J. Brokken-Zijp, Separation and characterization of functional poly (n-butyl acrylate) by critical liquid chromatography, *J. Chromatogr. A* 1055 (1–2) (2004) 123–133.
- [101] M. Adler, F. Rittig, S. Becker, H. Pasch, Multidimensional chromatographic and hyphenated techniques for hydrophilic copolymers, *Macromol. Chem. Phys.* 206 (22) (2005) 2269–2277.