

- [102] K.A. Alberty, T.E. Hogen-Esch, S. Carloti, Synthesis and characterization of macrocyclic vinyl-aromatic polymers, *Macromol. Chem. Phys.* 206 (10) (2005) 1035–1042.
- [103] M.A. Bashir, A. Brüll, W. Radke, Fast determination of critical eluent composition for polymers by gradient chromatography, *Polymer* 46 (10) (2005) 3223–3229.
- [104] D. Cho, K. Masuoka, K. Koguchi, T. Asari, D. Kawaguchi, A. Takano, Y. Matsushita, Preparation and characterization of cyclic polystyrenes, *Polym. J.* 37 (7) (2005) 506–511.
- [105] L. Coulier, E. Kaal, T. Hankemeier, Comprehensive two-dimensional liquid chromatography and hyphenated liquid chromatography to study the degradation of poly(bisphenol A)carbonate, *J. Chromatogr. A* 1070 (1–2) (2005) 79–87.
- [106] L.-C. Heinz, S. Graef, T. Macko, R. Brüll, S. Balk, H. Keul, H. Pasch, Development of high-temperature separation techniques for the chemical composition analysis of ethylene-methyl methacrylate block copolymers, *E-Polym.* 5 (1) (2005).
- [107] X. Jiang, A. van der Horst, V. Lima, P.J. Schoenmakers, Comprehensive two-dimensional liquid chromatography for the characterization of functional acrylate polymers, *J. Chromatogr. A* 1076 (1–2) (2005) 51–61.
- [108] V. Lima, X. Jiang, J. Brokken-Zijp, P.J. Schoenmakers, B. Klumperman, R. Van Der Linde, Synthesis and characterization of telechelic polymethacrylates via RAFT polymerization, *J. Polym. Sci. Part A: Polym. Chem.* 43 (5) (2005) 959–973.
- [109] K. Min, H. Gao, K. Matyjaszewski, Preparation of homopolymers and block copolymers in miniemulsion by ATRP using activators generated by electron transfer (AGET), *J. Am. Chem. Soc.* 127 (11) (2005) 3825–3830.
- [110] H. Pasch, M. Adler, D. Knecht, F. Rittig, R. Lange, New chromatographic and hyphenated techniques for hydrophilic copolymers, *Macromol. Symp.* 231 (1) (2005) 166–177.
- [111] W. Radke, K. Rode, A.V. Gorshkov, T. Biela, Chromatographic behavior of functionalized star-shaped poly(lactide)s under critical conditions of adsorption. Comparison of theory and experiment, *Polymer* 46 (15) (2005) 5456–5465.
- [112] B. Trathnigg, Characterization of amphiphilic polymers: Independent analysis of blocks in poloxamers by liquid chromatography under critical conditions, *Polymer* 46 (22) (2005) 9211–9223.
- [113] B. Trathnigg, C. Rappel, S. Fraydl, A. Gorbunov, Liquid chromatography of polyoxyethylenes under critical conditions: A thermodynamic study, *J. Chromatogr. A* 1085 (2) (2005) 253–261.
- [114] M.A. Bashir, W. Radke, Comparison of retention models for polymers, *J. Chromatogr. A* 1131 (1–2) (2006) 130–141.
- [115] L. Coulier, E. Kaal, T. Hankemeier, Towards early detection of the hydrolytic degradation of poly(bisphenol A)carbonate by hyphenated liquid chromatography and comprehensive two-dimensional liquid chromatography, *Polym. Degrad. Stab.* 91 (2) (2006) 271–279.
- [116] L. Coulier, E. Kaal, T. Hankemeier, Hyphenation of infrared spectroscopy to liquid chromatography for qualitative and quantitative polymer analysis: Degradation of poly(bisphenol A)carbonate, *J. Chromatogr. A* 1130 (1) (2006) 34–42.
- [117] H. Gao, K. Min, K. Matyjaszewski, Characterization of linear and 3-arm star block copolymers by liquid chromatography at critical conditions, *Macromol. Chem. Phys.* 207 (19) (2006) 1709–1717.
- [118] K. Im, Y. Kim, T. Chang, K. Lee, N. Choi, Separation of branched polystyrene by comprehensive two-dimensional liquid chromatography, *J. Chromatogr. A* 1103 (2) (2006) 235–242.
- [119] D. Knecht, F. Rittig, R.F. Lange, H. Pasch, Multidimensional chromatographic techniques for hydrophilic copolymers, *J. Chromatogr. A* 1130 (1) (2006) 43–53.
- [120] B. Trathnigg, A. Gorbunov, Characterization of EO-PO block copolymers by liquid chromatography under critical conditions, *Macromol. Symp.* 237 (1) (2006) 18–27.
- [121] A. Albrecht, R. Brüll, T. Macko, H. Pasch, Separation of ethylene-vinyl acetate copolymers by high-temperature gradient liquid chromatography, *Macromol.* 40 (15) (2007) 5545–5551.
- [122] H. Gao, K. Matyjaszewski, Arm-first method as a simple and general method for synthesis of miktoarm star copolymers, *J. Am. Chem. Soc.* 129 (38) (2007) 11828–11834.
- [123] M.d. Geus, Enzymatic Catalysis in the Synthesis of New Polymer Architectures and Materials (PhD thesis), Technische Universiteit Eindhoven, 2007.
- [124] W. Hiller, P. Sinha, H. Pasch, On-line HPLC-NMR of PS-b-PMMA and blends of PS and PMMA: LCCC-NMR at critical conditions of PS, *Macromol. Chem. Phys.* 208 (18) (2007) 1965–1978.
- [125] M. Jacquin, P. Muller, G. Lizarraga, C. Bauer, H. Cottet, O. Théodoly, Characterization of amphiphilic diblock copolymers synthesized by MADIX polymerization process, *Macromol.* 40 (8) (2007) 2672–2682.
- [126] M. Jacquin, P. Muller, R. Talingting-Pabalan, H. Cottet, J. Berret, T. Futterer, O. Théodoly, Chemical analysis and aqueous solution properties of charged amphiphilic block copolymers PBA-b-PAA synthesized by MADIX®, *J. Colloid Interface Sci.* 316 (2) (2007) 897–911.
- [127] M.I. Malik, B. Trathnigg, C.O. Kappe, Selectivity of PEO-block-PPO diblock copolymers in the microwave-accelerated, anionic ring-opening polymerization of propylene oxide with PEG as initiator, *Macromol. Chem. Phys.* 208 (23) (2007) 2510–2524.
- [128] C. Petit, B. Luneau, E. Beaudoin, D. Gimes, D. Bertin, Liquid chromatography at the critical conditions in pure eluent: An efficient tool for the characterization of functional polystyrenes, *J. Chromatogr. A* 1163 (1–2) (2007) 128–137.
- [129] A. Takano, Y. Kushida, K. Aoki, K. Masuoka, K. Hayashida, D. Cho, D. Kawaguchi, Y. Matsushita, HPLC characterization of cyclization reaction product obtained by end-to-end ring closure reaction of a telechelic polystyrene, *Macromol.* 40 (3) (2007) 679–681.
- [130] S. Trimpin, S.M. Weidner, J. Falkenhagen, C.N. McEwen, Fractionation and solvent-free MALDI-MS analysis of polymers using liquid adsorption chromatography at critical conditions in combination with a multisample on-target homogenization/transfer sample preparation method, *Anal. Chem.* 79 (19) (2007) 7565–7570.
- [131] S.M. Weidner, J. Falkenhagen, S. Maltsev, V. Sauerland, M. Rinken, A novel software tool for copolymer characterization by coupling of liquid chromatography with matrix-assisted laser desorption/ionization time-of-flight mass spectrometry, *Rapid Commun. Mass Spectrom.* 21 (16) (2007) 2750–2758.
- [132] S. Weidner, J. Falkenhagen, R.-P. Krueger, U. Just, Principle of two-dimensional characterization of copolymers, *Anal. Chem.* 79 (13) (2007) 4814–4819.
- [133] E. Beaudoin, A. Favier, C. Galindo, A. Lapp, C. Petit, D. Gimes, S. Marquie, D. Bertin, Reduced sample recovery in liquid chromatography at critical adsorption point of high molar mass polystyrene, *Eur. Polym. J.* 44 (2) (2008) 514–522.
- [134] J. Falkenhagen, S. Weidner, Determination of critical conditions of adsorption for chromatography of polymers, *Anal. Chem.* 81 (1) (2008) 282–287.
- [135] H. Gao, K. Matyjaszewski, Synthesis of low-polydispersity miktoarm star copolymers via a simple ‘Arm-First’ method: Macromonomers as arm precursors, *Macromol.* 41 (12) (2008) 4250–4257.
- [136] M.d. Geus, R. Peters, C.E. Koning, A. Heise, Insights into the initiation process of enzymatic ring-opening polymerization from monofunctional alcohols using liquid chromatography under critical conditions, *Biomacromolecules* 9 (2) (2008) 752–757.
- [137] M. Girod, T.N.T. Phan, L. Charles, On-line coupling of liquid chromatography at critical conditions with electrospray ionization tandem mass spectrometry for the characterization of a nitroxide-mediated poly(ethylene oxide)/polystyrene block copolymer, *Rapid Commun. Mass Spectrom.* 22 (23) (2008) 3767–3775.
- [138] A.S. Goldmann, D. Quémener, P.-E. Millard, T.P. Davis, M.H. Stenzel, C. Barner-Kowollik, A.H. Müller, Access to cyclic polystyrenes via a combination of reversible addition fragmentation chain transfer (RAFT) polymerization and click chemistry, *Polymer* 49 (9) (2008) 2274–2281.
- [139] K. Im, H.-W. Park, Y. Kim, S. Ahn, T. Chang, K. Lee, H.-J. Lee, J. Ziebarth, Y. Wang, Retention behavior of star-shaped polystyrene near the chromatographic critical condition, *Macromol.* 41 (9) (2008) 3375–3383.
- [140] J. Listak, W. Jakubowski, L. Mueller, A. Plichta, K. Matyjaszewski, M.R. Bockstaller, Effect of symmetry of molecular weight distribution in block copolymers on formation of ‘metastable’ morphologies, *Macromol.* 41 (15) (2008) 5919–5927.
- [141] M.I. Malik, B. Trathnigg, C.O. Kappe, Microwave assisted synthesis and characterization of end functionalized poly(propylene oxide) as model compounds, *Eur. Polym. J.* 44 (1) (2008) 144–154.
- [142] V. Mass, V. Bellas, H. Pasch, Two-dimensional chromatography of complex polymers, 7 – detailed study of polystyrene-block-polyisoprene diblock copolymers prepared by sequential anionic polymerization and coupling chemistry, *Macromol. Chem. Phys.* 209 (19) (2008) 2026–2039.
- [143] S. Park, D.Y. Ryu, J.K. Kim, M. Ree, T. Chang, Property of diblock copolymer having extremely narrow molecular weight distribution, *Polymer* 49 (8) (2008) 2170–2175.
- [144] H. Pasch, L.-C. Heinz, T. Macko, W. Hiller, High-temperature gradient HPLC and LC-NMR for the analysis of complex polyolefins, *Pure Appl. Chem.* 80 (8) (2008) 1747–1762.
- [145] A.I. Abdulahad, C.Y. Ryu, Liquid chromatography at the critical condition: Thermodynamic significance and influence of pore size, *J. Polym. Sci. Part B: Polym. Phys.* 47 (24) (2009) 2533–2540.
- [146] H. Ahmed, B. Trathnigg, C. Oliver Kappe, R. Saf, Characterization of poly(ethylene glycol)-b-poly(ϵ -caprolactone) by liquid chromatography under critical conditions: Influence of catalysts and reaction conditions on product composition, *Eur. Polym. J.* 45 (8) (2009) 2338–2347.
- [147] H. Ahmed, B. Trathnigg, Characterization of poly(ethylene glycol)-b-poly(ϵ -caprolactone) by two-dimensional liquid chromatography, *J. Sep. Sci.* 32 (9) (2009) 1390–1400.
- [148] C. Galindo, E. Beaudoin, D. Gimes, D. Bertin, Polybutadiene-graft-polystyrene copolymer: Grafting quantification by liquid chromatography at critical conditions using single UV detection, *J. Chromatogr. A* 1216 (47) (2009) 8386–8390.
- [149] M. Girod, E. Beaudoin, L. Charles, Effect of salt in the mobile phase on the critical conditions of poly(ethylene glycol) in liquid chromatography-mass spectrometry coupling, *Anal. Methods* 1 (2) (2009) 128.