

# Appendix A

## Stochastic simulation of epidemics

In this appendix, we discuss some algorithms for performing stochastic simulations and comment on some issues related to interpreting the resulting output.

### A.1 Efficient simulations

We begin with techniques for simulating disease spread in networks. We consider two general algorithms: The Gillespie algorithm and an event-driven algorithm. The Gillespie algorithm is a well-known algorithm for simulating Markovian processes where objects change status. The algorithm calculates the time to the next event and then separately calculates what that event will be. It then jumps to that time, updates statuses and repeats. This is particularly efficient for processes where all that matters is the number of objects of each status, such as a chemical reaction in a well-mixed vessel or disease spread in a well-mixed, mass-action population. For these, the information update at each event simply adjusts the counts. In a network however, knowing the exact node changing status is important. Choosing which node changes status can be a computationally slow process.

An alternative event-driven approach involves a priority queue. Rather than calculating at each time what the next event might be (chosen from all available events), when a node is infected, we calculate when it will transmit to others and when it will recover. These events are placed into the priority queue, and then the next event is removed from the queue and processed. The computational time required to add and remove events is roughly proportional to the log of the number of events already in the queue. This significantly reduces the amount of processing per event, producing a more efficient algorithm. An implementation of these methods is available at

<https://springer-math.github.io/Mathematics-of-Epidemics-on-Networks/>

Although the implementations of the SIS and SIR event-driven algorithms are quite different from the Gillespie algorithms, they simulate the same stochastic processes.

Thus, either could be used for the same purpose. In general, the event-driven algorithms are faster than the equivalent Gillespie algorithms and are more flexible in the case of non-Markovian transmission and recovery processes.

### A.1.1 Gillespie algorithm

**Input:** Network  $G$ , per-edge transmission rate  $\tau$ , recovery rate  $\gamma$ , set of index node(s) initial infecteds, maximum time  $t_{\max}$ .

**Output:** Lists times,  $S$ ,  $I$ , and  $R$  giving number in each state at each time.

```

function Gillespie_network_epidemic( $G, \tau, \gamma$ , initial_infections,  $t_{\max}$ )
    times,  $S, I, R \leftarrow [0], [|G| - \text{len}(\text{initial\_infections})], [\text{len}(\text{initial\_infections})], [0]$ 
    infected_nodes  $\leftarrow$  initial_infections
    at_risk_nodes  $\leftarrow$  uninfected nodes with infected neighbours
    for each node  $u$  in at_risk_nodes do
        infection_rate[ $u$ ] =  $\tau \times$  number of infected neighbours
    total_infection_rate  $\leftarrow \sum_{u \in \text{at\_risk\_nodes}} \text{infection\_rate}[u]$ ,
    total_recovery_rate  $\leftarrow \gamma \times \text{len}(\text{infected\_nodes})$ 
    total_rate  $\leftarrow$  total_infection_rate + total_recovery_rate
    time  $\leftarrow$  exponential_variate(total_rate)
    while time <  $t_{\max}$  and total_rate > 0 do
         $r = \text{uniform\_random}(0, \text{total\_rate})$ 
        if  $r < \text{total\_recovery\_rate}$  then
             $u = \text{random.choice}(\text{infected\_nodes})$ 
            remove  $u$  from infected_nodes
            reduce infection_rate[ $v$ ] for  $u$ 's susceptible neighbours  $v$ 
        else
            choose  $u$  from at_risk_nodes with probability  $\frac{\text{infection\_rate}[u]}{\text{total\_infection\_rate}}$ 
            remove  $u$  from at_risk_nodes
            add  $u$  to infected_nodes
            for susceptible neighbours  $v$  of  $u$  do
                if  $v$  not in at_risk_nodes then
                    add  $v$  to at_risk_nodes
                update infection_rate[ $v$ ]
            update times,  $S, I$ , and  $R$ 
            update total_recovery_rate, total_infection_rate, and total_rate
            time  $\leftarrow$  time + exponential_variate(total_rate)
    return times,  $S, I, R$ 

```

Fig. A.1: Pseudocode for the Gillespie algorithm simulating an SIR epidemic in a network. Small changes are needed for simulating an SIS epidemic: when a node recovers, it needs to become susceptible and possibly needs to be placed into the at-risk node category.

Figure A.1 gives pseudocode adapting the well-known Gillespie algorithm [112, 113] (much of which was introduced earlier by Doob in [83]) to epidemics on networks. This algorithm uses iterative steps:

1. Find the rate of all possible events and compute the total rate of change occurring; this is done by finding the rate at which each infected node will recover and each at-risk susceptible node becomes infected;
2. Based on this rate of change, select the waiting time until the next event from an exponential distribution whose rate is the total rate of change;
3. Select which event occurs, with probability proportional to each event's rate; and
4. Update the rates based on which node changes status and repeat.

In more detail, these steps amount to determining the rate of infection of all susceptible nodes and the rate of recovery of all infectious nodes. Let these be denoted by  $r_u$ , where  $u = 1, 2, 3, \dots, N$  are the nodes. The infection rate of a susceptible node depends on how many infected neighbours it has, but the recovery rate of an infected node is independent of the network and status of neighbours. For example, Fig. A.2 illustrates the computation of some transition rates. Node 1 has one infected neighbour, so  $r_1 = \tau$ , and the two infected neighbours of node 3 yield  $r_3 = 2\tau$ . All infected nodes, e.g. nodes 5 and  $N$  have  $r_5 = r_N = \gamma$ . From this, the total rate of all transitions, denoted by  $T$  (**total\_rate** in the pseudocode), is calculated from the current status of all individuals across the whole network. Therefore,  $T = \sum_{u=1}^N r_u$ , and the waiting time until the next event,  $t_{\text{next}}$ , is chosen from an exponential distribution with rate  $T$ . Because  $r_u = \gamma$  if  $u$  is infected and 0 if  $u$  is susceptible and not in the at-risk nodes, we can write  $T = \gamma[I] + \sum_{u \in \text{at\_risk\_nodes}} r_u$ .




Fig. A.2: Illustrating the relation between the network, status of nodes and events rate vector used by the Gillespie algorithm. Susceptible nodes are denoted by (●) and infected nodes by (●).

Next, a single event is chosen with probability proportional to its rate. Because all recoveries occur with the same rate,  $\gamma$ , the total recovery rate is  $\gamma[I]$ . So the probability the event is a recovery is  $\gamma[I]/T$ . If so, we select a random infected node. If a recovery does not occur, we select a random at-risk node with probability proportional to its rate. Because not all at-risk nodes have the same rate, this step of selecting the node is more involved than simply choosing a random node. It can be done in several ways:

- We can select a random number  $r \in (0, T - \gamma[I])$ . We iterate through the at-risk nodes, each time subtracting  $r_i$  until the result becomes negative. This is the node that recovers. This approach can be slow because, on average, half of the at-risk nodes will be processed in each iteration;
- We can *a priori* artificially inflate the transmission rates, treating all nodes as having the same infection rate  $r^* \geq \max_{u \in \text{at\_risk\_nodes}} \{r_u\}$ . This increases  $T$  to  $\gamma[I] + r^*|\text{at\_risk\_nodes}|$ . When a transmission is predicted to happen, we select a random at-risk node  $u$  whose true rate is  $r_u \leq r^*$ . We then correct for the fact that we have overestimated the risk of  $u$  by selecting an additional random number  $p \in (0, 1)$ . If  $p < r_u/r^*$ , we “accept” the event:  $u$  is infected. Otherwise, we “reject” the event: no nodes change status, but the time  $t$  is increased. This can be slow if much processing time is spent on events in which nothing happens; and
- Alternately, we keep  $T$  as  $T = \gamma[I] + \sum_{u \in \text{at\_risk\_nodes}} r_u$ . We track the maximum transmission rate  $r^*$ . When a transmission event is going to occur, we choose a random node  $u$  from the at-risk nodes. We select an additional random number  $p \in (0, 1)$ . If  $p < r_u/r^*$ , we infect  $u$ . If not, we choose a new random node  $u'$  from the at-risk nodes and a new  $p'$  and repeat. This can be slow if some of these events require repeated selections before finding the newly infected node.

These last two options use a technique known as rejection sampling. Whichever option is chosen, this is generally the slowest step of the algorithm. Once the time to next event and the event itself have been found, necessary rate updates are performed and the process begins again. The Gillespie algorithm allows us to exactly simulate the process.

### A.1.2 Event-driven algorithm

We now consider another efficient algorithm for simulating SIR and SIS epidemics. This uses event-based simulation techniques to improve on Gillespie-style algorithms by avoiding the slow step of finding which node becomes infected. An additional advantage of this approach is that it can be easily generalised for non-Markovian processes.

A key observation is that when a node is infected, nothing that any of its neighbours does will affect when it recovers, who it transmits to or the timing of those transmissions. Thus, as soon as a node is infected, we can calculate when it will recover and when it will transmit to its neighbours. These events are inserted into a priority queue ordered by event time.<sup>1</sup> At each step of the simulation, the next event in the list is removed and processed. If it is a transmission, new events will be added to the queue corresponding to the recovery of the node and any transmissions from that node. This process iterates until a time specified by the user or no events remain in the queue (in which case no infection remains).

Although we assume here that the algorithm is performed using constant rates  $\tau$  and  $\gamma$ , it is straightforward to adapt event-based simulations to other rules. The key requirement is that when a node becomes infected, we can calculate transmission and recovery times without considering events that have not happened yet. Thus, this approach can handle a non-Markovian epidemic process, as in Chapter 9, by some relatively simple adjustments using a different calculation of the duration of infection and the waiting time before transmission.

## SIR epidemics

We begin by describing an event-based algorithm for continuous-time SIR disease transmission in an arbitrary unweighted, undirected network. The algorithm is based on the use of a priority queue  $Q$ . The underlying structure of  $Q$  allows for efficient addition of new events and removal of the earliest remaining event in the queue. Pseudocode for the algorithm is given in Figs. A.3 and A.4.

The algorithm repeatedly removes the earliest remaining event in  $Q$ . If the event is a recovery, the node recovers. If the event is a transmission and the recipient of the transmission has not yet been infected, the node is infected. Now that it is infected, the time of its recovery and the times at which it transmits to its neighbours will not be influenced by any external event. Thus, we can determine the future recovery time and times at which it may transmit to its neighbours immediately, without regard to any events that may happen between now and those events. We add these to  $Q$ , unless we can immediately rule them out of consideration. Since only the first transmission to a node  $v$  has any effect, we can rule out any transmissions for which the recipient has already been infected or has an earlier scheduled transmission. Similarly, we do not add events that would occur after the user-specified  $t_{\max}$  to  $Q$ .

---

<sup>1</sup> A “priority queue” is a data structure which allows for insertion of new events in such a way that we can easily remove the next event to occur. When implemented efficiently, the computational time required to add or remove events is logarithmic in the number of events in the queue [289].

**Input:** Network  $G$ , per-edge transmission rate  $\tau$ , recovery rate  $\gamma$ , set of index node(s) initial infecteds, and maximum time  $t_{\max}$ .

**Output:** Lists times,  $S$ ,  $I$ , and  $R$  giving number in each state at each time.

```

function fast_SIR( $G, \tau, \gamma$ , initial_infecteds,  $t_{\max}$ )
  times,  $S, I, R \leftarrow [0], [|G|], [0], [0]$ 
   $Q \leftarrow$  empty priority queue
  for  $u$  in  $G$ .nodes do
     $u$ .status  $\leftarrow$  susceptible
     $u$ .pred_inf_time  $\leftarrow \infty$ 
  for  $u$  in initial_infecteds do
    Event  $\leftarrow$  {node:  $u$ , time: 0, action: transmit}
     $u$ .pred_inf_time  $\leftarrow$  0
    add Event to  $Q$  ▷ ordered by time
  while  $Q$  is not empty do
    Event  $\leftarrow$  earliest remaining event in  $Q$ 
    if Event.action is transmit then
      if Event.node.status is susceptible then
        process_trans_SIR( $G$ , Event.node, Event.time,  $\tau, \gamma$ , times,  $S, I, R, Q, t_{\max}$ )
      else
        process_rec_SIR(Event.node, Event.time, times,  $S, I, R$ )
  return times,  $S, I, R$ 

```

Fig. A.3: An efficient algorithm simulating continuous-time SIR epidemics in static networks with rates  $\tau$  and  $\gamma$ . It relies on functions given in Fig. A.4. Events are stored in a priority queue  $Q$ , and the first is executed. If the event is a transmission and the recipient is susceptible, it becomes infected, its recovery is added to  $Q$  and transmissions to its neighbours may be added. If the event is a recovery, the node recovers.

## SIS epidemics

With some modest alterations, we can adapt the event-driven SIR disease algorithm to SIS disease. Figures A.5 and A.6 show the pseudocode. The major change is that we must consider the possibility of successful transmissions to a node after its first infection because it may recover to a susceptible state in between. This involves some changes to the algorithm to find the next transmission, and whenever we encounter a transmission event, after checking its effect, we have to consider the possibility that the original source may transmit again.

```

function process_trans_SIR( $G, u, t, \tau, \gamma, \text{times}, S, I, R, Q, t_{\max}$ )
  append times,  $S, I$ , and  $R$  with  $t, S.\text{last}-1, I.\text{last}+1$ , and  $R.\text{last}$ 
   $u.\text{status} \leftarrow \text{infected}$ 
   $u.\text{rec\_time} \leftarrow t + \text{exponential\_variate}(\gamma)$ 
  if  $u.\text{rec\_time} < t_{\max}$  then
    newEvent  $\leftarrow \{\text{node: } u, \text{time: } u.\text{rec\_time}, \text{action: recover}\}$ 
    add newEvent to  $Q$ 
    for  $v$  in  $G.\text{neighbours}(u)$  do
      find_trans_SIR( $Q, t, \tau, u, v, t_{\max}$ )
  function find_trans_SIR( $Q, t, \tau, \text{source}, \text{target}, t_{\max}$ )
    if target.status is susceptible then
      inf_time  $\leftarrow t + \text{exponential\_variate}(\tau)$ 
      if inf_time < minimum(source.rec_time, target.pred_inf_time,  $t_{\max}$ ) then
        newEvent  $\leftarrow \{\text{node: target, time: inf\_time, action: transmit}\}$ 
        add newEvent to  $Q$ 
        target.pred_inf_time  $\leftarrow \text{inf\_time}$ 
  function process_rec_SIR( $u, t, \text{times}, S, I, R$ )
    append times,  $S, I$ , and  $R$  with  $t, S.\text{last}, I.\text{last}-1$ , and  $R.\text{last}+1$ 
     $u.\text{status} \leftarrow \text{recovered}$ 

```

Fig. A.4: Auxiliary functions for fast\_SIR in Fig. A.3. process\_rec\_SIR handles a recovering node. process\_trans\_SIR takes a newly infected node  $u$  and creates a recovery event for  $u$  if the recovery happens before  $t_{\max}$ . For each neighbour  $v$  of  $u$ , it calls find\_trans\_SIR to determine when  $u$  would transmit to  $v$ , adding a transmission event to  $Q$  if the event might occur ( $u$  still infected and  $v$  not known to be already infected by then).

## A.2 Time shifting of simulation results

We now consider a technical issue related to comparing deterministic models with stochastic simulations. When a stochastic simulation begins with a small number of infections, it may die out, it may have unusually fast early growth or it may take a long time before successfully becoming established and growing. The deterministic models do not have this variability.

When we use a simulation to study epidemics, we must account for these stochastic delays and the fact that some realisations die out. If our goal is to predict the future spread of an epidemic that has already established itself, we should obviously discard the outbreaks that died out stochastically.

It is natural to consider the average of the simulations that do not die out. However, there is an additional issue. The stochastic effects that may accelerate or delay the onset of the epidemic mean that simulated epidemics become large at different times. In Fig. A.7, the epidemics are effectively time translations of one another. Thus, the peaks occur with some stochastic delay. If the population model and the

**Input:** Network  $G$ , transmission rate per edge  $\tau$ , recovery rate  $\gamma$ , set of index node(s) initial infecteds, and maximum time  $t_{\max}$ .

**Output:**  $t$ : list of times and  $I$ : list containing number infected at each time.

```

function fast_SIS( $G, \tau, \gamma$ , initial infecteds,  $t_{\max}$ )
    initialise  $Q$ , node statuses and return variables as in fast_SIR, but include a source for
    infections with recovery time 0.
    while  $Q$  is not empty do
        Event  $\leftarrow$  earliest remaining event in  $Q$ 
        if Event.action is transmit then
            if Event.node.status is susceptible then
                process_trans_SIS( $G$ , Event.node, Event.time,  $\tau, \gamma$ , times,  $S, I, Q, t_{\max}$ )
                find_next_trans_SIS( $Q$ , Event.source, Event.node,  $t$ )  ▷ needed for SIS model
            else
                process_rec_SIS(Event.node,  $t, S, I$ )
    return times,  $S, I$ 

```

Fig. A.5: An efficient event-based algorithm simulating continuous-time SIS epidemics in static networks. It relies on functions given in Fig. A.6. It is similar to the algorithm in Fig. A.3. When nodes recover, they become susceptible again. An important change compared to the SIR version is that when a transmission event occurs, we check the possibility that the source of that transmission might cause another transmission later. This requires that we include the source in every transmission event.

disease model accurately represent the real-world epidemic, we would expect the real epidemic to have the same shape, with some stochastic delay. If we try to capture this by taking the average of many simulations, the result will have a different shape. By averaging many peaks that do not align, we get a lower, wider peak which does not resemble any single realisation.

Thus, we have two options to align trajectories:

- We can take many simulations and shift time so that  $t = 0$  corresponds to the prevalence crossing some threshold size; or
- We can initialise our simulations with a large number of infections such that the stochastic noise is negligible.

Both of these assume that the epidemic is in a large enough population so that it is reasonable to talk about the deterministic phase of an epidemic.



```

function process_trans_SIS( $G, u, t, \tau, \gamma, \text{times}, S, I, Q, t_{\max}$ )
  append times,  $S$ , and  $I$  with  $t$ ,  $S.\text{last}-1$ , and  $I.\text{last}+1$ 
   $u.\text{status} \leftarrow \text{infected}$ 
   $u.\text{rec\_time} \leftarrow t + \text{exponential\_variate}(\gamma)$ 
  if  $u.\text{rec\_time} < t_{\max}$  then
    newEvent  $\leftarrow \{\text{node: } u, \text{time: } u.\text{rec\_time}, \text{action: recover}\}$ 
    add newEvent to  $Q$ 
  for  $v$  in  $G.\text{neighbours}(u)$  do
    find_next_trans_SIS( $Q, t, \tau, u, v, t_{\max}$ )
  function find_next_trans_SIS( $Q, t, \tau, \text{source}, \text{target}, t_{\max}$ )
    if  $\text{target.rec\_time} < \text{source.rec\_time}$  then
      transmission_time =  $\max(t, \text{target.rec\_time}) + \text{exponential\_variate}(\tau)$ 
      if transmission_time <  $\text{source.rec\_time}$  then
        newEvent  $\leftarrow \{\text{node: target, time: transmission\_time, action: transmit, source:}$ 
        source $\}$ 
        push( $Q$ , newEvent)
  function process_rec_SIS( $u, \text{times}, S, I$ )
    append times,  $S$ , and  $I$  with  $t$ ,  $S.\text{last}+1$ , and  $I.\text{last}-1$ 
     $u.\text{status} \leftarrow \text{susceptible}$ 

```

Fig. A.6: Auxiliary functions for event-based SIS simulation. Recovery is handled as in the SIR case except that the node becomes susceptible again. Each transmission is handled similarly, but when finding the next transmission event, there are fewer events we can exclude from adding to  $Q$ . We find the first transmission from the source to the recipient and add it to  $Q$  unless the recipient is infected and will not recover prior to the transmission or the source will recover prior to the transmission. Since multiple transmissions from a source to a target may cause infection, after each transmission event, `fast_SIS` also calls `find_next_trans`.




Fig. A.7: (Left) SIR simulations in a  $10^6$  node Configuration Model network with  $P(2) = P(3) = P(4) = P(5) = 1/4$  and  $\tau = \gamma = 1$ . Simulations were run until 200 epidemics occurred, using the algorithm in Fig. A.3. These epidemics are shown: three are highlighted, and the average of all 200 is dashed. The average does not resemble individual realisations. (Right) Shifting time so that  $t = 0$  when 1% of the population is infected aligns individual realisations and the average.

## References

1. Allard, A., Hébert-Dufresne, L., Young, J.G., Dubé, L.J.: General and exact approach to percolation on random graphs. *Phys. Rev. E* **92**(6), 062807 (2015)
2. Allard, A., Noël, P.A., Dubé, L.J., Pourbohloul, B.: Heterogeneous bond percolation on multitype networks with an application to epidemic dynamics. *Phys. Rev. E* **79**(3), 036113 (2009)
3. Andersson, H., Britton, T.: *Stochastic Epidemic Models and Their Statistical Analysis*, vol. 151. Springer Science & Business Media, New York (2012)
4. Anderson, R.M., May, R.M.: *Infectious Diseases of Humans: Dynamics and Control*, vol. 1. Oxford University Press, Oxford (1991)
5. Armbruster, B., Besenyei, A., Simon, P.L.: Bounds for the expected value of one-step processes. *arXiv preprint arXiv:1505.00898* (2015)
6. Artalejo, J.R.: On the time to extinction from quasi-stationarity: a unified approach. *Phys. A: Stat. Mech. Appl.* **391**(19), 4483–4486 (2012)
7. Artalejo, J.R., Economou, A., Lopez-Herrero, M.J.: Stochastic epidemic models with random environment: quasi-stationarity, extinction and final size. *J. Math. Biol.* **67**(4), 799–831 (2013)
8. Ball, F.: A unified approach to the distribution of total size and total area under the trajectory of infectives in epidemic models. *Adv. Appl. Probab.* **18**(2), 289–310 (1986)
9. Ball, F., Lyne, O.D.: Stochastic multi-type SIR epidemics among a population partitioned into households. *Adv. Appl. Probab.* **33**(1), 99–123 (2001)
10. Ball, F.G., Lyne, O.D.: Epidemics among a population of households. In: *Mathematical Approaches for Emerging and Reemerging Infectious Diseases: Models, Methods, and Theory*, pp. 115–142. Springer, New York (2002)
11. Ball, F.G., Lyne, O.D.: Optimal vaccination policies for stochastic epidemics among a population of households. *Math. Biosci.* **177**, 333–354 (2002)
12. Ball, F., Neal, P.: A general model for stochastic SIR epidemics with two levels of mixing. *Math. Biosci.* **180**(1), 73–102 (2002)
13. Ball, F., Neal, P.: Network epidemic models with two levels of mixing. *Math. Biol.* **212**(1), 69–87 (2008)
14. Ball, F., Sirl, D.: An SIR epidemic model on a population with random network and household structure, and several types of individuals. *Adv. Appl. Probab.* **44**(1), 63–86 (2012)
15. Ball, F., Sirl, D.: Acquaintance vaccination in an epidemic on a random graph with specified degree distribution. *J. Appl. Probab.* **50**(4), 1147–1168 (2013)
16. Ball, F., Sirl, D., Trapman, P.: Threshold behaviour and final outcome of an epidemic on a random network with household structure. *Adv. Appl. Probab.* **41**(3), 765–796 (2009)
17. Ball, F., Sirl, D., Trapman, P.: Analysis of a stochastic SIR epidemic on a random network incorporating household structure. *Math. Biosci.* **224**(2), 53–73 (2010)

18. Ball, F., Britton, T., Sirl, D.: A network with tunable clustering, degree correlation and degree distribution, and an epidemic thereon. *J. Math. Biol.* **66**(4–5), 979–1019 (2013)
19. Ball, F., Britton, T., Neal, P.: On expected durations of birth-death processes, with applications to branching processes and SIS epidemics. *J. Appl. Probab.* **53**(1), 203–215 (2016)
20. Ball, F., Britton, T., House, T., Isham, V., Mollison, D., Pellis, L., Tomba, G.S.: Seven challenges for metapopulation models of epidemics, including households models. *Epidemics* **10**, 63–67 (2015)
21. Bansal, S., Khandelwal, S., Meyers, L.A.: Exploring biological network structure with clustered random networks. *BMC Bioinformatics* **10**(1), 405 (2009)
22. Bansal, S., Read, J., Pourbohloul, B., Meyers, L.A.: The dynamic nature of contact networks in infectious disease epidemiology. *J. Biol. Dyn.* **4**(5), 478–489 (2010)
23. Barabási, A.L., Albert, R.: Emergence of scaling in random networks. *Science* **286**(5439), 509–512 (1999)
24. Barbour, A.D., Reinert, G.: Approximating the epidemic curve. *Electron. J. Probab* **18**(54), 1–30 (2013)
25. Barr, D.R., Thomas, M.U.: An eigenvector condition for Markov chain lumpability. *Oper. Res.* **25**, 1028–1031 (1977)
26. Barrat, A., Barthélemy, M., Vespignani, A.: Dynamical processes on complex networks. Cambridge University Press, New York (2008)
27. Barthélemy, M., Barrat, A., Pastor-Satorras, R., Vespignani, A.: Dynamical patterns of epidemic outbreaks in complex heterogeneous networks. *J. Theor. Biol.* **235**(2), 275–288 (2005)
28. Bátkai, A., Havasi, Á., Horváth, R., Kunszenti-Kovács, D., Simon, P.L.: PDE approximation of large systems of differential equations. *Oper. Matrices* **9**(1), 147–163 (2015)
29. Bátkai, A., Kiss, I.Z., Sikolya, E., Simon, P.L.: Differential equation approximations of stochastic network processes: an operator semigroup approach. *Netw. Heterog. Media* **7**(1), 43–58 (2012)
30. Battiston, F., Nicosia, V., Latora, V.: Structural measures for multiplex networks. *Phys. Rev. E* **89**(3), 032804 (2014)
31. Bauch, C.T.: The spread of infectious diseases in spatially structured populations: an invasy pair approximation. *Math. Biosci.* **198**(2), 217–237 (2005)
32. Bauch, C.T., Lloyd-Smith, J.O., Coffee, M.P., Galvani, A.P.: Dynamically modeling SARS and other newly emerging respiratory illnesses: past, present, and future. *Epidemiology* **16**(6), 791–801 (2005)
33. Bearman, P.S., Moody, J., Stovel, K.: Chains of affection: the structure of adolescent romantic and sexual networks. *Am. J. Sociol.* **110**(1), 44–91 (2004)
34. ben Avraham, D., Boltt, E.M., Tamon, C.: One-dimensional continuous-time quantum walks. *Quant. Inf. Process.* **3**(1–5), 295–308 (2004)
35. Bobrowski, A.: Functional Analysis for Probability and Stochastic Processes: An Introduction. Cambridge University Press, Cambridge (2005)

36. Boccaletti, S., Latora, V., Moreno, Y., Chavez, M., Hwang, D.U.: Complex networks: structure and dynamics. *Phys. Rep.* **424**(4), 175–308 (2006)
37. Boccaletti, S., Bianconi, G., Criado, R., Del Genio, C.I., Gómez-Gardeñes, J., Romance, M., Sendiña-Nadal, I., Wang, Z., Zanin, M.: The structure and dynamics of multilayer networks. *Phys. Rep.* **544**(1), 1–122 (2014)
38. Bodó, Á., Katona, G.Y., Simon, P.L.: SIS epidemic propagation on hypergraphs. *Bull. Math. Biol.* **78**(4), 713–735 (2016)
39. Boguñá, M., Pastor-Satorras, R.: Epidemic spreading in correlated complex networks. *Phys. Rev. E* **66**(4), 047104 (2002)
40. Boguñá, M., Pastor-Satorras, R., Vespignani, A.: Absence of epidemic threshold in scale-free networks with degree correlations. *Phys. Rev. Lett.* **90**(2), 028701 (2003)
41. Boguñá, M., Castellano, C., Pastor-Satorras, R.: Nature of the epidemic threshold for the susceptible-infected-susceptible dynamics in networks. *Phys. Rev. Lett.* **111**(6), 068701 (2013)
42. Boguñá, M., Lafuerza, L.F., Toral, R., Serrano, M.A.: Simulating non-Markovian stochastic processes. *Phys. Rev. E* **90**(4), 042108 (2014)
43. Bollobás, B.: *Random Graphs*, 2nd edn. Cambridge University Press, Cambridge (2001)
44. Bollobás, B., Kozma, R., Miklos, D.: *Handbook of Large-Scale Random Networks*, vol. 18. Springer Science & Business Media, Berlin, Heidelberg (2010)
45. Bornholdt, S., Schuster, H.G.: *Handbook of Graphs and Networks: From the Genome to the Internet*. Wiley, Berlin (2006)
46. Box, G.E.P., Hunter, J.S., Hunter, W.G.: Statistics for experimenters: design, innovation, and discovery. *AMC* **10**, 12 (2005)
47. Brauer, F., Castillo-Chavez, C.: *Mathematical Models in Population Biology and Epidemiology*, 2nd edn. Springer, New York (2012)
48. Brauer, F., van den Driessche, P., Wu, J.: *Mathematical Epidemiology*, vol. 1945. Springer, Berlin, Heidelberg (2008)
49. Britton, T.: Stochastic epidemic models: a survey. *Math. Biosci.* **225**(1), 24–35 (2010)
50. Britton, T., Trapman, P.: Inferring global network properties from egocentric data with applications to epidemics. *Math. Med. Biol.* **32**(1), 101–114 (2015)
51. Britton, T., Deijfen, M., Martin-Löf, A.: Generating simple random graphs with prescribed degree distribution. *J. Stat. Phys.* **124**(6), 1377–1397 (2006)
52. Broder, A., Kumar, R., Maghoul, F., Raghavan, P., Rajagopalan, S., Stata, R., Tomkins, A., Wiener, J.: Graph structure in the web. *Comput. Netw.* **33**(1), 309–320 (2000)
53. Brown, R.F.: *A Topological Introduction to Nonlinear Analysis*. Springer, Berlin (1993)
54. Butler, B.K., Siegl, P.H.: Sharp bounds on the spectral radius of nonnegative matrices and digraphs. *Linear Algebra Appl.* **439**(5), 1468–1478 (2013)
55. Caldarelli, G.: *Scale-Free Networks: Complex Webs in Nature and Technology*. Oxford University Press, Oxford (2007)

56. Cardy, J.L., Grassberger, P.: Epidemic models and percolation. *J. Phys. A: Math. Gen.* **18**(6), L267–L271 (1985)
57. Castellano, C., Pastor-Satorras, R.: Thresholds for epidemic spreading in networks. *Phys. Rev. Lett.* **105**(21), 218701 (2010)
58. Castellano, C., Fortunato, S., Loreto, V.: Statistical physics of social dynamics. *Rev. Mod. Phys.* **81**(591), 591–646 (2009)
59. Castillo-Chavez, C., Song, B.: Dynamical models of tuberculosis and their applications. *Math. Biosci. Eng.* **1**(2), 361–404 (2004)
60. Cator, E., Van Mieghem, P.: Nodal infection in Markovian susceptible-infected-susceptible and susceptible-infected-removed epidemics on networks are non-negatively correlated. *Phys. Rev. E* **89**(5), 052802 (2014)
61. Cator, E., van de Bovenkamp, R., Van Mieghem, P.: Susceptible-infected-susceptible epidemics on networks with general infection and cure times. *Phys. Rev. E* **87**(6), 062816 (2013)
62. Centers for Disease Control and Prevention: Severe acute respiratory syndrome — Singapore, 2003. *Morb. Mortal. Wkly. Rep.* **52**(18), 405–411 (2003)
63. Chatterjee, S., Durrett, R.: Contact processes on random graphs with power law degree distributions have critical value 0. *Ann. Probab.* **37**(6), 2332–2356 (2009)
64. Christakis, N.A., Fowler, J.H.: Social network sensors for early detection of contagious outbreaks. *PLoS ONE* **5**(9), e12948 (2010)
65. Clancy, D.: SIR epidemic models with general infectious period distribution. *Stat. Probab. Lett.* **85**, 1–5 (2014)
66. Clancy, D., Mendy, S.T.: Approximating the quasi-stationary distribution of the SIS model for endemic infection. *Methodol. Comput. Appl. Probab.* **13**(3), 603–618 (2011)
67. Clarke, J., White, K.A.J., Turner, K.: Approximating optimal controls for networks when there are combinations of population-level and targeted measures available: chlamydia infection as a case-study. *Bull. Math. Biol.* **75**(10), 1747–1777 (2013)
68. Cohen, R., Havlin, S.: *Complex Networks: Structure, Robustness and Function*. Cambridge University Press, Cambridge (2010)
69. Courtesy of Salathé Lab, Penn State University: Flu Outbreaks Modeled by new Study of Classroom Schedules. <http://science.psu.edu/news-and-events/2013-news/Salathe2-2013>. Accessed 31 July 2016
70. Cowan, J.D.: *Proceedings of the 1990 Conference on Advances in Neural Information Processing Systems 3: Stochastic Neurodynamics*. Morgan Kaufmann Publishers Inc., San Francisco, CA (1990)
71. Cowan, N.J., Chastain, E.J., Vilhena, D.A., Freudenberg, J.S., Bergstrom, C.T.: Nodal dynamics, not degree distributions, determine the structural controllability of complex networks. *PLoS ONE* **7**(6), e38398 (2012)
72. Cozzo, E., Banos, R.A., Meloni, S., Moreno, Y.: Contact-based social contagion in multiplex networks. *Phys. Rev. E* **88**(5), 050801 (2013)

73. Cvetkovski, Z.: *Inequalities: Theorems, Techniques and Selected Problems*. Springer Science & Business Media, Berlin/Heidelberg (2012)
74. Daley, D.J., Gani, J.: *Epidemic Modelling: An Introduction*, vol. 15. Cambridge University Press, Cambridge (2001)
75. Danon, L., Ford, A.P., House, T., Jewell, C.P., Keeling, M.J., Roberts, G.O., Ross, J.V., Vernon, M.C.: Networks and the epidemiology of infectious disease. *Interdiscip. Perspect. Infect. Dis.* **2011**, Article ID 284909, 1–28. (2011).
76. Decreusefond, L., Dhersin, J.S., Moyal, P., Tran, V.C.: Large graph limit for an SIR process in random network with heterogeneous connectivity. *Ann. Appl. Probab.* **22**(2), 541–575 (2012)
77. de Oliveira, M.J., Mendes, J.F.F., Santos, M.A.: Nonequilibrium spin models with Ising universal behaviour. *J. Phys. A: Math. Gen.* **26**, 2317 (1993)
78. Diekmann, O., Heesterbeek, J.A.P.: *Mathematical Epidemiology of Infectious Diseases: Model Building, Analysis and Interpretation*. Wiley, New York (2000)
79. Diekmann, O., Heesterbeek, J.A.P., Metz, J.A.J.: On the definition and the computation of the basic reproduction ratio  $\mathcal{R}_0$  in models for infectious diseases in heterogeneous populations. *J. Math. Biol.* **28**(4), 365–382 (1990)
80. Diekmann, O., De Jong, M.C.M., Metz, J.A.J.: A deterministic epidemic model taking account of repeated contacts between the same individuals. *J. Appl. Probab.* **35**(2), 448–462 (1998)
81. Diekmann, O., Heesterbeek, H., Britton, T.: *Mathematical Tools for Understanding Infectious Disease Dynamics*. Princeton University Press, Princeton (2012)
82. Diestel, R.: *Graph Theory*. Springer, Heidelberg, New York (2005)
83. Doob, J.L.: Markoff chains—denumerable case. *Trans. Am. Math. Soc.* **58**(3), 455–473 (1945)
84. Dorogovtsev, S.N., Mendes, J.F.F., Samukhin, A.N.: Giant strongly connected component of directed networks. *Phys. Rev. E* **64**(2), 025101 (2001)
85. Dorogovtsev, S.N., Goltsev, A.V., Mendes, J.F.F.: Ising model on networks with an arbitrary distribution of connections. *Phys. Rev. E* **66**(1), 016104–1–016104-5 (2002)
86. Draief, M., Massoulié, L.: *Epidemics and Rumours in Complex Networks*. Cambridge University Press, New York (2010)
87. Durrett, R.: *Random Graph Dynamics*. Cambridge University Press, Cambridge (2007)
88. Durrett, R.: Some features of the spread of epidemics and information on a random graph. *Proc. Natl. Acad. Sci.* **107**(10), 4491–4498 (2010)
89. Eames, K.T.D.: Modelling disease spread through random and regular contacts in clustered populations. *Theor. Popul. Biol.* **73**(1), 104–111 (2008)
90. Eames, K.T.D., Keeling, M.J.: Modeling dynamic and network heterogeneities in the spread of sexually transmitted diseases. *Proc. Natl. Acad. Sci.* **99**(20), 13330–13335 (2002)
91. Eames, K.T.D., Keeling, M.J.: Contact tracing and disease control. *Proc. R. Soc. Lond. B: Biol. Sci.* **270**(1533), 2565–2571 (2003)

92. Easley, D., Kleinberg, J.: *Networks, Crowds, and Markets: Reasoning About a Highly Connected World*. Cambridge University Press, Cambridge (2010)
93. Estrada, E.: *The Structure of Complex Networks: Theory and Applications*. Oxford University Press, Oxford (2011)
94. Ethier, S.N., Kurtz, T.G.: *Markov Processes: Characterization and Convergence*, vol. 282. Wiley, Hoboken/New Jersey (2009)
95. Euler, L.: *Solutio problematis ad geometriam situs pertinentis*. *Commentarii academiae scientiarum Petropolitanae* **8**, 128–140 (1741)
96. Feld, S.L.: Why your friends have more friends than you do. *Am. J. Sociol.* **96**(6), 1464–1477 (1991)
97. Fennell, P.G., Melnik, S., Gleeson, J.P.: Limitations of discrete-time approaches to continuous-time contagion dynamics. *Phys. Rev. E* **94**(5), 052125 (2016)
98. Ferreira, S.C., Castellano, C., Pastor-Satorras, R.: Epidemic thresholds of the susceptible-infected-susceptible model on networks: a comparison of numerical and theoretical results. *Phys. Rev. E* **86**(4), 041125 (2012)
99. Fiedler, M.: *Special Matrices and Their Applications in Numerical Mathematics*. Dover Publications, New York (2008)
100. Filliger, R., Hongler, M.O.: Lumping complex networks. In: *Lectures and Gallery of Madeira Math Encounters XXXV* (2008). <http://ccm.uma.pt/mme35/>
101. Fotouhi, B., Rabbat, M.G.: Degree correlation in scale-free graphs. *Eur. Phys. J. B* **86**(12), 1–19 (2013)
102. Frasca, M., Sharkey, K.J.: Discrete-time moment closure models for epidemic spreading in populations of interacting individuals. *J. Theor. Biol.* **399**, 13–21 (2016)
103. Frieze, A., Karoński, M.: *Introduction to Random Graphs*. Cambridge University Press, Cambridge (2015)
104. Fu, X., Small, M., Chen, G.: *Propagation Dynamics on Complex Networks: Models, Methods and Stability Analysis*. Wiley, Chichester/UK (2013)
105. Funk, S., Jansen, V.A.A.: Interacting epidemics on overlay networks. *Phys. Rev. E* **81**(3), 036118 (2010)
106. Funk, S., Gilad, E., Watkins, C., Jansen, V.A.A.: The spread of awareness and its impact on epidemic outbreaks. *Proc. Natl. Acad. Sci.* **106**(16), 6872–6877 (2009)
107. Funk, S., Salathé, M., Jansen, V.A.A.: Modelling the influence of human behaviour on the spread of infectious diseases: a review. *J. R. Soc. Interface* **7**(50), 1247–1256 (2010)
108. Gantmacher, F.R.: *The Theory of Matrices*. Taylor & Francis, Providence/Rhode Island (1964)
109. Gershgorin, S.A.: Über die abgrenzung der eigenwerte einer matrix. *Izvestiya Rossiiskoi Akademii Nauk, Seriya Matematicheskaya* **6**, 749–754 (1931)
110. Gertsbakh, I.B.: Epidemic process on a random graph: some preliminary results. *J. Appl. Probab.* **14**(03), 427–438 (1977)

111. Ghoshal, G., Zlatic, V., Caldarelli, G., Newman, M.E.J.: Random hypergraphs and their applications. *Phys. Rev. E* **79**, 066118 (2009)
112. Gillespie, D.T.: A general method for numerically simulating the stochastic time evolution of coupled chemical reactions. *J. Comput. Phys.* **22**(4), 403–434 (1976)
113. Gillespie, D.T.: Exact stochastic simulation of coupled chemical reactions. *J. Phys. Chem.* **81**(25), 2340–2361 (1977)
114. Gleeson, J.P.: Bond percolation on a class of clustered random networks. *Phys. Rev. E* **80**(3), 036107 (2009)
115. Gleeson, J.P.: High-accuracy approximation of binary-state dynamics on networks. *Phys. Rev. Lett.* **107**(6), 068701 (2011)
116. Gleeson, J.P.: Binary-state dynamics on complex networks: pair approximation and beyond. *Phys. Rev. X* **3**(2), 021004 (2013)
117. Gleeson, J.P., Melnik, S., Hackett, A.: How clustering affects the bond percolation threshold in complex networks. *Phys. Rev. E* **81**(6), 066114 (2010)
118. Goldstein, E., Paur, K., Fraser, C., Kenah, E., Wallinga, J., Lipsitch, M.: Reproductive numbers, epidemic spread and control in a community of households. *Math. Biosci.* **221**(1), 11–25 (2009)
119. Golub, G.H., Van Loan, C.F.: *Matrix Computations*, vol. 3. JHU Press, Baltimore/Maryland (2012)
120. Granell, C., Gómez, S., Arenas, A.: Dynamical interplay between awareness and epidemic spreading in multiplex networks. *Phys. Rev. Lett.* **111**(12), 128701 (2013)
121. Grassberger, P.: On the critical behavior of the general epidemic process and dynamical percolation. *Math. Biosci.* **63**, 157–172 (1983)
122. Green, D.M., Kiss, I.Z.: Large-scale properties of clustered networks: implications for disease dynamics. *J. Biol. Dyn.* **4**(5), 431–445 (2010)
123. Griffeath, D.: *Additive and Cancellative Interacting Particle Systems*. Springer, Berlin (1979)
124. Grimmett, G., Stirzaker, D.: *Probability and Random Processes*, 3rd edn. Oxford University Press, Oxford (2001)
125. Grindrod, P., Higham, D.J.: Evolving graphs: dynamical models, inverse problems and propagation. *Proc. R. Soc. Lond. A: Math. Phys. Eng. Sci.* **466**(2115), 753–770 (2010)
126. Gross, T., Blasius, B.: Adaptive coevolutionary networks: a review. *J. R. Soc. Interface* **5**(20), 259–271 (2008)
127. Gross, T., Kevrekidis, I.G.: Robust oscillations in SIS epidemics on adaptive networks: coarse graining by automated moment closure. *EPL (Europhys. Lett.)* **82**(3), 38004 (2008)
128. Gross, T., D’Lima, C.J.D., Blasius, B.: Epidemic dynamics on an adaptive network. *Phys. Rev. Lett.* **96**(20), 208701 (2006)
129. Hadjichrysanthou, C., Sharkey, K.J.: Epidemic control analysis: designing targeted intervention strategies against epidemics propagated on contact networks. *J. Theor. Biol.* **365**, 84–95 (2015)



130. Hagberg, A.A., Schult, D.A., Swart, P.J.: Exploring network structure, dynamics, and function using NetworkX. In: *Proceedings of the 7th Python in Science Conferences (SciPy 2008)*, vol. 2008, pp. 11–16 (2008)
131. Hale, J.K.: *Ordinary Differential Equations*. Dover Books on Mathematics Series. Dover Publications, New York (2009)
132. Halliday, D.M., Rosenberg, J.R., Amjad, A.M., Breeze, P., Conway, B.A., Farmer, S.F.: A framework for the analysis of mixed time series/point process data—theory and application to the study of physiological tremor, single motor unit discharges and electromyograms. *Progr. Biophys. Mol. Biol.* **64**(2), 237–278 (1995)
133. Harada, Y., Ezoe, H., Iwasa, Y., Matsuda, H., Sato, K.: Population persistence and spatially limited social interaction. *Theor. Popul. Biol.* **48**(1), 65–91 (1995)
134. Harris, T.E.: On a class of set-valued Markov processes. *Ann. Probab.* **4**(2), 175–194 (1976)
135. Hastings, M.B.: Systematic series expansions for processes on networks. *Phys. Rev. Lett.* **96**(14), 148701 (2006)
136. Hatzopoulos, V., Taylor, M., Simon, P.L., Kiss, I.Z.: Multiple sources and routes of information transmission: implications for epidemic dynamics. *Math. Biosci.* **231**(2), 197–209 (2011)
137. Hayden, R.A., Horváth, I., Telek, M.: Mean field for performance models with generally-distributed timed transitions. In: *Quantitative Evaluation of Systems*, pp. 90–105. Springer, Heidelberg/New York (2014)
138. Hébert-Dufresne, L., Patterson-Lomba, O., Goerg, G.M., Althouse, B.M.: Pathogen mutation modeled by competition between site and bond percolation. *Phys. Rev. Lett.* **110**, 108103 (2013)
139. Heesterbeek, H., Anderson, R.M., Andreasen, V., Bansal, S., De Angelis, D., Dye, C., Eames, K.T.D., Edmunds, W.J., Frost, S.D.W., Funk, S., Hollingsworth, T.D., House, T., Isham, V., Klepac, P., Lessler, J., Lloyd-Smith, J.O., Metcalf, C.J.E., Mollison, D., Pellis, L., Pulliam, J.R.C., Roberts, M.G., Viboud, C.: Modeling infectious disease dynamics in the complex landscape of global health. *Science* **347**(6227), aaa4339 (2015)
140. Hethcote, H.W., Yorke, J.A.: *Gonorrhea Transmission Dynamics and Control*. Lecture Notes in Biomathematics, vol. 56. Springer, Berlin/Heidelberg (1984)
141. Hethcote, H.W., Yorke, J.A., Nold, A.: Gonorrhea modeling: a comparison of control methods. *Math. Biosci.* **58**(1), 93–109 (1982)
142. Hirsch, M.W., Smith, H.: Monotone dynamical systems. In: A. Cañada, P. Drábek, A. Fonda (eds.) *Handbook of Differential Equations: Ordinary Differential Equations*, vol. 2, pp. 239–357. Elsevier BV, Amsterdam (2005)
143. Hladish, T., Melamud, E., Barrera, L.A., Galvani, A., Meyers, L.A.: Epifire: an open source C++ library and application for contact network epidemiology. *BMC Bioinformatics* **13**(1), 1 (2012)
144. Hoen, A.G., Hladish, T.J., Eggo, R.M., Lenczner, M., Brownstein, J.S., Meyers, L.A.: Epidemic wave dynamics attributable to urban community struc-

- ture: a theoretical characterization of disease transmission in a large network. *J. Med. Internet Res.* **17**(7) (2015).
145. Holley, R.A., Liggett, T.M.: Ergodic theorems for weakly interacting infinite systems and the voter model. *Ann. Probab.* **3**(4), 643–663 (1975)
  146. Holme, P., Saramäki, J.: Temporal networks. *Phys. Rep.* **519**(3), 97–125 (2012)
  147. Holme, P., Saramäki, J.: Temporal Networks. Springer, Berlin (2013)
  148. House, T., Keeling, M.J.: Deterministic epidemic models with explicit household structure. *Math. Biosci.* **213**(1), 29–39 (2008)
  149. House, T., Keeling, M.J.: The impact of contact tracing in clustered populations. *PLoS Computat. Biol.* **6**(3), e1000721 (2010)
  150. House, T., Keeling, M.J.: Epidemic prediction and control in clustered populations. *J. Theor. Biol.* **272**(1), 1–7 (2011)
  151. House, T., Keeling, M.: Insights from unifying modern approximations to infections on networks. *J. R. Soc. Interface* **8**(54), 67–73 (2011)
  152. House, T., Davies, G., Danon, L., Keeling, M.J.: A motif-based approach to network epidemics. *Bull. Math. Biol.* **71**(7), 1693–1706 (2009)
  153. Isham, V., Medley, G.: Models for Infectious Human Diseases: Their Structure and Relation to Data, vol. 6. Cambridge University Press, Cambridge (1996)
  154. Jackson, M.O.: Social and Economic Networks, vol. 3. Princeton University Press, Princeton (2008)
  155. Jacobi, M.N., Görnerup, O.: A spectral method for aggregating variables in linear dynamical systems with application to cellular automata renormalization. *Adv. Complex Syst.* **12**(1–25) (2009)
  156. Janson, S., Luczak, M., Windridge, P.: Law of large numbers for the SIR epidemic on a random graph with given degrees. *Rand. Struct. Alg.* **45**(4), 726–763 (2014)
  157. Jones, P.W., Smith, P.: Stochastic Processes: An Introduction, 2nd edn. CRC Press, Boca Raton/FL (2012)
  158. Juher, D., Ripoll, J., Saldaña, J.: Outbreak analysis of an SIS epidemic model with rewiring. *J. Math. Biol.* **67**(2), 411–432 (2013)
  159. Juher, D., Kiss, I.Z., Saldaña, J.: Analysis of an epidemic model with awareness decay on regular random networks. *J. Theor. Biol.* **365**, 457–468 (2015)
  160. Kamke, E.: Zur theorie der systeme gewöhnlicher differentialgleichungen. II. *Acta Math.* **58**(1), 57–85 (1932)
  161. Karlin, S., Taylor, H.M.: A First Course in Stochastic Processes, 2nd edn. Academic Press, New York (1975)
  162. Karlin, S., Taylor, H.M.: A Second Course in Stochastic Processes. Academic Press, New York (1981)
  163. Karrer, B., Newman, M.E.J.: Message passing approach for general epidemic models. *Phys. Rev. E* **82**(1), 016101 (2010)
  164. Karrer, B., Newman, M.E.J.: Random graphs containing arbitrary distributions of subgraphs. *Phys. Rev. E* **82**(6), 066118 (2010)
  165. Keeling, M.J.: The ecology and evolution of spatial host-parasite systems. Ph.D. thesis, University of Warwick (1995)

166. Keeling, M.J.: The effects of local spatial structure on epidemiological invasions. *Proc. R. Soc. Lond. Ser. B: Biol. Sci.* **266**(1421), 859–867 (1999)
167. Keeling, M.J., Eames, K.T.D.: Networks and epidemic models. *J. R. Soc. Interface* **2**(4), 295–307 (2005)
168. Keeling, M.J., Rohani, P.: *Modeling Infectious Diseases in Humans and Animals*. Princeton University Press, Princeton (2008)
169. Keeling, M.J., Ross, J.V.: On methods for studying stochastic disease dynamics. *J. R. Soc. Interface* **5**(19), 171–181 (2008)
170. Keeling, M.J., Rand, D.A., Morris, A.J.: Correlation models for childhood epidemics. *Proc. R. Soc. Lond. Ser. B: Biol. Sci.* **264**(1385), 1149–1156 (1997)
171. Kemeny, J.G., Snell, J.L.: *Finite Markov Chains*, 2nd edn. Springer, New York (1976)
172. Kenah, E., Miller, J.C.: Epidemic percolation networks, epidemic outcomes, and interventions. *Interdiscip. Perspect. Infect. Dis.* (2011)
173. Kenah, E., Robins, J.M.: Second look at the spread of epidemics on networks. *Phys. Rev. E* **76**(3), 036113 (2007)
174. Kermack, W.O., McKendrick, A.G.: A contribution to the mathematical theory of epidemics. *R. Soc. Lond. Proc. Ser. A* **115**, 700–721 (1927)
175. Kirkwood, J.D.: Statistical mechanics of fluid mixtures. *J. Chem. Phys.* **3**, 300–313 (1935)
176. Kiss, I.Z., Simon, P.L.: New moment closures based on *a priori* distributions with applications to epidemic dynamics. *Bull. Math. Biol.* **74**(7), 1501–1515 (2012)
177. Kiss, I.Z., Green, D.M., Kao, R.R.: The effect of contact heterogeneity and multiple routes of transmission on final epidemic size. *Math. Biosci.* **203**(1), 124–136 (2006)
178. Kiss, I.Z., Green, D.M., Kao, R.R.: The network of sheep movements within Great Britain: network properties and their implications for infectious disease spread. *J. R. Soc. Interface* **3**(10), 669–677 (2006)
179. Kiss, I.Z., Simon, P.L., Kao, R.R.: A contact-network-based formulation of a preferential mixing model. *Bull. Math. Biol.* **71**(4), 888–905 (2009)
180. Kiss, I.Z., Berthouze, L., Taylor, T.J., Simon, P.L.: Modelling approaches for simple dynamic networks and applications to disease transmission models. *Proc. R. Soc. Lond. A: Math. Phys. Eng. Sci.* **468**(2141), 1332–1355 (2012)
181. Kiss, I.Z., et al.: Exact deterministic representation of Markovian SIR epidemics on networks with and without loops. *J. Math. Biol.* **70**(3), 437–464 (2015)
182. Kiss, I.Z., Röst, G., Vizi, Z.: Generalization of pairwise models to non-Markovian epidemics on networks. *Phys. Rev. Lett.* **115**(7), 078701 (2015)
183. Kivelä, M., Arenas, A., Barthélemy, M., Gleeson, J.P., Moreno, Y., Porter, M.A.: Multilayer networks. *J. Complex Netw.* **2**(3), 203–271 (2014)
184. Koch, D., Illner, R., Ma, J.: Edge removal in random contact networks and the basic reproduction number. *J. Math. Biol.* **67**(2), 217–238 (2013)

185. Kolmogorov, A.N.: The local structure of turbulence in incompressible viscous fluid for very large Reynolds numbers. *Dokl. Akad. Nauk SSSR* **30**, 299–303 (1941)
186. Krings, G., Karsai, M., Bernhardsson, S., Blondel, V.D., Saramäki, J.: Effects of time window size and placement on the structure of an aggregated communication network. *EPJ Data Sci.* **1**(4), 1–16 (2012)
187. Kuehn, C.: Moment closure—a brief review: control of self-organizing nonlinear systems. Springer, 253–271 (2016)
188. Kurant, M., Thiran, P.: Layered complex networks. *Phys. Rev. Lett.* **96**(13), 138701 (2006)
189. Kurtz, T.G.: Solutions of ordinary differential equations as limits of pure jump Markov processes. *J. Appl. Probab.* **7**(1), 49–58 (1970)
190. Kuulasmaa, K.: The spatial general epidemic and locally dependent random graphs. *J. Appl. Probab.* **19**(4), 745–758 (1982)
191. Kuulasmaa, K., Zachary, S.: On spatial general epidemics and bond percolation processes. *J. Appl. Probab.* **21**(4), 911–914 (1984)
192. Lajmanovich, A., Yorke, J.A.: A deterministic model for Gonorrhea in a non-homogeneous population. *Math. Biosci.* **28**(3), 221–236 (1976)
193. Lanchier, N., Neufer, J.: Stochastic dynamics on hypergraphs and the spatial majority rule model. *J. Stat. Phys.* 21–45 (2013)
194. Leone, M., Vazquez, A., Vespignani, A., Zecchina, R.: Ferromagnetic ordering in graphs with arbitrary degree distribution. *Eur. Phys. J. B* **28**, 191–197 (2002)
195. Leung, K.Y.: Dangerous connections: the spread of infectious diseases on dynamic networks. Ph.D. thesis, Utrecht University (2016)
196. Leung, K.Y., Kretzschmar, M., Diekmann, O.: *SI* infection on a dynamic partnership network: characterization of  $R_0$ . *J. Math. Biol.* **71**(1), 1–56 (2015)
197. Lindquist, J., Ma, J., van den Driessche, P., Willeboordse, F.H.: Effective degree network disease models. *J. Math. Biol.* **62**(2), 143–164 (2011)
198. Liu, Y.Y., Slotine, J.J., Barabási, A.L.: Controllability of complex networks. *Nature* **473**(7346), 167–173 (2011)
199. Lloyd, A.: Realistic distributions of infectious periods in epidemic models: changing patterns of persistence and dynamics. *Theor. Popul. Biol.* **60**, 59–71 (2001)
200. Lokhov, A.Y., Mézard, M., Ohta, H., Zdeborová, L.: Inferring the origin of an epidemic with a dynamic message-passing algorithm. *Phys. Rev. E* **90**(1), 012801 (2014)
201. Lovász, L.: Large Networks and Graph Limits, vol. 60. American Mathematical Society, Providence/Rhode Island (2012)
202. Ludwig, D.: Final size distributions for epidemics. *Math. Biosci.* **23**, 33–46 (1975)
203. Lusher, D., Koskinen, J., Robins, G.: Exponential Random Graph Models for Social Networks: Theory, Methods, and Applications. Cambridge University Press, Cambridge (2012)
204. Ma, J., Earn, D.J.D.: Generality of the final size formula for an epidemic of a newly invading infectious disease. *Bull. Math. Biol.* **68**(3), 679–702 (2006)

205. Marceau, V., Noël, P.A., Hébert-Dufresne, L., Allard, A., Dubé, L.J.: Adaptive networks: coevolution of disease and topology. *Phys. Rev. E* **82**(3), 036116 (2010)
206. Marceau, V., Noël, P.A., Hébert-Dufresne, L., Allard, A., Dubé, L.J.: Modeling the dynamical interaction between epidemics on overlay networks. *Phys. Rev. E* **84**(2), 026105 (2011)
207. Matsuda, H., Ogita, N., Sasaki, A., Satō, K.: Statistical mechanics of population the Lattice Lotka-Volterra Model. *Progr. Theor. Phys.* **88**(6), 1035–1049 (1992)
208. May, R.M., Anderson, R.M.: Transmission dynamics of HIV infection. *Nature* **326**, 137 (1987)
209. Melnik, S., Hackett, A., Porter, M.A., Mucha, P.J., Gleeson, J.P.: The unreasonable effectiveness of tree-based theory for networks with clustering. *Phys. Rev. E* **83**(3), 036112 (2011)
210. Meyers, L.A., Pourbohloul, B., Newman, M.E., Skowronski, D.M., Brunham, R.C.: Network theory and SARS: predicting outbreak diversity. *J. Theor. Biol.* **232**(1), 71–81 (2005)
211. Miller, J.C.: Epidemic size and probability in populations with heterogeneous infectivity and susceptibility. *Phys. Rev. E* **76**(1), 010101(R) (2007)
212. Miller, J.C.: Bounding the size and probability of epidemics on networks. *J. Appl. Probab.* **45**, 498–512 (2008)
213. Miller, J.C.: Percolation and epidemics in random clustered networks. *Phys. Rev. E* **80**(2), 020901(R) (2009)
214. Miller, J.C.: Spread of infectious disease through clustered populations. *J. R. Soc. Interface* **6**(41), 1121–1134 (2009)
215. Miller, J.C.: A note on a paper by Erik Volz: SIR dynamics in random networks. *J. Math. Biol.* **62**(3), 349–358 (2011)
216. Miller, J.C.: A note on the derivation of epidemic final sizes. *Bull. Math. Biol.* **74**(9), 2125–2141 (2012)
217. Miller, J.C.: Epidemics on networks with large initial conditions or changing structure. *PLoS ONE* **9**(7), e101421 (2014)
218. Miller, J.C.: Complex contagions and hybrid phase transitions. *J. Complex Netw.* **4**(2), 201–223 (2016)
219. Miller, J.C., Kiss, I.Z.: Epidemic spread in networks: Existing methods and current challenges. *Math. Modell. Nat. Phenom.* **9**(2), 4 (2014)
220. Miller, J.C., Volz, E.M.: Incorporating disease and population structure into models of SIR disease in contact networks. *PLoS ONE* **8**(8), e69162 (2013)
221. Miller, J.C., Volz, E.M.: Model hierarchies in edge-based compartmental modeling for infectious disease spread. *J. Math. Biol.* **67**(4), 869–899 (2013)
222. Miller, J.C., Slim, A.C., Volz, E.M.: Edge-based compartmental modelling for infectious disease spread. *J. R. Soc. Interface* **9**(70), 890–906 (2012)
223. Mollison, D.: *Epidemic Models: Their Structure and Relation to Data*, vol. 5. Cambridge University Press, Cambridge (1995)
224. Molloy, M., Reed, B.: A critical point for random graphs with a given degree sequence. *Rand. Struct. Alg.* **6**(2), 161–179 (1995)

225. Molloy, M., Reed, B.: The size of the giant component of a random graph with a given degree sequence. *Comb. Probab. Comput.* **7**(03), 295–305 (1998)
226. Moreno, Y., Pastor-Satorras, R., Vespignani, A.: Epidemic outbreaks in complex heterogeneous networks. *Eur. Phys. J. B: Condens. Matter Complex Syst.* **26**(4), 521–529 (2002)
227. Morris, A.J.: Representing spatial interactions in simple ecological models. Ph.D. thesis, University of Warwick (1997)
228. Müller, M.: Über das fundamentaltheorem in der theorie der gewöhnlichen differentialgleichungen. *Mathematische Zeitschrift* **26**(1), 619–645 (1927)
229. Nagy, N., Kiss, I.Z., Simon, P.L.: Approximate master equations for dynamical processes on graphs. *Math. Modell. Nat. Phenom.* **9**(02), 43–57 (2014)
230. Nåsell, I.: The quasi-stationary distribution of the closed endemic SIS model. *Adv. Appl. Probab.* **28**(03), 895–932 (1996)
231. Nåsell, I.: On the quasi-stationary distribution of the stochastic logistic epidemic. *Math. Biosci.* **156**(1), 21–40 (1999)
232. Nåsell, I.: Extinction and quasi-stationarity in the Verhulst logistic model. *J. Theor. Biol.* **211**(1), 11–27 (2001)
233. Neal, P.: Endemic behaviour of SIS epidemics with general infectious period distributions. *Adv. Appl. Probab.* **46**(1), 241–255 (2014)
234. Newman, M.E.J.: Spread of epidemic disease on networks. *Phys. Rev. E* **66**(1), 016128 (2002)
235. Newman, M.E.J.: Component sizes in networks with arbitrary degree distributions. *Phys. Rev. E* **76**(4), 045101 (2007)
236. Newman, M.E.J.: *Networks: An Introduction*. Oxford University Press, Oxford (2009)
237. Newman, M.E.J.: Random graphs with clustering. *Phys. Rev. Lett.* **103**(5), 058701 (2009)
238. Newman, M., Barabasi, A.L., Watts, D.J.: *The Structure and Dynamics of Networks*. Princeton University Press, Princeton (2006)
239. Noël, P.A., Allard, A., Hébert-Dufresne, L., Marceau, V., Dubé, L.J.: Propagation on networks: an exact alternative perspective. *Phys. Rev. E* **85**(3), 031118 (2012)
240. Noël, P.A., Davoudi, B., Brunham, R.C., Dubé, L.J., Pourbohloul, B.: Time evolution of disease spread on finite and infinite networks. *Phys. Rev. E* **79**(2), 026101 (2009)
241. Nold, A.: Heterogeneity in disease-transmission modeling. *Math. Biosci.* **52**(3), 227–240 (1980)
242. Nowzari, C., Preciado, V.M., Pappas, G.J.: Analysis and control of epidemics: a survey of spreading processes on complex networks. *IEEE Control Syst.* **36**(1), 26–46 (2016)
243. Pastor-Satorras, R., Vespignani, A.: Epidemic dynamics and endemic states in complex networks. *Phys. Rev. E* **63**(6), 066117 (2001)
244. Pastor-Satorras, R., Vespignani, A.: Epidemic spreading in scale-free networks. *Phys. Rev. Lett.* **86**, 3200–3203 (2001)

245. Pastor-Satorras, R., Vespignani, A.: Epidemic dynamics in finite size scale-free networks. *Phys. Rev. E* **65**(3), 035108 (2002)
246. Pastor-Satorras, R., Rubi, M., Diaz-Guilera, A.: *Statistical Mechanics of Complex Networks*, vol. 625. Springer Science & Business Media, Berlin/Heidelberg/New York (2003)
247. Pastor-Satorras, R., Castellano, C., Van Mieghem, P., Vespignani, A.: Epidemic processes in complex networks. *Rev. Mod. Phys.* **87**, 925 (2015)
248. Pellis, L., Ball, F., Bansal, S., Eames, K., House, T., Isham, V., Trapman, P.: Eight challenges for network epidemic models. *Epidemics* **10**, 58–62 (2015)
249. Pellis, L., House, T., Keeling, M.J.: Exact and approximate moment closures for non-Markovian network epidemics. *J. Theor. Biol.* **382**, 160–177 (2015)
250. Perko, L.: *Differential Equations and Dynamical Systems*, vol. 7. Springer Science & Business Media, New York (2001)
251. Perra, N., Gonçalves, B., Pastor-Satorras, R., Vespignani, A.: Activity driven modeling of time varying networks. *Sci. Rep.* **2**, Article No. 469, 1–7 (2012)
252. Picard, P.: Sur les modèles stochastique logistiques en démographie. *Ann. Inst. Henri Poincaré B* **II**, 151–172 (1965)
253. Porter, M.A., Gleeson, J.P.: *Dynamical Systems on Networks: A Tutorial*. Springer International Publishing, Heidelberg/New York (2016)
254. Pourbohloul, B., Brunham, R.C.: Network models and transmission of sexually transmitted diseases. *Sex. Transm. Dis.* **31**(6), 388–390 (2004)
255. Prakash, B.A., Chakrabarti, D., Valler, N.C., Faloutsos, M., Faloutsos, C.: Threshold conditions for arbitrary cascade models on arbitrary networks. *Knowl. Inf. Syst.* **33**(3), 549–575 (2012)
256. Rand, D.A.: Advanced ecological theory: principles and applications. In: *Correlation Equations and Pair Approximations for Spatial Ecologies*, pp. 100–142. Blackwell Science, Oxford (1999)
257. Rattana, P., Blyuss, K.B., Eames, K.T.D., Kiss, I.Z.: A class of pairwise models for epidemic dynamics on weighted networks. *Bull. Math. Biol.* **75**(3), 466–490 (2013)
258. Rattana, P., Miller, J.C., Kiss, I.Z.: Pairwise and edge-based models of epidemic dynamics on correlated weighted networks. *Math. Modell. Nat. Phenom.* **9**(02), 58–81 (2014)
259. Renshaw, E.: *Modelling Biological Populations in Space and Time*. Cambridge University Press, Cambridge (1991)
260. Ribeiro, B., Perra, N., Baronchelli, A.: Quantifying the effect of temporal resolution on time-varying networks. *Sci. Rep.* **3**, Article No. 3006, 1–5 (2013)
261. Riley, S., Fraser, C., Donnelly, C.A., Ghani, A.C., Abu-Raddad, L.J., Hedley, A.J., Leung, G.M., Ho, L.M., Lam, T.H., Thach, T.Q., Chau, P., Chan, K.P., Lo, S.V., Leung, P.Y., Tsang, T., Ho, W., Lee, K.H., Lau, E.M.C., Ferguson, N.M., Anderson, R.M.: Transmission dynamics of the etiological agent of SARS in Hong Kong: impact of public health interventions. *Science* **300**(5627), 1961–1966 (2003)
262. Risau-Gusmán, S., Zanette, D.H.: Contact switching as a control strategy for epidemic outbreaks. *J. Theor. Biol.* **257**(1), 52–60 (2009)

263. Risken, H.: *The Fokker-Planck Equation: Methods of Solution and Applications*. Springer Series in Synergetics. Springer, Berlin, Heidelberg (2012)
264. Ritchie, M., Berthouze, L., House, T., Kiss, I.Z.: Higher-order structure and epidemic dynamics in clustered networks. *J. Theor. Biol.* **348**, 21–32 (2014)
265. Ritchie, M., Berthouze, L., Kiss, I.Z.: Beyond clustering: mean-field dynamics on networks with arbitrary subgraph composition. *J. Math. Biol.* **72**(1–2), 255–281 (2016)
266. Ritchie, M., Berthouze, L., Kiss, I.Z.: Generation and analysis of networks with a prescribed degree sequence and subgraph family: higher-order structure matters. *J. Complex Netw.* (2016)
267. Rock, K., Brand, S., Moir, J., Keeling, M.J.: Dynamics of infectious diseases. *Rep. Progr. Phys.* **77**(2), 026602 (2014)
268. Rogers, L.C.G., Pitman, J.W.: Markov functions. *Ann. Probab.* **9**, 537–711 (1981)
269. Rogers, T., Clifford-Brown, W., Mills, C., Galla, T.: Stochastic oscillations of adaptive networks: application to epidemic modelling. *J. Stat. Mech.: Theor. Exp.* **2012**(08), P08018 (2012)
270. Röst, G., Vizi, Z., Kiss, I.Z.: Impact of non-Markovian recovery on network epidemics. In: *Biomat 2015: Proceedings of the International Symposium on Mathematical and Computational Biology*. World Scientific, New York (2015)
271. Röst, G., Vizi, Z., Kiss, I.Z.: Pairwise approximation for SIR type network epidemics with non-Markovian recovery. *arXiv preprint arXiv:1605.02933* (2016)
272. Salinelli, E., Tomarelli, F.: *Discrete Dynamical Models*, vol. 76. Springer, Heidelberg/New York (2014)
273. Saramäki, J., Kaski, K.: Modelling development of epidemics with dynamic small-world networks. *J. Theor. Biol.* **234**(3), 413–421 (2005)
274. Saumell-Mendiola, A., Serrano, M.Á., Boguñá, M.: Epidemic spreading on interconnected networks. *Phys. Rev. E* **86**(2), 026106 (2012)
275. Scott, M.: *Applied stochastic processes in science and engineering*. University of Waterloo, eBook (2013)
276. Sedgewick, R.: *Algorithms in C, Part 5: Graph Algorithms*. Addison-Wesley (2002)
277. Sélley, F., Besenyei, Á., Kiss, I.Z., Simon, P.L.: Dynamic control of modern, network-based epidemic models. *SIAM J. Appl. Dyn. Syst.* **14**(1), 168–187 (2015)
278. Sellke, T.: On the asymptotic distribution of the size of a stochastic epidemic. *J. Appl. Probab.* **20**(02), 390–394 (1983)
279. Sharkey, K.J.: Deterministic epidemiological models at the individual level. *J. Math. Biol.* **57**, 311–331 (2008)
280. Sharkey, K.J.: Deterministic epidemic models on contact networks: correlations and unbiological terms. *Theor. Popul. Biol.* **79**, 115–129 (2011)
281. Sharkey, K.J., Wilkinson, R.R.: Complete hierarchies of SIR models on arbitrary networks with exact and approximate moment closure. *Math. Biosci.* **264**, 74–85 (2015)



282. Sharkey, K.J., Fernandez, C., Morgan, K.L., Peeler, E., Thrush, M., Turnbull, J.F., Bowers, R.G.: Pair-level approximations to the spatio-temporal dynamics of epidemics on asymmetric contact networks. *J. Math. Biol.* **53**(1), 61–85 (2006)
283. Sharkey, K.J., et al.: Exact equations for SIR epidemics on tree graphs. *Bull. Math. Biol.* **77**(4), 614–645 (2015)
284. Sherborne, N., Blyuss, K.B., Kiss, I.Z.: Dynamics of multi-stage infections on networks. *Bull. Math. Biol.* **77**(10), 1909–1933 (2015)
285. Shkarayev, M.S., Tunc, I., Shaw, L.B.: Epidemics with temporary link deactivation in scale-free networks. *J. Phys. A: Math. Theor.* **47**(45), 455006 (2014)
286. Simon, P.L., Kiss, I.Z.: From exact stochastic to mean-field ODE models: a new approach to prove convergence results. *IMA J. Appl. Math.* **78**(5), 945–964 (2013)
287. Simon, P.L., Kiss, I.Z.: Super compact pairwise model for SIS epidemic on heterogeneous networks. *J. Complex Netw.* **4**(2), 187–200 (2016)
288. Simon, P.L., Taylor, M., Kiss, I.Z.: Exact epidemic models on graphs using graph-automorphism driven lumping. *J. Math. Biol.* **62**(4), 479–508 (2011)
289. Skiena, S.S.: *The Algorithm Design Manual*, 2nd edn. Springer Science & Business Media, New York (2009)
290. Smith, H.L.: *Monotone Dynamical Systems: An Introduction to the Theory of Competitive and Cooperative Systems*, vol. 41. American Mathematical Society, Providence/Rhode Island (2008)
291. Sood, V., Redner, S.: Voter model on heterogeneous graphs. *Phys. Rev. Lett.* **94**(17), 178701-1–178701-4 (2005)
292. Startsev, A.N.: On the distribution of the size of an epidemic in a non-Markovian model. *Theor. Probab. Appl.* **41**(4), 730–740 (1997)
293. Startsev, A.N.: Asymptotic analysis of the general stochastic epidemic with variable infectious periods. *J. Appl. Probab.* **38**(01), 18–35 (2001)
294. Szabó, A., Simon, P.L., Kiss, I.Z.: Detailed study of bifurcations in an epidemic model on a dynamic network. *Differ. Equ. Appl.* **4**, 277–296 (2012)
295. Szabó-Solticzky, A., et al.: Oscillating epidemics in a dynamic network model: stochastic and mean-field analysis. *J. Math. Biol.* **72**(5), 1153–1176 (2016)
296. Szarski, J.: *Differential inequalities*. Instytut Matematyczny Polskiej Akademii Nauk (Warszawa) (1965)
297. Tarjan, R.: Depth-first search and linear graph algorithms. *SIAM J. Comput.* **1**(2), 146–160 (1972)
298. Taylor, T.J., Kiss, I.Z.: Interdependency and hierarchy of exact and approximate epidemic models on networks. *J. Math. Biol.* **69**(1), 183–211 (2014)
299. Taylor, M., Simon, P.L., Green, D.M., House, T., Kiss, I.Z.: From Markovian to pairwise epidemic models and the performance of moment closure approximations. *J. Math. Biol.* **64**(6), 1021–1042 (2012)
300. Taylor, M., Taylor, T.J., Kiss, I.Z.: Epidemic threshold and control in a dynamic network. *Phys. Rev. E* **85**, 016103 (2012)
301. Trapman, P.: On analytical approaches to epidemics on networks. *Theor. Popul. Biol.* **71**(2), 160–173 (2007)

302. Trapman, P.: Reproduction numbers for epidemics on networks using pair approximation. *Math. Biosci.* **210**(2), 464–489 (2007)
303. Traud, A.L., Kelsic, E.D., Mucha, P.J., Porter, M.A.: Comparing community structure to characteristics in online collegiate social networks. *SIAM Rev.* **53**(3), 526–543 (2011)
304. Tunc, I., Shkarayev, M.S., Shaw, L.B.: Epidemics in adaptive social networks with temporary link deactivation. *J. Stat. Phys.* **151**(1–2), 355–366 (2013)
305. Valdez, L.D., Macri, P.A., Braunstein, L.A.: Temporal percolation of the susceptible network in an epidemic spreading. *PLoS ONE* **7**(9), e44188 (2012)
306. van Baalen, M.: Pair approximations for different spatial geometries. In: *The Geometry of Ecological Interactions: Simplifying Spatial Complexity*, pp. 359–387. Cambridge University Press, Cambridge (2000)
307. van de Bovenkamp, R., Van Mieghem, P.: Survival time of the susceptible-infected-susceptible infection process on a graph. *Phys. Rev. E* **92**(3), 032806 (2015)
308. van den Driessche, P., Watmough, J.: Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission. *Math. Biosci.* **180**(1), 29–48 (2002)
309. van Kampen, N.G.: *Stochastic Processes in Physics and Chemistry*, vol. 1. Elsevier, Amsterdam (1992)
310. Van Mieghem, P.: The n-intertwined SIS epidemic network model. *Computing* **93**(2–4), 147–169 (2011)
311. Van Mieghem, P., van de Bovenkamp, R.: Non-Markovian infection spread dramatically alters the susceptible-infected-susceptible epidemic threshold in networks. *Phys. Rev. Lett.* **110**(10), 108701 (2013)
312. Van Mieghem, P., Omic, J., Kooij, R.: Virus spread in networks. *IEEE/ACM Trans. Netw.* **17**(1), 1–14 (2009)
313. Van Mieghem, P., Sahneh, F.D., Scoglio, C.: An upper bound for the epidemic threshold in exact Markovian SIR and SIS epidemics on networks. In: *2014 IEEE 53rd Annual Conference on Decision and Control (CDC)*, pp. 6228–6233. IEEE (2014)
314. Vazquez, F., Eguíluz, V.M.: Analytical solution of the voter model on uncorrelated networks. *New J. Phys.* **10**(6), 063011, 1–19 (2008)
315. Volz, E.M.: Random networks with tunable degree distribution and clustering. *Phys. Rev. E* **70**(5), 056115 (2004)
316. Volz, E.M.: SIR dynamics in random networks with heterogeneous connectivity. *J. Math. Biol.* **56**(3), 293–310 (2008)
317. Volz, E.M., Meyers, L.A.: Susceptible–infected–recovered epidemics in dynamic contact networks. *Proc. R. Soc. Lond. B: Biol. Sci.* **274**(1628), 2925–2934 (2007)
318. Volz, E.M., Miller, J.C., Galvani, A., Meyers, L.A.: Effects of heterogeneous and clustered contact patterns on infectious disease dynamics. *PLoS Comput. Biol.* **7**(6), e1002042 (2011)

319. Wallinga, J., Lipsitch, M.: How generation intervals shape the relationship between growth rates and reproductive numbers. *Proc. R. Soc. Lond. B: Biol. Sci.* **274**(1609), 599–604 (2007)
320. Wang, H., Li, Q., D’Agostino, G., Havlin, S., Stanley, H.E., Van Mieghem, P.: Effect of the interconnected network structure on the epidemic threshold. *Phys. Rev. E* **88**(2), 022801 (2013)
321. Wang, W., Tang, M., Zhang, H.F., Gao, H., Do, Y., Liu, Z.H.: Epidemic spreading on complex networks with general degree and weight distributions. *Phys. Rev. E* **90**(4), 042803 (2014)
322. Wang, W., Liu, Q.H., Zhong, L.F., Tang, M., Gao, H., Stanley, H.E.: Predicting the epidemic threshold of the susceptible-infected-recovered model. *Sci. Rep.* **6**, 24676, 1–12 (2016)
323. Watts, D.J.: Networks, dynamics, and the small-world phenomenon 1. *Am. J. Soc.* **105**(2), 493–527 (1999)
324. Watts, D.J., Strogatz, S.H.: Collective dynamics of ‘small-world’ networks. *Nature* **393**(6684), 440–442 (1998)
325. Wearing, H.J., Rohani, P., Keeling, M.J.: Appropriate models for the management of infectious diseases. *PLoS Med.* **2**(7), 621 (2005)
326. Wilkinson, R.R., Sharkey, K.J.: An exact relationship between invasion probability and endemic prevalence for Markovian SIS dynamics on networks. *PLoS ONE* **8**(7), e69028 (2013)
327. Wilkinson, R.R., Sharkey, K.J.: Message passing and moment closure for susceptible-infected-recovered epidemics on finite networks. *Phys. Rev. E* **89**(2), 022808-1–022808-6 (2014)
328. Wilkinson, R.R., Ball, F.G., Sharkey, K.J.: The relationships between message passing, pairwise, Kermack-McKendrick and stochastic SIR epidemic models. *arXiv preprint arXiv:1605.03555* (2016)
329. Yan, G., Tsekenis, G., Barzel, B., Slotine, J.J., Liu, Y.Y., Barabási, A.L.: Spectrum of controlling and observing complex networks. *Nat. Phys.* **11**(9), 779–786 (2015)
330. Yap, H.P.: Some Topics in Graph Theory. London Mathematical Society, Lecture Notes, vol. 108. Cambridge University Press, Cambridge (1986)
331. Youssef, M., Scoglio, C.: Mitigation of epidemics in contact networks through optimal contact adaptation. *Math. Biosci. Eng.* **10**(4), 1227–1251 (2013)

# Index

## A

$\mathcal{A}$ , *see* attack rate, *see* final epidemic size  
 adjacency matrix, 14, 73  
 annealed network, 170  
 approximate master equations, 195  
 approximation error, 143, 363  
 assortative, 170  
 assortative mixing, 16  
 attack rate, *see* final epidemic size, 207,  
     210–215  
     Configuration Model networks, 239–243  
     definition, 213  
 automorphism, 51  
 average degree, 16, 166

## B

Barabási–Albert networks, 374–377  
 basic reproductive ratio, 6, 7, 9, 136, 140, 163,  
     219, 222, 305, 314  
     growth-based, 136–140  
     heterogeneous mean-field at single level,  
     180  
     non-Markovian pairwise, 319  
 bimodal random network, 16, 19, 114, 147,  
     175, 179, 184, 186  
 binary dynamics, 32, 155, 329  
 birth-death process, 327  
 bond percolation, 226  
 bottom-up model, 73  
     exact, SIR, 74  
     exact, SIS, 73  
 bow-tie, 235  
 bridge, 102

## C

class, 44  
     induced, 44  
 closure, 67, 70–73  
     binomial, 143, 349  
     clustered, 127  
     exact, 101  
     pair, 23, 72, 123, 143, 168, 169  
     triple, 23, 72, 101, 124, 143, 168, 170, 173,  
     197  
 clustering, 17  
 clustering coefficient, 17, 126  
     average, 368  
 cobweb diagrams, 218  
 compact pairwise  
     model, 24  
     SIS, analytical results, 183, 192  
 compact pairwise equations  
     SIR, 176, 257  
     SIS, 175  
 compact pairwise model, 174–176  
 compartmental model, 5  
     SIR, 7  
     SIR extended, 305  
     SIR with fixed recovery time, 314  
     SIS, 6  
 complete network, 94, 141  
 complex contagion, 253  
 Configuration Model, 18  
     Generation algorithm, 19  
 connected, 18  
 connected component  
     definition, 226

conservation relation, 158, 182, 192  
 cooperative, 76  
 correlation, 71, 133, 161  
 critical value of  $\tau$ , 145, 184  
   compact pairwise, 182, 193  
   heterogeneous mean-field at single level, 180, 189  
   preferential mixing, 205  
 cut-vertex, 102, 104  
 cycle, 18

**D**

degree, 16  
 degree distribution, 16, 165  
   second moment, 166  
 degree heterogeneity, 24, 183, 186, 295  
 degree-based mean-field model, 204  
 delay differential equation, 315  
 density dependent, 327  
 differential inequality, 75  
 directed graph, 14  
 disassortative, 170  
 disassortative mixing, 16  
 disease-free steady state, 81  
   bottom up, 85, 86  
   compact pairwise, 182, 193  
   compartmental model, 6  
   heterogeneous mean-field at single level, 179, 189  
   homogeneous mean-field at single level, 129  
   homogeneous pairwise, 133, 159  
   link-status dependent rewiring, 288  
   random link activation and deletion, 281  
 distribution  
   exponential, 233

**E**

EBCM, *see* edge-based compartmental model  
 ED, *see* effective degree  
 edge, 13  
 edge-based compartmental model, 24, 243–251, 257  
   dynamic network, 296–300  
 effective degree model, 24, 195–204  
   basic, 196  
   compact, 200  
   compact, SIR, 203, 257  
   compact, SIS, 204  
   contact preserving rewiring, 279  
   random link activation and deletion, 284  
   SIR, 199  
   SIS, 198  
 endemic steady state  
   bottom up, 81, 86, 88

compact pairwise, 183, 194  
 compartmental model, 6  
 contact preserving rewiring, 276  
 heterogeneous mean-field at single level, 179, 190  
 homogeneous mean-field at single level, 129, 130  
 homogeneous pairwise, 133, 159  
 link-status dependent rewiring, 289  
 random link activation and deletion, 281  
 rewiring on fixed networks, 295  
 epidemic, 209  
   definition, 210, 213  
 epidemic probability, 9, 207, 210–215  
   Configuration Model networks, 215–226  
   definition, 213  
 epidemic threshold, *see* critical value of  $\tau$ , 209, 218, 219  
   random link activation and deletion, 286  
 Erdős-Rényi random graph, 15, 19, 114, 145, 149, 184, 185  
 excess degree, 266

**F**

Facebook, 367  
 final epidemic size, 140, 210–215  
   bottom up, 99  
   deterministic compartmental model, 7  
   extended compartmental model, 305  
   extended pairwise model, 310  
   heterogeneous mean-field at single level, 181, 191  
   homogeneous mean-field equations at single level, 131  
   homogeneous pairwise, 135  
   non-Markovian pairwise, 317  
   stochastic compartmental model, 10  
 fixed point, 218  
   theorem, Brouwer, 82  
 Fokker-Planck equation, 25  
   boundary condition, 335  
   density dependent case, 336  
   derivation, 334  
   Gaussian approximation of steady state, 358  
   general definition, 333  
   one-step process with linear coefficients, 338  
   random walk, 341  
   solving by Fourier method, 357  
   steady state, 358

**G**

Galton-Watson, 216  
 $\gamma$ , *see* recovery rate

giant component, 227  
 giant strongly connected component, 235  
 Gillespie algorithm, 382  
 global behaviour, 160, 182  
 growth-based basic reproductive ratio  
   compact pairwise, 182

**H**

heterogeneous susceptibility/infectiousness,  
   224  
 HetMF, *see* mean field, heterogeneous  
 HomMF, *see* mean field, homogeneous  
 homogeneous graph, 219  
 homogeneous susceptibility/infectiousness,  
   224

**I**

in-component, 234  
 individual-based model, 78, 91  
   mean-field, 23  
   pairwise, 23  
   SIR, 98  
   SIS, 78  
 infection rate, *see* transmission rate  
 initial condition, 126  
 Ising model, 32

**K**

Kamke-Müller condition, 75  
 Kolmogorov equations, *see* master equations

**L**

line graph, 13  
 linear, 155  
 linked list, 15  
 lollipop network, 102  
 lumpable, 47  
 lumping, 30, 42–62  
   by equivalence classes, 53  
   recipe, 54

**M**

Malthusian parameter, 137  
 Markov chains, 8, 30–31  
   continuous-time, 327  
 master equations, 9, 23, 30–42, 121, 140, 152,  
   327  
 mean-field  
   exact at pair level, 123, 167  
   exact at single level, 121, 166  
   heterogeneous networks, 165–205  
   homogeneous networks, 117–164  
 mean-field at single level  
   SIR heterogeneous, 172

SIR heterogeneous, analytical results, 182  
 SIR homogeneous, 125  
 SIR homogeneous, analytical results, 131  
 SIS heterogeneous, 171  
 SIS heterogeneous, analytical results, 181  
 SIS homogeneous, 124  
 SIS homogeneous, analytical results, 130  
 mean-field model, 23, 117  
   heterogeneous, 24  
 message passing, 239, 253  
 moments of a probability distribution, 332

**N**

network  
   directed, 73–89, 164  
   multiplex, 205  
   weighted, 15, 73–89, 164  
 next generation matrix, 140  
 node, 13  
 non-Markovian epidemics, 24

**O**

one-step process, 25, 327  
 orbit, 52  
 oscillation, 24  
   contact preserving rewiring, 277  
   link-status dependent rewiring, 290, 291  
 oscillatory cycle in adaptive networks, 291  
 out-component, 234  
 outbreak  
   epidemic, 209  
   small, 209

**P**

$\mathcal{P}$ , *see* epidemic probability  
 pair  
   closure, 23, 72, 123, 143, 169  
   conservation, 132  
   count, 119, 166  
 pair-based model  
   SIR, 112  
   SIS, 92  
 pairwise equations  
   contact preserving rewiring, 275  
   link-status dependent activation and  
     deletion, 287  
   random link activation and deletion, 280  
   rewiring on fixed networks, 294  
   SIR heterogeneous, 174  
   SIR homogeneous, 126  
   SIR homogeneous, analytical results, 136  
   SIS heterogeneous, 173  
   SIS homogeneous, 125  
   SIS homogeneous, analytical results, 134

pairwise model, 23  
     heterogeneous, 24  
     SIR extended, 305  
     SIR with fixed recovery time, 316  
 partial differential equation, 25  
 partition, 44  
     induced, 44  
     refinement, 49  
 path, 18  
 percolated network  
     definition, 226  
 percolation, 24, *see* bond percolation  
     directed, 231–239  
 pgf, *see* probability generating function  
 phase plane analysis, 159  
 phase portrait, 130  
 $P_n(k)$ , 19, 215  
 power law degree distribution, 20, 95, 114,  
     147, 184, 185  
 preferential attachment, 374  
 preferential mixing, 16, 170, 173  
 probability generating function, 25, 215  
     definition for one-step processes, 332  
     one-step processes with linear coefficients,  
         337  
     one-step processes with polynomial  
         coefficients, 351  
     random walk, 341  
 $\psi$ , *see* probability generating function  
 $\hat{\psi}$ , 172, 244  
 PW, *see* pairwise

## Q

$q(G, p, d)$ , 214  
 $Q(\mathcal{G}, p, d, N)$ , 214  
 quadruple, 128  
 quasi-steady state, 143  
 quenched network, 170  
 quiescent-active-quiescent model, 32

## R

$R_0$ , *see* basic reproductive ratio  
 $\bar{R}_0$ , *see* basic reproductive ratio, growth-based  
 random graph, 15  
 random regular network, 19, 114, 145, 149  
 recovery rate, 21, 29  
 refinement, 49  
 regular graph, 16, 219  
 rewiring, 24  
     contact preserving, 274  
     link-status dependent, 274  
     on a fixed network, 274  
     random link activation and deletion, 274  
 rumour spread, 155

## S

saddle-node bifurcation  
     contact preserving rewiring, 276  
 scale-free random network, 20  
 simulation  
     continuous-time, 381  
     efficient algorithms, 381–386  
     event-based, 384  
     Gillespie, 382  
     individual-based stochastic, 22, 145, 183,  
         285  
     individual-based stochastic non-Markovian,  
         316  
     SIR, 385  
     SIS, 386  
     stochastic, 141  
 SIR, 5  
 SIS, 5  
 small outbreak  
     definition, 210  
 small-world networks, 371–373  
 star network, 94, 95, 141, 144  
 state, 27  
 state space, 27, 30  
 strongly connected, 18  
 super compact pairwise model, 24, 176–179  
     SIR, 178  
     SIS, 177  
 survivor function, 308

## T

$\tau$ , *see* transmission rate  
 test node, 239  
 time shifting, 387–388  
 toast network, 15  
 transcritical bifurcation  
     backward, 277, 295  
     bottom up, 86  
     compact pairwise, 183, 194  
     contact preserving rewiring, 276  
     heterogeneous mean-field at single level,  
         180, 189  
     homogeneous mean-field at single level,  
         129  
     homogeneous pairwise, 134, 159  
     random rewiring, 281  
     rewiring on fixed networks, 295  
 transition, 154  
 transition function  
     linear, 155, 331  
 transmissibility, 307  
 transmission probability  
     average, 223

- in, 223
- out, 223
- transmission rate, 21, 29
- tree network, 18, 109, 112
- triple
  - closure, 23, 72, 124, 143, 170, 173, 197
  - clustered closure, 127
  - count, 119
  - number of, 127

**U**

- un-directed graph, 14

**V**

- vertex, 13
- voter model, 32, 155, 330

**W**

- Watts–Strogatz networks, 371–373
- weighted graph, *see* network, weighted

**X**

- $x_{G,p}(m)$ , 208

**Y**

- $y_{G,p}(\rho)$ , 213