



# Finite Element Approach to Photoemission of Complex Molecules

Tobias Möhle

December 12, 2016

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Calculation of Photoelectron spectra</b>	<b>6</b>
2.1	The Green's Function Approach . . . . .	8
2.2	Combined Bound and Continuum State Representation . . . . .	9
2.3	The Dyson Orbital Formalism . . . . .	11
2.3.1	Time-dependent Dyson Orbitals . . . . .	12
2.3.2	Time-independent Dyson Orbitals . . . . .	12
2.3.3	Sudden Approximation . . . . .	14
<b>3</b>	<b>Description of Free Particles</b>	<b>16</b>
3.1	Continuum Waves . . . . .	16
3.1.1	Stieltjes Imaging . . . . .	18
3.2	Finite Differences and Finite Volumes . . . . .	19
3.3	Pseudospectral Methods . . . . .	21
3.4	Radial Basis Functions . . . . .	22
3.5	Finite Elements . . . . .	22
3.6	Hybrid Methods . . . . .	25
3.7	Wavelets . . . . .	25
<b>4</b>	<b>Finite Element Methods</b>	<b>27</b>
4.1	Finite Element Calculations in Quantum Chemistry . . . . .	27
4.2	Integration of Matrix Elements and Formulations of the Equation System .	28
4.3	Element Types and Mesh Types . . . . .	28
4.4	Boundary Conditions . . . . .	30
4.4.1	Complex Absorbing Potential . . . . .	31
4.4.2	Mode-matching Schemes . . . . .	31
4.4.3	The Boundary Element Method . . . . .	32
4.4.4	Infinite Elements . . . . .	32
4.5	Solving Large Eigenvalue Problems . . . . .	34
4.5.1	Generalised Eigenproblem . . . . .	35
4.5.2	Stabilisation of Eigenproblems . . . . .	35
4.5.3	Solving Large Eigenproblems . . . . .	37
<b>5</b>	<b>Applied Protocol</b>	<b>38</b>
5.1	Bound State Functions . . . . .	38
5.1.1	Ground State Density . . . . .	38

5.1.2	Excited State Properties . . . . .	40
5.1.3	OTRSH-scheme . . . . .	40
5.1.4	Computing the Dyson orbital . . . . .	41
5.2	Free electron function . . . . .	41
5.2.1	Setup of the Grid . . . . .	41
5.2.2	Obtaining ESP . . . . .	44
5.3	Obtaining the DOs . . . . .	44
<b>6</b>	<b>Results</b>	<b>46</b>
6.1	Benchmarks and Tests on the Formalism . . . . .	46
6.1.1	Comparison of Formulations . . . . .	46
6.1.2	Numerical Convergence . . . . .	47
6.2	atomic Lithium . . . . .	49
6.3	Triatomic Linear Molecules . . . . .	49
6.4	water . . . . .	49
<b>7</b>	<b>Resumee</b>	<b>51</b>
<b>A</b>	<b>Appendix</b>	<b>70</b>
A.1	Additional Data . . . . .	70
A.2	Delaunay Triangulation . . . . .	70

## Notation:

- $\mathbf{r}$  vector in 3 dimensions
- $\mathbb{R}$  General matrix
- $|\Psi\rangle$  one-particle wave function
- $|\Phi^{\text{DO}}\rangle$  Dyson orbital
- $|\Phi^{\text{el}}\rangle$  Free electron function
- $|\Psi^N\rangle$   $N$ -electron wave function
- $\hat{a}/\hat{a}^\dagger$  annihilation/creation operators:..

## Abbreviations:

- CAP complex absorbing potential
- CI configuration interaction
- DO Dyson orbital
- DFT density functional theory
- DVR discrete variable approximation
- eV electron volt
- FD finite difference
- FEF free electron function
- FEM finite element method
- FE-DVR finite element discrete variable approximation
- FV finite volume
- GASCI
- LCAO linear combination of atomic orbitals
- LUMO lowest unoccupied molecular orbital
- RBF radial basis function
- SCF self-consistent field
- SD Slater determinant
- SA sudden approximation
- SE Schrödinger equation
- TDDFT time dependent density functional theory

# 1 | Introduction

Steady state as well as time-resolved photoelectron spectroscopy have become widely used tools to study the composition of gases and liquids [1–3], the structure of solids [4] as well as chemical reactions such as electron transfer [3]. The process studied by photoelectron spectra is the absorption of a high-energetic photon (usually in the ultra-violet to hard X-ray regime) by an  $N$ -electron system which leads to the release of an electron whose kinetic energy is measured. This process is sketched in Figure 1.1 where the bound states are depicted by horizontal bars and the states of the outgoing electron are visualised by the gradient-filled block, indicating that its energy is continuous. Measuring the kinetic energy of the photoelectron yields information on the energy of its previous states due to energy-conservation over the process.

One of the main reasons for the broad success of this particular type of spectroscopy is that these spectra are very sensitive to small changes in the chemical environment and hence yield information not only about the chemical structure of molecules but also about intermolecular interactions as for example solvation effects [1, 5–8]. Furthermore, photoelectron spectroscopy provides a more direct access to the energy levels than optical absorption and emission spectroscopy since the transition energies are obtained with respect to the vacuum level and no “dark” states occur due to different selection rules. Finally the ease of handling charged particles experimentally makes this method appealing and a good temporal resolution can be achieved by varying the path in the time of flight spectrometer. Moreover, besides its capabilities in steady state spectroscopy, photoelectron spectra are the standard tool to study attosecond physics due to the naturally high energy of the short pulses which ionise the systems under




Figure 1.1: Schematic representation of a photoelectron transition : an incoming photon with energy  $h\nu$  ionises the  $N$ -electron system, transferring it into a free electron in a continuum state and a system of  $N - 1$  bound electrons. The bars denote bound states, the color-gradient depicts the energy of the outgoing electron.

study [9–14]. However, a limitation of photoelectron spectroscopy is that the short free-path of electrons in condensed phases limits the probe depth significantly.

Having performed a measurement of a photoelectron spectrum (PES) it is, at least for complex systems, rich of features and hence the interpretation requires theoretical methods. This especially applies to systems with strong electron correlation which manifests itself in the appearance of combination transitions. Over the decades a large variety of methods to calculate photoelectron spectra have been developed at different levels of theory. In literature often the theoretical spectra are estimated on the basis of Koopmans' theorem [15], (or its density functional theory (DFT) counterpart [16],) assuming equal intensities for all transitions [17–20]. Even though this is a quite successful approach for solid states [21, 22] and gives an easy interpretation, it is too simplistic in many cases since it neglects electron relaxation and correlation effects and hence may give an invalid picture as shown by Cederbaum *et al.* even for small systems such as various diatomics [23, 24]. Furthermore in this model no reliable information about intensities can be obtained. To retrieve quantitative transition strengths, more advanced models are needed. Since simulations in time-domain are very demanding they are only applied to small molecules, mainly to study strong-field effects such as high harmonic generation [10, 25–28] which can not be described in frequency-domain. In this thesis however the focus is on complex molecular systems and a wide range of kinetic energies, while more moderate field strengths should be applied. Hence frequency-domain methods derived from a perturbation theory with respect to the irradiating electromagnetic field are applicable here.

In this thesis, the PESs are calculated within the Dyson orbital (DO) formalism which is derived and explained in more detail. This formalism is based on Fermis' Golden Rule [29] and allows a reduction of the dipole moment matrix element from an  $N$ -electron integral to effective one-particle quantities. Thereby the initial state and the bound part of the final state are represented by a one-electron quantity denoted as DO. The other wave function entering the dipole moment operator is the free electron function (FEF). In this formalism electron relaxation and correlation effects between bound states of the unionised and ionised systems are included, allowing the description of combination transitions. Per contra, this formalism neglects the correlation of the outgoing electron with the ionic remainder and thus may neglect certain transitions [30].

Since the bound state wave functions can be obtained from standard quantum-chemical tools, the computation of the DO is straightforward even though technically demanding [31]. The computation of the FEF is in general not trivial since analytic solutions are known only for few special cases such as hydrogen-like atoms [32] and general basis sets as they are used for bound states are not available. Among others, three analytic expressions have been suggested, each based on an expansion of spherically symmetric functions in plane waves [33]. One of them is the spherical wave basis which is a set of solutions of the Schrödinger equation (SE) without any potential and hence is especially well-suited for photodetachment from negative ions, leaving systems in an uncharged state [33, 34]. The other two expansions are based on Coulomb waves which assume a Coulomb potential and hence are exact for the ionisation of hydrogen-like atoms [32]. Thereby one of them is an expansion in Coulomb waves for the given momentum vector  $\mathbf{k}$  while the other is an asymptotic expression for Coulomb waves with vanishing momentum (Coulomb  $|\mathbf{k}| = 0$ ).

The functions each are expanded in a series of increasing quantum numbers for angular momentum  $l$  and its projection  $m$ . Since all three functions assume spherical symmetry of the potential, they are expected to give a good approximation to the real FEF for spherically symmetric systems only. Ionisation *e. g.* from a delocalised orbital of a linear molecule have a different symmetry and thus can be expected to be described only crudely by this approach. The effects due to lower molecular symmetry become especially important for low kinetic energies because the interaction with the ionic remainder becomes strong in this case.

To analyse the quality of a given expression for a FEF, it is instructive to study the intensity of a given transition as a function of kinetic energy. Such a study is examined in Figure 1.2, showing the intensity of the photoionisation transition from the highest occupied molecular orbital (HOMO) of water for different kinetic energies of the photo-electron, estimated with the above-mentioned expansions. The comparison shows that all three expansions yield different intensities for most transition energies. Besides leading a wrong wrong symmetry of the FEF, these expansions yiled even in the limit of infinite terms only a plane-wave, neglecting the ionic potential and hence are not asymptotically correct.

To overcome the restrictions in symmetry and kinetic energy of the photoelectron function, an explicit formulation is needed, taking the molecular electrostatic potential experienced by the outgoing electron into account. Assuming that the correlation between the FEF and the bound states of the ion is weak (which is a prerequisite for the DO formalism anyway), one can use the mean-field potential of the molecular remainder. This allows for obtaining the FEF from the one-electron SE with an appropriate potential and thus reduces the complexity compared to the exact case where a coupled  $N$ -electron equation needs to be solved, considerably.

In the thesis at hand a good approximation to the FEF is aimed which should be applicable to a wide range of molecules and photon energies. Such a flexible description is possible exploiting the finite element method for solving the SE. In the finite element method, the space of interest is subdivided into small volume elements and solved variationally with stepwise polynomials whose support spans over one or few elements only [35, 36]. The finite element description is especially efficient here since the size of the elements




Figure 1.2: The intensity of the lowest-lying transition of water depending on the kinetic energy of the photoelectron. Each expansion is performed up to  $l = 10$ .

can be locally adapted to reproduce finer structures or broader shapes [35, 37]. Using finite elements, the one-particle SE is formulated a generalised eigenvalue problem with sparse matrices.

An important characteristic of the exact FEF is its spacially infinite extend which can not be handled by the finite element method. To describe the function in a good approximation, the finite elements are extended by infinite elements [38–41] which are volume elements with infinite radial extend that are connected to the outer surface of the finite element region. Thereby the radial function is a polynomial in  $\frac{1}{r}$  multiplied by  $e^{ikr}$ , resembling the asymptotic behaviour of known analytic solutions and fulfilling the Sommerfeld radiation condition [42].

The bound states can be obtained on the level of density functional theory (DFT) which is a formally exact method based on the Hohenberg-Kohn theorem [43], stating that the total electron density determines all system properties such as electronic binding energies. However, the correct form of the exchange potential as well as the kinetic energy are not known as functionals of the electron density and hence approximate functionals need to be used. While the kinetic energy can be estimated reasonably by the Kohn-Sham scheme [44], the expressions used for the exchange-correlation potential yield electron densities that have a wrong long-range asymptotic decay which affects the observables of these systems [18, 19, 45]. To reduce this error, range-separated hybrid functionals are used where the usual DFT exchange functionals are used at small distances while a Hartree-Fock exact exchange ensures the correct behaviour at larger distances. The interchange between these contributions is modelled via an error function with a characteristic distance that is optimized for each system separately. This procedure has been observed to enhance the accuracy of the predicted properties such as the orbital energies [45–48].

While the schemes described above are well established, their combination is only rarely used. Especially the FEF is found in literature to be approximated only crudely as plane-wave functions [49] or in some expansion as described above [31, 33, 46]. The goal of this thesis is to use the DO formalism with bound states being described by DFT using the above-mentioned optimized range-separated hybrid (OTRSH) functionals to obtain accurate orbital energies and complement it with a FEF that accounts the molecular electrostatic potential explicitly using the finite and infinite element methods which have been applied only to very few quantum mechanical problems [50, 51].

In the protocol used the DFT and time-dependent DFT calculations for excited states are done using a locally modified version of **NWChem** [52] and the **Gaussian 09** package [53]. From those, the DO is calculated with the in-house code **DYSON** developed previously [31]. A self-written interface extracts the required data as *e.g.* the molecular orbital coefficients and overlap matrix of atomic orbitals from the output of these programs. For the computation of the FEF, the implementation that is developed in the framework of this thesis uses the finite element library **Libmesh** [54]. It is an open source library that provides a broad range of capabilities and interfaces to several high-performance linear algebra libraries [55–57]. Moreover, it supports MPI parallelisation and implements a recent formulation of the above-mentioned infinite elements [41]. Especially the latter is to the best of my knowledge a unique option. Furthermore it has an automated procedure to adaptively refine or coarsen the elements according to local error estimates.

The goal of this thesis is to find a systematic way to setup the finite element mesh

for any given molecule to describe the photoelectron with a reasonable accuracy that goes beyond the capabilities of existing programs in this field. This procedure will be tested on different model systems.

- Lithium: light atomic system, many experimental and experimental reference data available, the `ezDyson` calculations are expected to give a good approximation.
- CO<sub>2</sub>: linear easy molecule with some reference data available, used to test some further properties
- Benzene (?): More complex molecule with reference data, well investigated
- Some other system(s) for which `ezDyson` gives poor results (?)

## 2 | Calculation of Photoelectron spectra

The large variety of systems and effects studied with photoelectron spectroscopy led to the existence of diverse methods for theoretical modelling of the spectra [58, 59]. In this work, the focus is put on steady-state photoelectron spectra of molecular systems with a size up to some tens of atoms. As light source, here a classical ultra-violet or soft X-ray source with a discrete spectrum such as a gas discharge lamp or a laser is assumed where the strength of the applied electromagnetic field is weak and the kinetic energy of the outgoing electrons reaches at most tens of electronvolt but may become arbitrarily small.

A summary of methods that can be used to describe the most important scenarios arising in photoelectron spectroscopy is given in the Table 2.1 and some of them are described in more detail in the following sections. The main classification of these methods can be done dividing them into time and frequency domain. These domains are related to each other via the Fourier transform but provide different pictures and properties. Using a frequency-domain method, the Hamiltonian has to be diagonalised to obtain the states' energies and wave functions. In contrast to this, in time domain the Hamiltonian is applied to an initial state to propagate the system in time. Usually such a simulation needs several thousands of propagation steps to obtain reasonable accuracy. Moreover, the joint treatment of bound and continuum states demands a large and flexible basis which makes time domain methods usually computationally very demanding and restricts their applicability to small systems as shown in the Table 2.1. On the other hand, in frequency-domain the nonlinear response properties are neglected so that strong-field effects such as multiphoton ionisation and high harmonic generation (HHG) can not be treated. Due to the different treatment of the system in both domains, they each have a set of methods that can be used and which are more or less specific to the domain as shown in the third column of Table 2.1. The same holds for the quantum chemical methods available to treat the system.

	Field Str.	Method	System Size	Typical Problems	QC <sup>(a)</sup>
Time-domain	weak to strong	TDDO [60] SE Green's function	atoms [61], diatomics [25, 62, 63] triatomics [64]	HHG [26–28], Multiph. ionisation attosec. dynam.[9–14]	(TD)DFT, GASCI [61], EOM-CC [65], CASSCF
Frequency-domain	weak	R-Matrix [66–70] DO [34, 49] Green's Function [71, 72] Koopmans'[16, 73, 74]	up to biomolecules [75], solid state [21, 22]	steady-state, angle-res. PES time-res. PES	RASSCF [31, 46], TD-DFT

Table 2.1: Overview of time- and frequency-domain methods and theyr typical applications.

<sup>(a)</sup>QC: quantum chemical; GASCI: generalised active space configuration interaction; CASSCF: complete active space self-consistent field; EOM-CC: equation of motion- coupled cluster; RASSCF: restricted active space self-consistent field.

Besides the distinction according to the domain, the methods can be also categorised according to the partitioning of the system they use which is sketched in Figure 2.1. The simplest and crudest method is an approach derived from Koopman's theorem where only the ground state of the un-ionised state is considered which is visualised in Figure 2.1 by the fact that the box for this method is only at the  $N$ -electron system. Similarly, also the Green's function approach treats only the initial state but accounts for electron relaxation and thus correlation effects during ionisation due to its quasi-particle picture. In the DO formalism, in contrast, the final state is treated explicitly, using a partitioning into the bound  $N - 1$ -electron state  $|\Psi^{N-1}\rangle$  and the FEF  $|\Psi^{\text{el}}\rangle$ . Finally a large class of methods treat the system uniformly, using a joint treatment for the bound and continuum states as indicated by the single box in Figure 2.1.




Figure 2.1: Scheme of the system-representation used in the different methods.

The Koopmans' approach is, at least among the frequency domain methods, the most prominent representative [18, 73, 74, 76]. In this scheme, the systems ground state is computed with a self-consistent field quantum mechanical method to obtain the one-electron binding energies. The photoelectron spectrum is than estimated using the orbital energies as the transition energies and using uniform intensities. Although this method has shown to be in qualitative agreement with experiments for different systems [18, 19, 76–78], it breaks down in case of strong electron correlation [23, 72] due to relaxation of the orbitals upon ionisation. This method is characterised by its low computational costs and robustness and, thus, well-suited for very large systems such as solid state problems where calculations beyond ground-state DFT are very demanding or not feasible at all.

In the following sections, the most important methods will be briefly introduced according to their affiliation to the groups distinguished in Figure 2.1. First, in section 2.1 the working equations of the Green's function approach are introduced to give a fundamental understanding of this group of methods. Thereafter, in section 2.2 several approaches both in time- and frequency-domain are described that use a unified description of the complete  $N$ -electron system. Finally the DO formalism is derived in more detail, since it is the method of choice in this work, in section 2.3.

## 2.1 The Green's Function Approach

From a formal point of view, the computation of photoelectron spectra using the Green's function approach is similar to the use of Koopmans' approach since in both methods only the ground state of the unionised system is considered. However, the level of theory achievable when using Green's functions is much higher.

In contrast to most other quantum-chemical methods, in the Greens' function approach expectation values for a given operator  $\hat{O}$  are not computed as a scalar product  $\langle \Psi^N | \hat{O} | \Psi^N \rangle$  with the wave function  $|\Psi^N\rangle$  of the  $N$  electron state of interest, but by contour integrals with the Greens' function [79, 80]. Thereby the (one particle) Greens' function is a matrix  $\mathbb{G}$  whose elements are defined as

$$G_{i,j}(t, t') = -i\langle \Psi^N | \hat{T} \left( \hat{a}_i(t) \hat{a}_j^\dagger(t') \right) | \Psi^N \rangle \quad (2.1)$$

with the creation operator  $\hat{a}_j^\dagger(t) = e^{i\hat{H}t} \hat{a}_j e^{-i\hat{H}t}$  of an electron in state  $j$  at time  $t$  in Heisenberg picture and the annihilation operator  $\hat{a}(t) = e^{-i\hat{H}t} \hat{a}_j e^{i\hat{H}t}$  respectively.  $\hat{T}$  is the Dyson time ordering operator that orders the operators  $\hat{a}$  and  $\hat{a}^\dagger$  by their time arguments to ensure that the operator with smaller time argument acts first [79]. Hence, the Greens' function can be interpreted as an additional electron (or hole, depending on the time ordering) propagating from  $t'$  to  $t$  in a system described by the Hamiltonian  $\hat{H}$  [79].

To find an expression of the Green's function (2.1) from which the transition energies and intensities can be extracted, it needs to be reformulated. In the first step, the time ordering operator is written explicitly which results in

$$G_{i,j}(t, t') = i\langle \Psi^N | \hat{a}_j(t) \hat{a}_i^\dagger(t') | \Psi^N \rangle \Theta(t - t') - i\langle \Psi^N | \hat{a}_i^\dagger(t') \hat{a}_j(t) | \Psi^N \rangle \Theta(t' - t). \quad (2.2)$$

Inserting the closure relation  $\hat{1} = \sum_k |\Psi_k^M\rangle \langle \Psi_k^M|$ , where  $M = N \pm 1$  and  $|\Psi_k^M\rangle$  describes a bound state, between the operators in both terms of (2.2), the Lehmann representation [79] is obtained whose Fourier transform is

$$G_{i,j}(\omega) = i \sum_k \frac{\left| \langle \Psi^N | \hat{a}_j | \Psi_k^{N+1} \rangle \right|^2}{\omega - (E_k^{N+1} - E^N) + i\nu} - i \sum_k \frac{\left| \langle \Psi^N | \hat{a}_i^\dagger | \Psi_k^{N-1} \rangle \right|^2}{\omega + (E_k^{N-1} - E^N) - i\nu}, \quad (2.3)$$

where  $\nu$  is a small parameter arising from calculation of principal value and  $\omega$  denotes the argument of the Fourier transform while  $E^N$  and  $E_k^M$  are the energies of the  $N$ -electron ground state  $|\Psi^N\rangle$  and of the  $k$ -th  $M$ -electron state  $|\Psi_k^M\rangle$ , respectively. In this form, the second sum corresponds to transitions in the PES: the nodes of the denominator (poles of the Greens' function) can be easily assigned to the ionisation potentials and, thus, the transition energies in photoelectron spectra. Further, the integrals in the nominator are equivalent to the sudden approximation derived in chapter 2.3.3 and hence provide a good approximation to the transition strengths. The terms in the first sum correspond to the respective quantities of electron detachment [80].

However, computing the Greens' function is a demanding task which is of similar complexity as the computation of a solution to the SE. Over the years several approaches were developed of which the algebraic diagrammatic construction [80] and the equation


of motion [76, 80] are the most prominent. In the diagrammatic construction one starts with an initial zeroth order Greens' function  $\mathbb{G}^0(\omega)$  constructed in a Hartree-Fock basis and corrects it iteratively by the term  $\mathbb{G}^0(\omega)\Sigma(\omega)\mathbb{G}(\omega)$ , where  $\Sigma(\omega)$  is the self-energy, an effective potential that is used to recover electron correlation and relaxation effects [71]. The self-energy usually is expanded in a perturbation series with respect to Feynman diagrams with increasing number of vertices and is exact in the limit of infinite terms [79, 81]. The one-particle energies, Coulomb matrix elements and overlap integrals are obtained from self-consistent field (SCF) calculations [80] *e.g.* on the HF-level [71] but (TD)DFT or any other quantum chemical method can be used as well [18]. Being much easier to compute, the Hartree-Fock basis has the disadvantage that only single configurational electronic states can be treated.

On the other hand, an important advantage of the Greens' function method is that the transition energies are computed directly whereas in most other methods they are calculated as the difference of the initial and final state energies. The latter approach however can lead to errors in the electronvolt range when the difference in the correlation energy is badly estimated [80].

## 2.2 Combined Bound and Continuum State Representation

In contrast to the Greens' function method, in which the explicit description of the FEF omitted, a large group of methods describes the full  $N$ -electron system before and after the ionisation, respectively (see Figure 2.1). In this section a selection of such methods is presented. The treatment of bound and continuum states, however, requires a very flexible formalism whose large computational costs restrict its applicability to small systems that often fulfil certain symmetry-requirements and have a small amount of electrons.

The most prominent representative of this class in frequency domain is the R-matrix method [68, 69, 82]. Its general idea is to conduct a partition of space into regions which are treated differently and connected by explicit boundary conditions to ensure smoothness of the wave function. These regions are constructed with concentric spheres, restricting the symmetry of wave functions in this scheme. Nonetheless it is found to be applied not only to atoms [66, 67, 83] but also to small molecules [70, 84]. In the R-matrix formalism the inner region is chosen large enough to contain the bound part of the  $N$ -electron function that is usually represented by a configuration interaction (CI) expansion of Slater determinants (SDs), using a linear combination of atomic orbitals (LCAO) basis or a grid representation [68]. The FEF is commonly described by a linear combination of bound orbital type functions and continuum functions such as Coulomb waves [69, 82]. In the outer region, an expansion in Coulomb waves is used to ensure the asymptotically correct behaviour. In addition to the inner region and the asymptotic outer region further intermediate regions can be added where the FEF is represented by a multipole-radiation expansion which is a polynomial of



inverse powers of the distance to the centre [68].

A generalisation to non-spherical systems is employed, *e.g.*, by Johnson [85] who used different kinds of non-concentric spheres. One set spheres is centred at atoms while others are placed in the interatomic regions such that the space is filled as dense as possible as shown in Figure 2.2 for a molecule with four atoms. Each atom is located in the centre of a circle denoted as I while the circles II fill the interatomic space. An outer sphere surrounds the molecule to account for the asymptotic region similar to the asymptotic region in R-matrix theory. In this scheme, the exchange-correlation is treated on an approximate level [86] and is spherically averaged, resulting in a description that is equivalent to the muffin-tin potential which is a well-known model in solid state physics [87, 88]. The continuity of the wave-functions as well as their derivatives is ensured over the regions via multiple-scattered-wave theory [85]. The wave-functions are chosen in each region as a one-centre expansion in spherical coordinates of the form

$$\Psi(r, \theta, \phi) = \sum_{l=0}^{l_{\max}} \sum_{m=-l}^l c_{l,m} R_l(r) Y_l^m(\theta, \phi) \quad (2.4)$$

where  $r, \theta, \phi$  are the spherical coordinates and  $Y_l^m(\theta, \phi)$  are the spherical Harmonics [32] and the radial function  $R_l(r)$  is a solution of the radial SE

$$\frac{\partial^2 R(r)}{\partial r^2} + \left( E - V(r) + \frac{l(l+1)}{r^2} \right) R(r) = 0 \quad (2.5)$$

with the respective spherically averaged potential  $V(r)$  [85].

An approach applied by DeCleva *et al.* to  $H_2^+$  [25] and benzene [89] refrains from the use of spheres, allowing for more general boundaries between the inner and outer regions. Here the FEFs are globally represented as the one-centre expansion (2.4) where  $R(r)$  is expressed by a B-spline basis. An advantage of the spline-based description is that smoothness at the interface between the regions is ensured intrinsically.

Another scheme in frequency domain is used by Richards and Larkins [90] with a hybrid ansatz: The bound states of the  $H_2$  molecule are described in the common LCAO scheme and the FEF is described by the product ansatz  $\Psi(\mathbf{r}) = R(r, \theta) e^{im\phi}$ , which is a generalisation of (2.4), where  $R(r, \theta)$  is obtained from a two-dimensional SE on a regular grid using a finite difference (FD) scheme. In contrast to the previously described methods, here no partition of space is performed but instead a finite box with Dirichlet boundary conditions is used. Moreover, the FEF is treated on the HF-level, neglecting correlation effects [90].

Similar descriptions are used in time domain by several authors [61, 65, 91]. Even though they are much more demanding than calculations in frequency domain, time do-

main methods can predict strong-field effects and are used to simulate attosecond dynamic which is both not achievable using frequency-domain methods.

An important difference between time- and frequency-domain methods of practical use is that in time-domain unbound particles are described by a wave packet and hence as a localised function. Moreover, at short times after ionisation the continuum states often can be assumed to be of similar spacial extend as bound states, allowing the basis to be a linear combination of bound state functions but having a complex energy whose real part introduces oscillations and the imaginary part damps the wave function, leading to a finite lifetime of the function [65]. Moreover a complex absorbing potential (CAP) (discussed in chapter 4.4.1 in more detail) is most often applied, ensuring that the continuum states can be described by bound-state functions by cutting off that part of the wave function which is not localised at the molecule anymore. Due to these considerations the LCAO basis can be used for the description of the free particles as well, see *e.g.* Jagau *et al.* [65]. In other simulations, grid-based descriptions are chosen using symmetry-adapted coordinate systems [64, 91–93], allowing for a product ansatz similar to (2.4) and hence a reduction in dimensionality. On the remaining one-dimensional grids often a discrete variable representation (DVR) (described in section 3.3 of this thesis) is chosen. As an example, Yip *et al.* [94] simulate double ionisation of atomic beryllium in spherical coordinates, using the expansion (2.4) where  $R(r)$  is separated into two regions in which a DVR and a finite element DVR (FE-DVR) scheme are used respectively. As other examples, Tao *et al.* [91] as well as Bauch *et al.* [61, 62] use the FE-DVR basis in spheroidal coordinates.

The examples introduced above represent only a small fraction of the methods used to describe photoionisation in frequency and time domain. But a drawback that is common to these methods is that a large number of continuum functions is required which are correlated with the bound electrons and thus lead to a computationally expensive description of the system. To overcome this drawback, in the DO formalism that is presented in the coming section the description of bound and continuum states is separated into separate problems.

## 2.3 The Dyson Orbital Formalism

The Dyson orbital (DO) formalism can be considered as an approximation to the methods described above. Here the free and bound states are described separately, see Figure 2.1, using a product ansatz. Using such a separation leads to a neglect of correlation effects between the outgoing electron and the bound states and is often denoted as sudden ionisation limit [31, 33]. Moreover, non-linear effects as well as recombination transitions are not considered within DO theory. An important advantage, however, is that the overlap between the initial state and the bound part of the final state is formulated as a one-electron quantity, called DO which can be used to simplify the description of the photoionisation process. Usually the DO scheme is considered in the frequency domain, but a time domain formulation exists as well and is described in the following section [60].

### 2.3.1 Time-dependent Dyson Orbitals

A good starting point for the DO formalism is an expansion of the time-dependent  $N$ -electron function  $|\Psi^N(t)\rangle$  (omitting the spacial coordinates for brevity) in the form

$$|\Psi^N(t)\rangle = \frac{1}{\sqrt{N}} \sum_k |\Psi_k^{DO}(t)\rangle |\Psi_k^{N-1}\rangle e^{iE_k^{N-1}t} \quad (2.6)$$

where  $E_k^{N-1}$  are the energies of the  $N - 1$ -electron bound states described by the time-independent wave functions  $|\Psi_k^{N-1}\rangle$  that are complete in the space of  $N - 1$ -electron wavefunctions. The expansion coefficients  $|\Psi_k^{DO}(t)\rangle$  have the dimensionality of a one-particle function and are denoted as time-dependent DO (TDDO). An important feature of the expansion (2.6) is that the dynamics of the  $N$ -electron system is reduced to a system of one-electron quantities  $|\Psi_k^{DO}(t)\rangle$ , propagated according to the electronic SE [60]. In this approach the interaction of the DO with the bound states is approximated by the mean-field electrostatic potential [60], neglecting exchange and correlation.

The physical interpretation of the TDDO becomes clear when regarding its definition, given by the  $N - 1$ -electron integral

$$|\Psi_k^{DO}(t)\rangle = e^{iE_k^{N-1}t} \sqrt{N} \langle \Psi_k^{N-1} | \Psi^N(t) \rangle_{N-1}. \quad (2.7)$$

The remaining coordinate (2.7) belongs to the photoelectron since  $|\Psi^N(t)\rangle$  is restricted to the description of bound states. Considering eq. 2.7 in a frozen orbital approximation, *i.e.* one-electron ionisation, the TDDO corresponds to the photoelectron. However, relaxation effects lead to non-orthogonal orbitals of the  $N$ - and  $N - 1$ -electron systems and thus lead to additional contributions to the TDDO. Therefore, when considering relaxation effects, the TDDO is a quasi-particle describing the electron that is ionised, including relaxation and correlation effects [33, 60].

### 2.3.2 Time-independent Dyson Orbitals

Working in frequency domain, no propagation needs to be considered. Instead, the quantity of interest here is the stationary DO which corresponds to the TDDO at  $t = 0$ . From eq. 2.7 the DO can be interpreted as quasi-particle that is ejected by the irradiating light [33].

While for the TDDO only the general expressions were shown, in this section, since the time-independent DO formalism is used within this work, a more detailed derivation of the expressions is done.

The main advantage of this formalism in frequency-domain becomes clear when the photoelectron cross-section is considered. In the Fermis' Golden Rule [29] formulation, which assumes the wavelength to be much larger than the characteristic size of the system under study and weak irradiating light-field, the cross-section for the transition between an initial  $i$  and final  $k$  states is in atomic units [31, 90]

$$\sigma(\epsilon) = \frac{2}{3} \sum_k (\epsilon + E_k - E_i) \left| \langle \Psi_i^N | \hat{\mathbf{d}} | \Psi_k^N \rangle \rho(\epsilon) \right|^2 \propto \sum_k |\mathbf{D}_k|^2 \quad (2.8)$$

where  $\epsilon = h\nu - (E_k - E_i)$  is the kinetic energy of the photoelectron that is determined by the photon energy  $h\nu$  and the energies  $E_\alpha$  of the initial unionised state  $|\Psi_i^N\rangle$  and the final ionised state  $|\Psi_k^N\rangle$  which includes all electrons, further  $\mathbf{D}_k = \langle \Psi_i^N | \hat{\mathbf{d}} | \Psi_k^N \rangle$  is the transition dipole moment and  $\rho(\epsilon)$  is the density of final states which is in the following assumed to be a delta function. Writing the initial and final states each as SDs

$$|\Psi_i^N\rangle = \hat{A}_N |\Phi_{i,1} \dots \Phi_{i,N}\rangle \quad (2.9a)$$

$$|\Psi_k^N\rangle = \hat{A}_N |\Phi_{k,1} \dots \Phi_{k,N-1} \Psi_k^{\text{el}}\rangle \quad (2.9b)$$

where  $\hat{A}_N$  is an  $N$ -electron antisymmetrisation operator,  $|\Phi_{k,j}\rangle$  are the  $j$ -th (Kohn-Sham) orbitals and  $\Psi_k^{\text{el}}$  is the FEF. The index  $k$  enumerates the final states which can have an arbitrary electron configuration here. The dipole operator  $\hat{\mathbf{d}}$  is a one-electron operator that can be written as  $\hat{\mathbf{d}} = \sum_{j=1}^N \hat{\mathbf{d}}_j$  where  $\hat{\mathbf{d}}_j = \mathbf{r}_j$  in length gauge or  $\hat{\mathbf{d}}_j = \nabla_j / (\epsilon + E_k - E_i)$  in velocity gauge respectively [90].

Using the SD representations (2.9), the integral  $\mathbf{D}_k$  in eq. (2.8) can be written as

$$\mathbf{D}_k = \langle \Phi_{i,1} \dots \Phi_{i,N} | \hat{A}_N \sum_{j=1}^N \hat{\mathbf{d}}_j \hat{A}_N | \Phi_{k,1} \dots \Phi_{k,N-1} \Psi_{\text{el}} \rangle, \quad (2.10)$$

where hermiticity of the antisymmetrisation operator is used. The expression (2.10) can be further expanded taking into account that  $\hat{A}_N$  commutes with the dipole operator and making use of the relation  $\hat{A}_N \hat{A}_N = \sqrt{N!} \hat{A}_N = \sum_P (-1)^p \hat{P}$  where the sum goes over all permutations  $\hat{P}$  of electron indices with parity  $p$

$$\mathbf{D}_k = \sqrt{N!} \sum_P (-1)^p \sum_{j=1}^N \langle \Phi_{i,P(1)} \dots \Phi_{i,P(N)} | \hat{\mathbf{d}}_j | \Phi_{k,1} \dots \Phi_{k,N-1} \Psi_{\text{el}} \rangle \quad (2.11)$$

$$= \sqrt{N!} \sum_P (-1)^p \sum_{j=1}^N \langle \Phi_{i,P(j)} | \hat{\mathbf{d}}_j | \Phi_{k,j} \rangle \langle \Phi_{i,P(1)} | \Phi_{k,1} \rangle \dots \langle \Phi_{i,P(j-1)} | \Phi_{k,j-1} \rangle \quad (2.12)$$

$$\times \langle \Phi_{i,P(j+1)} | \Phi_{k,j+1} \rangle \langle \Phi_{i,P(N-1)} | \Phi_{k,N-1} \rangle \dots \langle \Phi_{i,P(N)} | \Psi_{\text{el}} \rangle \quad (2.13)$$

where  $P(j)$  is a permutation of the  $j$ -th orbital. Thereby the term  $j = N$  differs qualitatively from the others since the dipole operator acts on the FEF. Hence the sum can be reordered to obtain

$$\begin{aligned} \mathbf{D}_k &= \underbrace{\sqrt{N!} \sum_P (-1)^p \langle \Phi_{i,P(1)} | \Phi_{k,1} \rangle \dots \langle \Phi_{i,P(N-1)} | \Phi_{k,N-1} \rangle \langle \Phi_{i,P(N)} | \hat{\mathbf{d}}_j | \Psi_{\text{el}} \rangle}_{\langle \Psi_k^{\text{DO}} |} + \\ &\quad \underbrace{\sqrt{N!} \sum_P (-1)^p \sum_{j=1}^{N-1} \langle \Phi_{i,P(1)} | \Phi_{k,1} \rangle \dots \langle \Phi_{i,P(j-1)} | \Phi_{k,j-1} \rangle \langle \Phi_{i,P(j)} | \hat{\mathbf{d}}_j | \Phi_{k,j} \rangle \times} \\ &\quad \langle \Phi_{i,P(j+1)} | \Phi_{k,j+1} \rangle \dots \langle \Phi_{i,P(N-1)} | \Phi_{k,N-1} \rangle \langle \Phi_{i,P(N)} | \Psi_k^{\text{el}} \rangle \end{aligned} \quad (2.14)$$

where the first sum is denoted as DO and the second as conjugate DO respectively [63]. To reduce the large amount of integrals therein, the strong orthogonality approximation

is applied under the assumption that the overlap

$$\langle \Phi_{i,j} | \Psi_{\text{el}} \rangle = 0 \quad \forall j = 1, \dots, N \quad (2.15)$$

of the FEF with all bound states vanishes. This is strictly valid if  $|\Phi_{i,j}\rangle$  and  $|\Psi^{\text{el}}\rangle$  would correspond to the same Hamiltonian. But in most cases the relaxation of the orbital  $|\Phi_{i,j}\rangle$  is small, leading to a small but non-zero overlap with the FEF [46, 63]. Applying the strong orthogonality condition leads to the drop out of the second sum in eq. 2.14 and the transition dipole moment simplifies to

$$\begin{aligned} \mathbf{D}_k &= \sum_p (-1)^P \langle \Phi_{i,P(1)} | \Phi_{k,1} \rangle \dots \langle \Phi_{i,P(N-1)} | \Phi_{k,N-1} \rangle \langle \Phi_{i,P(N)} | \hat{\mathbf{d}}_j | \Psi_k^{\text{el}} \rangle \\ &= \langle \Psi_k^{\text{DO}} | \hat{\mathbf{d}}_j | \Psi_k^{\text{el}} \rangle \end{aligned} \quad (2.16)$$

which corresponds to the definition of the TDDO in eq. (2.7) for  $t = 0$  as mentioned above. The expression for the PES cross-section in the DO formalism simplifies to

$$\sigma(\epsilon) = \frac{2}{3} \sum_k (\epsilon + E_k - E_i) \left| \langle \Psi_k^{\text{DO}} | \hat{\mathbf{d}} | \Psi_{\text{el}} \rangle \right|^2. \quad (2.17)$$

In the derivation given above, it is assumed that initial and final states can be represented by single SD. In practice, states often are described by a linear combination of SDs with different electronic configurations. A respective generalisation of the DO is straight forward but introduces additional summations and thus leads to more complex terms [46].

### 2.3.3 Sudden Approximation

Further simplification of the DO formalism can be obtained by applying the so-called sudden approximation (SA) where the computation of the FEF is omitted. In the SA it is assumed that the transition to the continuum with corresponding energy has a constant probability [95]. This can be justified for high kinetic energies where the oscillations of the FEF are much stiffer than the structures of bound states so that small changes in the energy of the FEF do not play an important role. Thus, the transition dipole moment (2.14) reduces to the scalar

$$D_k = \sum_p (-1)^P \langle \Phi_{i,P(1)} | \Phi_{k,1} \rangle \dots \langle \Phi_{i,P(N-1)} | \Phi_{k,N-1} \rangle \quad (2.18)$$

which involves only the bound  $N - 1$ -electron system [96]. With this assumption the kinetic energy of the free electron dependence of the PES is neglected, assuming error introduced is a similar factor for all transitions. This has shown to be a valid assumption if the nature and spacial extend of the Dyson orbitals is similar for different transitions. Note that the expression (2.18) corresponds to the nominator of the Greens' function in (2.3) and thus is the level of theory at which the Greens' function PES transitions are computed.

Finally, the expression (2.18) can be further simplified to get the approach described in the introduction to this chapter, based on Koopmans' theorem. Therefore the electron

relaxation is neglected and thus the  $N - 1$ -electron state can be written as  $|\Psi_k^{N-1}\rangle = \hat{a}_k |\Psi_i^N\rangle$ . Thus, the sum over all permutations  $\hat{P}$  in (2.18) reduces to the case  $\delta_{P(j),j}$  due to orthogonality of the orbitals. Moreover, due to the normalisation of the orbital functions, all transitions have a probability of one and thus corresponds to the scheme described earlier.

# 3 | Description of Free Particles

The description of unbound particles has several complications of conceptual and technical kind. In this chapter, different numerical methods are introduced that can be used to solve the one-electron SE. In section 3.1, the conceptual differences between bound and continuum states as well as the numerical treatment of the latter in a finite basis are discussed. Thereafter, in the sections 3.2 - 3.7, different numerical methods are introduced that can be used to solve the one-particle SE on a molecular domain.

## 3.1 Continuum Waves

Continuum waves are discussed only sparsely in lectures on quantum mechanics, even though they posses certain fundamental differences compared to bound states. The states considered here have a spacially infinite extend as it is well-known for plane waves

$$\Psi^{\text{plan}}(\mathbf{r}) = \sqrt{\frac{|\mathbf{k}|}{(2\pi)^3}} e^{i\mathbf{k}\mathbf{r}} \quad (3.1)$$

with wave-vector  $\mathbf{k}$  as well as for spherical waves. Moreover, these waves are not square integrable and hence can not be normalised according to  $\int \Psi^\dagger(\mathbf{r})\Psi(\mathbf{r})d\mathbf{r} = 1$  and thus the probability interpretation is invalid [97]. Instead, these functions are sharp in a continuous variable (namely the momentum) and hence should rather be interpreted as probability densities, suggesting the normalisation  $\int \Psi^\dagger(\mathbf{r})\Psi(\mathbf{r}')d\mathbf{r} = \delta(\mathbf{k} - \mathbf{k}')$  where  $\delta(\mathbf{k})$  is the Dirac delta distribution [97]. This property distinguishes the analytical FEF from those obtained with numerical methods that often have a finite extent and are obtained from an approximate Hamiltonian matrix which has no continuous spectrum due to finiteness of its basis. Especially the difference in the normalisation of wave functions and by this in the dimensionality of overlap integrals also affects the integral evaluated when calculating the transition dipole matrix elements [98]. However, the Stieltjes imaging approach described in section 3.1.1 gives a formal verification for the use of square-integrable function for the representation of a continuum state. Moreover, it provides a formalism to enhance the quality of spectra obtained with numerical methods.

A further aspect of numerical treatment of continuum functions that is only sparsely discussed in literature concerns the question of the physical interpretation of the numerical solution. When dealing with approximate continuum functions, it is often assumed that the analytic function of interest  $|\Psi_a(\mathbf{k})\rangle$  is approximated by that particular numeric solution  $|\Psi_n(\mathbf{k}')\rangle$  which is closest in energy to the desired one [25]. An alternative interpretation

would be to assume that the numerical solution corresponds to an approximation to  $|\Psi_n\rangle \approx \frac{1}{k^+ - k^-} \int_{k^-}^{k^+} |\Psi_a(k)\rangle dk$  which would change the interpretation and normalisation.

In other works simulating PES in frequency domain, mainly two different representations of the photoelectron are found which are based on plane-wave expansions in spherically symmetric bases [33, 34, 99]. One of the expansions is that in spherical waves [32]

$$\Psi_k^{\text{sph}}(\mathbf{r}) = 4\pi \sum_{l=0}^{\infty} \sum_{m=-l}^l i^l j_l(kr), Y_l^m(\theta, \phi) Y_l^{\dagger m}(\theta_k, \phi_k) \quad (3.2)$$

where  $k = |\mathbf{k}|$  is the length of the wave vector  $\mathbf{k}$ ,  $j_l(kr)$  are spherical Bessel functions that solve the radial SE (eq. (2.5) with  $V(r) = 0$ ) and the arguments to the spherical Harmonics  $Y_l^m$ ,  $(\theta, \phi)$  and  $(\theta_k, \phi_k)$ , are the angles of the spherical coordinate system in real-space and Fourier-space, respectively [33]. Another expansion of a plane wave often used that in Coulomb waves

$$\Psi_k^{\text{Coul}}(\mathbf{r}) = \frac{1}{(2\pi)^{\frac{3}{2}}} \frac{1}{r} \sum_{l=0}^{\infty} \sum_{m=-l}^l i^l (2kr)^l e^{-\pi \frac{Z}{2k}} \frac{|\Gamma(l+1+i\frac{Z}{k})|}{\Gamma(2l+2)} e^{ikr} F_1(l+1-i\frac{Z}{k}, 2l+2, 2ikr) Y_l^m(\theta, \phi) Y_l^{\dagger m}(\theta_k, \phi_k) \quad (3.3)$$

where  $Z$  is the charge of the nucleus and  $F_1$  is the confluent hypergeometric function of the first kind [99, 100]<sup>1</sup>. Assuming  $Z = -1$ , *i.e.* the case of the hydrogen atom, the limit  $k \rightarrow 0$  the confluent hypergeometric function is

$$\lim_{k \rightarrow 0} F_1(l+1+\frac{i}{k}, 2l+2, 2ikr) = (2l+1)!(2r)^{-l-\frac{1}{2}} J_{2l+1}(\sqrt{8r}) \quad (3.4)$$

where  $J_{2l+2}(x)$  is the Bessel function that is connected with the spherical Bessel functions  $j_l(x)$  via the relation  $j_l(x) = \sqrt{\frac{\pi}{2x}} J_{l+\frac{1}{2}}(x)$  [32]. This case is separately available in **ezDyson** [33].

The latter approach accounts for the electrostatic potential of the remaining anion at larger distances to the molecule but does not account for the Coulomb singularities due to the particular molecular structure respectively. To correct for this error which becomes significant especially for lower kinetic energies where the influence of the potential is stronger, *e.g.* an effective charge  $Z^* < 1$  is used [99] or the ansatz function is orthonormalised with respect to the DO [101–103] which can be shown to be equivalent to a respective shift of the coordinate system such that the expectation value of the dipole moment of the DO vanishes [101].

In this work the spherical waves (3.2) are used as reference basis for the computed solutions to

- elaborate on Bessel- and hypergeometric functions
- show figures of radial functions
- Write out  $Y_l^m$ , describe orthonormality, dipole selection rules

Another conceptual question is about the infinite degeneracy of the analytic solution which becomes clear when considering an example of the ionisation of a hydrogen-like atom whose analytical FEF is well-known and has the form [32]

---

<sup>1</sup> $F_1(\alpha, \gamma, z) = 1 + \frac{\alpha}{\gamma} z + \frac{\alpha(\alpha+1)}{\gamma(\gamma+1)} \frac{z^2}{2!} + \dots$




Figure 3.1: Radial part of spherical waves and Coulompart of spherical waves and Coulomb waves (confluent hypergeometric function approximated with 35 terms each) for  $l = 0$  (left),  $l = 1$  (centre) and  $l = 2$  (right).

The function (3.3) is, given its energy  $E = \frac{1}{2}|\mathbf{k}|^2$ , infinitely degenerate in the direction of  $\mathbf{k}$  as well as in the quantum numbers  $l$  and  $m$ . Hence the wave function of a photoelectron can consist of any superposition of the form (3.3), choosing the coefficients  $c_{l,m}$  accordingly. The coefficients  $c_{l,m}$  of the physically observable FEF are determined by the matrix elements of the dipole operator with the DO (2.16). Considering as an example the ionisation of hydrogen in its ground state, the FEF is in one of the three  $p$ -waves because the probability to access any of the other vacuum-states with the correct energy vanishes due to dipole selection rules for spherically symmetric systems.

Unfortunately, as will be shown below in section 6.1, the considerations made above are not true for the numerical solution anymore where a finite set of superpositions is obtained which, due to numerical treatment, differ in energy and thus can not be assigned to a common value of  $\mathbf{k}$ . Choosing the energetically best-fitting solution though might in principle result in a function with vanishing contribution of the  $p$ -type orbitals while, for slightly changed computational scheme or target energy could yield a solution with large  $p$ -contributions with.

### 3.1.1 Stieltjes Imaging

The Stieltjes imaging provides an elegant way to still use the numerically obtained solutions of a non-continuous spectrum by using moment theory. Starting from the expression for the photoelectron spectrum

$$\sigma(\varepsilon) \propto \left| \langle \Psi_0 | \hat{d} | \Psi_{\text{el}} \rangle \right|^2 \quad (3.5)$$

where  $\Psi_{\text{el}}$  is a continuum function and as such, as discussed above, not square integrable which is an important characteristic for the dimensionality of this expression [98]. To repair this, the Stieltjes imaging approach uses spectral moments, where the  $n$ -th spectral moment is of the form

$$M_n = \langle \Psi_0 | \hat{d} (\hat{H} - E_0)^n \hat{d} | \Psi_0 \rangle \quad (3.6)$$

where  $\hat{H}$  is the systems Hamiltonian and  $E_0$  is the ground state energy [104]. Inserting the closure relation  $\sum_{\alpha} |\Psi_{\alpha}\rangle\langle\Psi_{\alpha}| + \int |\Psi_{\varepsilon}\rangle\langle\Psi_{\varepsilon}|d\varepsilon$  for the full (anatlytic) Hamiltonian results in

$$M_n = \sum_{\alpha} \langle\Psi_0|\hat{d}(\hat{H} - E_0)^n|\Psi_{\alpha}\rangle\langle\Psi_{\alpha}|\hat{d}|\Psi_0\rangle + \int \langle\Psi_0|\hat{d}(\hat{H} - E_0)^n|\Psi_{\varepsilon}\rangle\langle\Psi_{\varepsilon}|\hat{d}|\Psi_0\rangle d\varepsilon \quad (3.7)$$

$$= \sum_{\alpha} (E_{\alpha} - E_0)^n \left| \langle\Psi_0|\hat{d}|\Psi_{\alpha}\rangle \right|^2 + \int_0^{\infty} \varepsilon^n \left| \langle\Psi_0|\hat{d}|\Psi_{\varepsilon}\rangle \right|^2 d\varepsilon. \quad (3.8)$$

However, inserting the unity expression for the numerical Hamiltonian which does not have continuum functions but an everywhere discrete spectrum and hence  $L^2$ -integrable eigenfunctions results in

$$M_n = \sum_{j=0}^N (E_j - E_0) \left| \langle\Psi_0|\hat{d}|\Psi_j\rangle \right|^2 \quad (3.9)$$

where  $N$  denotes the number of eigenfunctions  $|\Psi_j\rangle$  [98, 105]. For a good numerical approximation, the expressions (3.9) and (3.7) should be similar for as many  $n$  as possible where, due to convergence of the spectral moments,  $n < 2$  [98]. These considerations are used in the Stieltjes imaging approach with moment theory [106, 107].

In the protocol derived from this scheme thus the oscillator strength  $|\langle\Psi_0|\hat{d}|\Psi_{\text{el}}\rangle|^2$  is approximated by the histogram

$$\sigma(\varepsilon) = \frac{2}{3} \frac{df}{d\varepsilon} = \frac{2}{3} \begin{cases} 0 & 0 < \varepsilon < \varepsilon_1(n) \\ \frac{1}{2} \frac{(f_i + f_{i+1})}{\varepsilon_{i+1} - \varepsilon_i} & \varepsilon_j(n) < \varepsilon < \varepsilon_{j+1}(n) \\ 0 & \varepsilon_n(n) < \varepsilon \end{cases} \quad (3.10)$$

where  $(\varepsilon_i, f_i)_{i=1,\dots,l}$  are the principal pseudo spectra [105] that are obtained from

$$\mathbf{M} = \mathbb{E}\mathbf{f} \quad (3.11)$$

where  $\mathbf{f}_i = f_i$ ,  $\mathbf{M}_i = M_i$  and  $\mathbb{E}_{i,j} = \varepsilon_i^j$  [104]. It can be shown moreover that  $M_n$  is only for  $n < 2$  finite and therefore negative moments are used [98]. Moreover, it can be shown that the expression (3.9) converges already for small  $N$ , so only a small fraction of the spectrum needs to be considered in practice and the matrix equation (3.11) remains at a reasonable size. Usually not more than 10 – 12 moments are used to avoid spurious oscillations due to unconverged high order moments [108].

In practice this method becomes very demanding if energetically narrow transitions occur since many terms in expression (3.9) are needed for convergence, making this method effective mainly for low energy range [25]. Another disadvantage that no asymptotic information about this scheme present [25].

## 3.2 Finite Differences and Finite Volumes

The most common and straightforward approach to solve differential equations numerically is via the so-called finite difference (FD) scheme. Starting from the one-particle SE in

atomic units

$$\left( \frac{1}{2} \nabla^2 + V(\mathbf{r}) \right) \Psi(\mathbf{r}) = E \Psi(\mathbf{r}) \quad (3.12)$$

where  $V(\mathbf{r})$  is an arbitrary potential at this point. Considering a general problem where the low symmetry does not support a product ansatz which would reduce the dimensionality and thus the complexity of the problem, the kinetic energy operator has at least 6 non-diagonal terms which lead to an Hamiltonian of a non-banded structure [109]. Moreover, the finite-difference schemes require the evaluation points to be on a regular grid whose size is governed by the sharpest features in the system while local refinement is hardly available. To resemble the expected fine structure close to the nuclei, a dense grid is needed but any FD scheme beyond the first order requires the evaluation points to be on a regular grid, making the computation of a FEF with a reasonable box size very expensive [90]. Nonetheless, there are applications to this scheme to the SE [109, 110], some of them with massive parallelisation using MPI and multiple GPUs [111].

To overcome this bottleneck for systems with non-uniform parameters, often the finite volume method is used. The finite volume method is an integral method that is not based on a regular grid. Instead, for the estimation of the kinetic energy Gauß's theorem  $\int_V \nabla \mathbf{u}(\mathbf{r}) dV = \int_{\partial V} \mathbf{u}(\mathbf{r}) \mathbf{n} dS$  is used where  $\mathbf{u}(\mathbf{r})$  is a vector-valued function and  $\mathbf{n}$  is the normal vector on the surface  $S$  of the volume  $V$  under consideration. Chosing  $\mathbf{u}(\mathbf{r}) = \nabla \Psi(\mathbf{r})$  yields the relation

$$\nabla^2 \Psi(\mathbf{r}) = \lim_{V \rightarrow 0} \frac{\int_{\partial V} \nabla \Psi(\mathbf{r}) \mathbf{n} dS}{V}. \quad (3.13)$$

This scheme becomes especially interesting when the finite volume elements are chosen to be Voronoi cells [112]. A Voronoi-tesselation is associated with a grid and is constructed such that the Voronoi cell  $T_i$  consists of all points in space that are closer to the point  $i$  than to any other points in the grid [113–115].

This description has the advantage that the quantities on the right-hand side of (3.13) can be associated with properties of the Voronoi-Cell, leading to the first-order approximation




Figure 3.2: 2-dimensional Voronoi diagram on an arbitrary point-distribution. [116]

$$\nabla^2 \Phi(\mathbf{r}_i) = \frac{1}{v_i} \sum_j^{\text{neighbours}} \frac{\Psi(\mathbf{r}_j) - \Psi(\mathbf{r}_i)}{h_{ij}} s_{ij} a \quad (3.14)$$

where  $v_i$  is the volume of the  $i$ -th Voronoi-cell,  $h_{ij} = |\mathbf{r}_i - \mathbf{r}_j|$  is the distance between the centres of the  $i$ -th and  $j$ -th Voronoi-cell and  $s_{ij}$  is the length of the common facet of the  $i$ -th and  $j$ -th Voronoi cell.

Such a scheme is applied *e.g.* by Son and Chu [112, 116] to the time-dependent SE, studying multi-photon ionisation of several molecular systems.

### 3.3 Pseudospectral Methods

Under the term (pseudo)spectral methods a large group of methods is comprised which treat the differential equation of interest variationally using an orthogonal basis  $\text{span}\{\varphi_i(\mathbf{r})\}$ , hence seeking for a solution of the form

$$\Psi(\mathbf{r}) = \sum_i^n c_i \varphi_i(\mathbf{r}) \quad (3.15)$$

where  $c_i$  are coefficients to be found [117]. In the pseudospectral methods the basis functions  $\varphi_i(\mathbf{r})$  are smooth global functions such as  $\varphi_{\mathbf{k}}(\mathbf{r}) = \frac{1}{(2\pi)^{\frac{3}{2}}} e^{i\mathbf{k}\mathbf{r}}$  resulting in the Fourier method [118]. Alternatively Jacobi, Chebychev or Legendre polynomials are commonly used [119].

To find the corresponding coefficients, the equation is evaluated on a grid, leading to a linear system of equations to be solved. Depending on the technique used for determining the coefficients, it can be seen as a high-order FD or high-order ( $p$ -) finite element method (FEM). One large advantage of this class of methods is that the error of the solution  $\psi(\mathbf{r})$  usually decays exponentially with the number  $n$  of basis functions and the grid can be chosen coarsely, making the numerical scheme very efficient [119, 120]. In addition, many formulations allow for an implementation making use of the fast fourier transform [119].

A special representative subgroup of the pseudospectral methods are the so-called discrete variable representation (DVR) schemes which are frequently used to study quantum chemical problems [94, 121, 122]. Since this scheme is applicable in one dimension only, often symmetry-adapted coordinates such as spherical coordinates are used and the ansatz

$$\varphi = \sum_{l,m} R_{l,m}(r) Y_l^m(\theta, \phi) \quad (3.16)$$

is applied where  $Y_l^m(\theta, \phi)$  are the well-known spherical harmonics [123] and  $R_{l,m}(r)$  are radial functions that are computed explicitly, solving the radial SE.

The basis functions used in this scheme commonly are Lagrange polynomials [91, 121, 122] whose nodes are chosen by a general Gaußquadrature rule [121] or from a quadrature rule for the radial Coulomb function which is in particular popular for quantum chemical problems [120, 122]. Even if the DVR provides a flexible and fast-convergent basis, an extension to higher dimensionality can be achieved by a product ansatz only and thus is applicable to problems with high symmetry only.

Moreover, as for the other pseudospectral methods, the system matrices are dense and thus the solution is numerically expensive. This is, at least in parts, solved by using a hybrid scheme of the finite element and DVR scheme where the DVR scheme is used on small segment only, connected by single bridge-functions each [91, 121].

## 3.4 Radial Basis Functions

The radial basis function (RBF) technique is a basis that is based on an arbitrary point distribution with very general properties. The basis functions  $\varphi_i(\mathbf{r})$  used in this technique need to be spherically symmetric, *i.e.*  $\varphi_i(\mathbf{r}) = \varphi_i(|\mathbf{r}|)$  and are placed at the grid points of a given domain. Among the commonly used functions are, besides linear and cubic functions, multiquadratic ( $\sqrt{r^2 + r_0^2}$ ), inverse multiquadratic ( $\frac{1}{\sqrt{r^2 + r_0^2}}$ ) and Gaussian ( $e^{-\frac{r^2}{2r_0^2}}$ ) functions [124, 125].

If the parameter  $r_0$  is chosen reasonably the scheme is numerically stable and fast-converging [124]. Another advantage is the straight-forward implementation and generality of this method. But the choice of  $r_0$  can be non-trivial especially for highly non-regular grids where the occurring large distances require broad and smooth functions which lead to instabilities for close-lying points. Moreover the resulting system matrices are dense due to the global definition of the ansatz functions, making it computationally expensive.

The RBF-scheme is used not only for solving differential equations but is also a useful tool for interpolation of scattered data in arbitrary dimensions [126] and surface reconstruction from scattered data [127].

An important disadvantage of the RBF scheme is that the space between two interpolation points is interpolated non-linearly and, depending on the distance of the two functions, the values are over- or underestimated. This error is visualised in Figure 3.3 for the case of a function that is one at both supporting points, represented by a Gaussian RBF basis. The intermediate course of the function depends strongly on the ratio of the broadening  $r_0$  and the difference between the supporting points.

While this behaviour is not problematic for visual purposes, it may introduce a sub-structure which can lead to unintended behaviour of the solution. Since on irregular grids strongly varying distances between supporting points occur, this behaviour is stronger the less regular the initial data are.




Figure 3.3: A function (blue) represented in a RBF basis with Gaussian functions, centred at different distances compared to the broadening parameter  $r_0$ .

## 3.5 Finite Elements

In the FEM the differential equations to be solved are formulated in their weak form that can be understood as a generalisation of it. Starting with the well-known (strong) form

of the SE defined on a domain  $\Gamma$  in atomic units which are used here and in the following chapters

$$-\frac{1}{2}\nabla^2\Psi(\mathbf{r}) + V(\mathbf{r})\Psi(\mathbf{r}) = E\Psi(\mathbf{r}) \quad \mathbf{r} \in \Gamma \quad (3.17)$$

with the condition for the boundary domain  $\partial\Gamma$  which is assumed to be Lipschitz continuous

$$\Psi(\mathbf{r}) = 0 \quad \mathbf{r} \in \partial\Gamma. \quad (3.18)$$

The first term in equation (3.17) corresponds to the kinetic energy and  $V(\mathbf{r})$  is a potential that is not further specified at this point. The unknowns of this equation are the wave function  $\Psi(\mathbf{r})$  and respective energy  $E$ . To bring equation (3.17) into the weak form, it is first multiplied by the complex conjugate of a test function  $\Phi(\mathbf{r})$  which fulfils the same boundary conditions and needs to be differentiable. In the second step, one integrates over the whole space of interest resulting in

$$\int d\mathbf{r} \left( -\frac{1}{2} (\nabla^2\Phi^*(\mathbf{r})) \Psi(\mathbf{r}) + V(r)\Phi^*(\mathbf{r})\Psi(\mathbf{r}) \right) = E \int d\mathbf{r} \Phi^*(\mathbf{r})\Psi(\mathbf{r}). \quad (3.19)$$

The kinetic energy term can be symmetrised using Green's first identity  $\int_{\Gamma}(\nabla^2\Psi)\Phi^*d\mathbf{r} = \int_{\partial\Gamma}(\nabla\Psi)\Phi^*ds - \int_{\Gamma}(\nabla\Psi)(\nabla\Phi^*)$  to obtain the final expression

$$\int d\mathbf{r} \left( \frac{1}{2} (\nabla\Phi^*(\mathbf{r})) (\nabla\Psi(\mathbf{r})) + V(r)\Phi^*(\mathbf{r})\Psi(\mathbf{r}) \right) = E \int d\mathbf{r} \Phi^*(\mathbf{r})\Psi(\mathbf{r}) \quad (3.20)$$

$$\Phi(\mathbf{r}) = 0 \quad \Psi(\mathbf{r}) = 0 \quad \mathbf{r} \in \Gamma \quad (3.21)$$

which is denoted as the weak form of the SE. A function  $\Psi(\mathbf{r})$  is considered as a solution of (3.20) if the equation holds for any test function  $\Phi(\mathbf{r})$ . Since usually only real-valued bases functions are used, in the literature the complex conjugation is omitted. From a mathematical point of view, since  $\Phi(\mathbf{r})$  is an arbitrary function, complex conjugation does not change anything here but it will be shown later that it is more reasonable from a physical point of view if the weak formulation is defined as given in equation (3.20).

The weak formulation (3.20) has two main differences compared to the strong formulation (3.17). The first is that any solution of (3.17) solves (3.20) while the space of solutions of the weak form is larger since only first derivatives need to be defined, which lead to the terms weak and strong respectively. In addition, a weak solution is defined only up to a cardinal number of zero; hence changing the values of a given solution  $\Psi(\mathbf{r})$  along a finite number of planes in three dimensions yields a function that still solves (3.20). These properties play an important role here since they allow *e.g.* for piecewise linear basis functions as they are commonly used in the FEM scheme which do not have a second derivative and whose first derivative is undefined on areas with a cardinal number of zero.

To obtain the finite element formulation, one does not work on the whole space of differentiable and quadratically integrable functions defined on  $\Gamma$  but restricts the space to subspace spanned by a finite set of test functions for both  $\Psi(\mathbf{r})$  and  $\Phi(\mathbf{r})$ , leading to the Petrov-Galerkin scheme. To get best flexibility for the wave function, the domain  $\Gamma$  is subdivided into finite volume elements with a close packing. These volume elements, denoted as finite elements, give this method its name. Having this subdivision of space, the ansatz (basis) functions  $\varphi(\mathbf{r})$  used to construct  $\Psi(\mathbf{r})$  and  $\Phi(\mathbf{r})$  are defined as piecewise

polynomials being non-zero only on one or few elements.

Considering  $\Gamma$  to describe a plane, the elements can be for example triangles as shown in Figure 3.4. In this plane the vertices of the triangles can be enumerated as  $\mathbf{r}_i i = 1, \dots, N$ , leading to the definition of piecewise linear basis functions

$$\varphi_i(\mathbf{r}_j) = \begin{cases} 1 & j = i \\ 0 & j \neq i \end{cases} \quad (3.22)$$

resulting in two dimensional “hat” functions as depicted in Figure 3.4.

The so defined basis ensures continuity of the solution over  $\Gamma$  while the first derivatives are continuous only within each element. To assure continuity of the first derivatives as well, second order polynomials are needed.

As usual in variational schemes, the wave function of interest is a linear combination of these basis functions  $\Psi(\mathbf{r}) = \sum_i c_i \varphi_i(\mathbf{r})$  where  $c_i$  are weighting coefficients. The test functions are chosen from the same space of ansatz functions, testing each basis separately  $\Phi(\mathbf{r}) = \varphi_j(\mathbf{r}), j \in 1, \dots, N$ . Hence, the SE (3.20) is rewritten as

$$\sum_i \int d\mathbf{r} \left( \frac{1}{2} (\nabla \varphi_i(\mathbf{r})) (\nabla \varphi_j(\mathbf{r})) + V(r) \varphi_i(\mathbf{r}) \varphi_j(\mathbf{r}) \right) c_i = E \sum_i \int d\mathbf{r} \varphi_i(\mathbf{r}) \varphi_j(\mathbf{r}) c_i \quad \forall j \in \{1, \dots, N\} \quad (3.23)$$

which is the  $j$ -th component of the generalised matrix eigenvalue problem

$$\left( \frac{1}{2} \mathbb{A} + \mathbb{V} \right) \mathbf{c} = E \mathbb{M} \mathbf{c}. \quad (3.24)$$

with

$$\mathbb{A}_{ij} = \int d\mathbf{r} (\nabla \varphi_i(\mathbf{r})) (\nabla \varphi_j(\mathbf{r})) \quad \mathbb{V}_{i,j} = \int d\mathbf{r} V(r) \varphi_i(\mathbf{r}) \varphi_j(\mathbf{r}) \quad \mathbb{M}_{i,j} = \int d\mathbf{r} \varphi_i(\mathbf{r}) \varphi_j(\mathbf{r}) \quad (3.25)$$

and the vector  $\mathbf{c}$  in equation (3.24) contains the coefficients  $c_i$  to be found.

The quality of this basis depends strongly on the size and shape of the finite elements: The stronger the solution varies, the smaller be should the elements, or it should have higher order ansatz functions to be able to represent the wave function. Knowledge about




Figure 3.4: Example of finite elements in 2D of triangular shape and linear basis functions  $\varphi_i(\mathbf{r})$  defined on them.

ranges of sharp structures and areas with smooth variation of the wave function  $\Psi(\mathbf{r})$  hence is crucial for the setup of a good mesh.

In the next chapter more details about the finite element formulation will be given with a special focus on how to setup a grid that is well suited for describing free particles in quantum mechanics.

## 3.6 Hybrid Methods

In addition to the three methods presented above also mixed forms of these exists, utilizing the advantages of the respective ideas. One of the most prominent combinations of the above-mentioned methods is the finite element discrete-variable representation (FE-DVR) [91, 121]. Therein a DVR scheme is used with a set of ansatz functions each defined on a given interval only, the finite element. A single function connects these intervals respectively, similarly to the hat functions in FEM. Using this approach, more flexibility is obtained by varying the size of the elements or the number of basis functions per element. Moreover, the usually dense matrix of DVR schemes changes to a block-structure, each coupled by only one row [91].

Another scheme resulting in a similar description to the above-mentioned which is used by several authors on quantum mechanical problems is the spectral element method [128]. In contrast to the FE-DVR-scheme it is based on the weak formulation and uses Chebychev or Lagrange polynomials and hence allows for larger elements than usual FEM. While the spectral elements are found to be well suited for linear problems they require a complex implementation and lead to denser system matrices. The larger density in general leads to a reduced numerical stability and is computationally more expensive. Finally, complex geometries usually benefit more from smaller elements than from higher orders [129].

A successfull variety of spectral elements and the finite element method is the so-called spectral difference method. In this scheme, a spacially uniform grid is used (as usual in the finite difference scheme) while employing a stepwise defined pseudospectral basis. The appealing properties of this method are an exponential convergence as known from finite difference methods by employing the sparsity of the equation system obtained in FEM [130].

More advanced methods based on the above-mentioned schemes are available as well [131, 132] but are not described in more detail for brevity.

## 3.7 Wavelets

The wavelet method was developed in the 1980s [133] and combines the advantages of the Fourier method and FEM by decomposing objects into features of different scales but using a local basis [133–135]. Even though wavelets posess very advantagous features, they are not very well-established untill today. The most prominent and widely distributed application of Wavelets is compression of data as used for example in the jpeg2000 standard [136].

In general, a wavelet is a basis of the function space  $L_p(\mathbb{R}^d)$  *i.e.*, the space of functions  $\phi(\mathbf{r})$  for which  $\int \phi(\mathbf{r})^p d\mathbf{r} < \infty$  where  $\mathbf{r} \in \mathbb{R}^d$ . A particular wavelet is defined by a finite




Figure 3.5: Daubechies wavelets of the orders 1, 2 and 7 (from left to right) [138].

set of orthonormal mother wavelets  $\varphi_i(\mathbf{r})$  for which

$$\varphi_{i,j,\alpha}(\mathbf{r}) = m^{\frac{j}{2}} \varphi_i(\mathbb{M}^j \mathbf{r} - \alpha) \quad (3.26)$$

with the expanding scaling matrix  $\mathbb{M}$  (*i.e.* all eigen values have modulus larger than one),  $m = |\det(\mathbb{M})|$  and the running indices  $j \in \mathbb{Z}$   $\alpha \in \mathbb{Z}^d$  span  $L_p(\mathbb{R}^d)$  [134]. In practice, this very general definition is often restricted by additional requirements such as orthogonality of  $\varphi_{i,j,\alpha}(\mathbf{r})$  and only one mother-wavelet  $\varphi(\mathbf{r}) = \varphi_i(\mathbf{r})$  is used. For brevity, in the following only the space  $L_2(\mathbb{R})$  with  $\mathbb{M} = 2$  is considered which corresponds to the initial definition of wavelets [137]. The most prominent representatives are the Haar-wavelet and Daubechies wavelets which are orthogonal functions with finite support as depicted in Figure 3.5.

Another set of orthogonal wavelets with compact support are the Daubechies wavelets that are a series with increasing smoothness and support. The Haar-wavelet, shown in Figure 3.5 on the left, can be seen as the first order Daubechies-wavelet. In contrast to most other bases, here no smoothness properties are required but are helpfull of course since the functions to be represented usually are smooth [138].

Besides their broad use in data compression and analysis, wavelet bases also provide a handy tool to solve partial differential equations and are found to have similar numerical properties as the FEM [135]. When applying a wavelet basis to solve a partial differential equation, the weak form (3.20) is used, leading to a similar description as in the FEM scheme [134]. However, not all wavelets provide a valid basis for partial derivatives since, for example, the Haar-wavelet has a constant derivative of zero and the Daubechies-wavelet 2 (shown in Figure 3.5) is only semi-differentiable at some points [139]. At the same time hand the generality of the wavelet formulation allows to use problem-specific bases that are *e.g.* especially adapted to certain pseudo-differential operators, increasing the convergence properties [134]. Finding such optimised bases in a systematic way is under current research but one can expect that they could obtain numerically stable, fast and accurate partial derivative solvers [134, 140].

# 4 | Finite Element Methods

As a conclusion of the previous chapter one can see that the number of methods that are currently available to describe a free electron function in presence of an intricate electrostatic background potential is not that large. In this work the method of choice to model the free electron function is the FEM which had been applied to quantum mechanical problems already by several authors [141–144], however, to the best of my knowledge, only to bound state problems so far. A brief review about these works is given in section 4.1. Besides its large flexibility and computational efficiency pointed out in section 3.5 already, the large amount of available libraries for FEM [54, 145–148] is another advantage of practical importance due to the complexity of the generation of a suitable mesh, assembling of matrices and solution of matrix equations.

In the following chapter the integration of the matrix elements (section 4.2) and set up of the mesh (section 4.3) will be described. Thereby the focus is put on the application to the one-particle SE that is to be solved with molecular electrostatic potential. Since the interest thereby is on free particle solutions, the spectrum is expected to be very dense and the wave function to be delocalised, requiring for well-designed boundary conditions. A discussion of various boundary conditions and asymptotic descriptions available for FEM is described in section 4.4.

## 4.1 Finite Element Calculations in Quantum Chemistry

The FEM is mainly known from engineering disciplines where it is used in a broad range of applications such as modelling of fluids [149, 150], heat transfer and flow [151–153] or material deformation under mechanical stress [154, 155]. However, also several different quantum chemical problems have been solved with this method: SE solvers for small systems such as light atoms [141, 156–159] or diatomics [160], vibrational model systems [142] and solid state problems [161, 162]. Moreover, even Hartree-Fock [143] and DFT calculations on systems up to the size of benzene [144, 163, 164] have been performed, yielding results comparable to those obtained by the usual linear combination of atomic orbitals (LCAO) approach.

The above-mentioned publications have shown that the FEM is able to obtain reasonable results for molecular systems where the errors were comparable to those obtained with standard quantum-chemistry schemes even though their computational costs are higher. This suggests that the FEM is a good tool for computations in the field of quantum chem-

istry for going beyond the capabilities of the established schemes such as the description of unbound states.

## 4.2 Integration of Matrix Elements and Formulations of the Equation System

In section 3.5 the basics of the FEM were described and the generalised eigen system shown in eq. (3.24) to solve the SE was derived. Here this is taken as starting point and a closer look at the computation of the matrix elements as well as solving strategies for the large sparse generalised eigen problems are taken.

The generalised eigenproblem as given in eq. (3.24) consists of three matrices. Since the ansatz functions (shape functions)  $\varphi_i(\mathbf{r})$  have only a small support, most of the matrix elements are zero. However, in two and three dimensions no distinct band structure is achievable and the matrices are irreducible. Thereby matrix elements are zero when the elements involved are not neighboured. The computation of the non-zero matrix elements involve an integration as *e.g.* the overlap integral  $\mathbb{M}_{i,j} = \int d\mathbf{r} \varphi_i(\mathbf{r}) \varphi_j(\mathbf{r})$  of ansatz functions,. Since these functions are the same for all elements, the evaluation of these integrals can be done via a lookup-table or an efficient numerical integration scheme whose required order is well-known and need only be multiplied by the Jacobian of the respective elements involved. The matrix elements  $\mathbb{A}_{i,j} = \int d\mathbf{r} (\nabla \varphi_i(\mathbf{r})) (\nabla \varphi_j(\mathbf{r}))$  consist similar to those of  $\mathbb{M}$  of overlap integrals of known functions. The only matrix containing system-specific information is the potential  $\mathbb{V}_{i,j} = \int d\mathbf{r} \varphi_i(\mathbf{r}) V(\mathbf{r}) \varphi_j(\mathbf{r})$  which requires numerical integration by which  $V(\mathbf{r})$  is approximated as a spline of given order.

After assembling the matrices the eigenpair  $(e_i, \mathbf{c}_i)$  of the system

$$\left( \frac{1}{2} \mathbb{A} + \mathbb{V} \right) \mathbf{c}_i = e_i \mathbb{M} \mathbf{c}_i \quad (4.1)$$

need to be found where  $e_i$  should be closest to the analytic value of the kinetic energy of the photoelectron. Since matrix eigenvalue equations with several thousands of dimensions occur in many fields, numerous schemes have been developed to solve them efficiently [165–168], a selection of them is described in section 4.5.3. Despite the numerical complexity due to the high dimensionality of this problem (several thousands of basis functions) the second problem is due to the fact that the eigenenergies  $e_i$  are expected to be close to each other since the corresponding analytical problem has a continuous spectrum in this range. It is well known in numerics that this leads to instabilities especially for the eigen vectors, making a regularisation of the problem (described in section 4.5.2) indispensable.

## 4.3 Element Types and Mesh Types

Among the FEM formulations several ‘flavours’ were designed for different purposes. Given a certain equation to be solved in FEM there are in general two systematic ways to increase the accuracy. The first way is to increase the number of elements which is referred to as the *h*-FEM approach [41]. The refinement of the mesh is in principle always possible but technically demanding since it is not known in which regions of a mesh are too coarse in

general [41]. To overcome this, some FEM implementations such as that of `Libmesh` [54] which is used here, provide an adaptive mesh refinement scheme, iteratively refining the mesh using local error estimations and thus fully automatic [54]. But these schemes are numerically demanding and hence can be only applied to benchmark systems.

The second strategy is called  $p$ -FEM. In the  $p$ -FEM scheme, the order  $p$  of the test functions is increased, resulting in smoother and more flexible solutions. This scheme requires a large set of functions to be implemented but is known to yield good results if the function is smooth [169]. While standard FEM-libraries usually have only  $p = 1, 2$  implemented, others have high-order polynomials implemented [147]. Combined schemes where the grid-size as well as the element order are adapted is referred to as  $hp$ -fem [170].

Besides these different refinement strategies, finite element types also vary in their use of the degrees of freedom. While Lagrange elements are integrated by evaluating certain inner or boarder points, *e.g.* in Argyris or Hsieh-Clough-Tocher elements also first or even second derivatives are evaluated [37, 171]. Moreover, the arrangement of elements one distinguishes conforming and the more general non-conforming meshes [172]. In case of the latter, more general structures are supported as *e.g.* hanging nodes which occur when a vertex of one element is on the edge of an other as can be found in Figure 4.1a [37]. The setup of the mesh is, as mentioned above already, critical to the quality of the solution and hence of special importance. Moreover, it is technically non trivial to set up a close packing of volume elements with the desired properties in a systematic way. There are different element types available, each with their techniques for set up. Although in principle any element shape can be chosen, in three dimensions only tetrahedral (simplex), prism- and pyramid-shaped as well as hexahedral elements commonly are used. By choosing the element shape and the polynomial order, also the type of shape functions is defined.

When considering meshes to describe molecular properties it is clear that the element size should be smaller in the vicinity of the nuclei while it may be broader at larger distances. One way to create a hexahedral mesh with local refinement is to start with a coarse uniform lattice and subdivide the hexahedra where necessary as shown in Figure 4.1a for a benzene molecule. Another way is to setup small regular cubic grids around the nuclei and expand them radially in boxes of growing size as shown in Figure 4.1b. These brick-shaped elements however have the disadvantages that the regular cube-like structures therein might be not well-suited for atoms and molecules that rely rather on spherical shapes.

Another approach used by Lehtovaara *et al.* [163] is to put layers of polyhedra around the atoms with increasing number of points and radii. Thereby the overlapping regions of these spheres are removed by deleting elements that are closer to another atom. The mesh obtained with this procedure for a benzene molecule [144] is shown in Figure 4.1c.

When restricting oneself to tetrahedral elements, other design principles are possible: Since they are simplexes in three dimensions, they can be designed from general grids using *e.g.* Voronoi [113] or Delaunay [173] tessellations (the latter is described in section A.2 in appendix) from a set of points with the required properties. Son and Chu [112, 116] constructed sets of points resembling molecular geometries by inserting  $N$  spherical grids with different radii  $r_i$  around the atoms and cutting off the overlapping regions. Thereby




Figure 4.1: 2D cuts through 3D meshes for molecular systems obtained with different schemes for local refinement: (a) hexahedral elements adapted for the benzene molecules [144] (b) hexahedral mesh for a diatomic [143] (c) Polyhedral mesh for a benzene geometry [163] (d) Adaptive refined tetrahedral Mesh for ethylene. [143]

the respective radii are chosen as

$$r_i = \frac{il}{N - i + \frac{lN}{r_{\max}}} \quad i = 1, \dots, N \quad (4.2)$$

where  $r_{\max}$  is the radius of the largest sphere and  $l$  is a parameter smoothly changing between a linear  $l \rightarrow \infty$  and a  $1/r$ -mapping. As spherical grids they suggested the use of Lebedev-grids [174] and a design of Womersley [175, 176]. A more detailed discussion about the choices of grids will be given in section 5.2.1.

A more entangled method is used by Alizadegan *et al.* [143]. They start with an initial guess for the wave function and create a grid whose distances are inverse proportional to the second gradient of the electron density

$$d \propto \left[ \max \left\{ \left| \frac{\partial^2 \rho}{\partial^2 x} \right|, \left| \frac{\partial^2 \rho}{\partial^2 y} \right|, \left| \frac{\partial^2 \rho}{\partial^2 z} \right| \right\} \right]^{-1}.$$

which gives an estimate for the error due to linear approximation within each element. After solving the eigenvalue equation on this grid, they recompute another mesh on the basis of the new function, iterating this procedure several times. A cut through a mesh obtained by this procedure is shown in Figure 4.1d.

## 4.4 Boundary Conditions

Boundary conditions have not been addressed in this thesis so far for any of the methods but play an important role for the properties of the solution. Hence there is a large number of boundary conditions. The simplest case applicable here are Dirichlet-boundaries, requiring the wave function to vanish at the boundaries of the finite element box. In the FEM this condition can be applied especially simple by setting the coefficients of the outermost shape functions to zero.

Due to the large extend of the free particle, these boundaries however are unphysical if they are not applied at distances that are several times larger than the wavelength. However, considering a particle with 0.1 eV kinetic energy, its wavelength is 4.4 Angstroms and

the box would need to have a diameter of several tens of Angstroms which is not feasible any more while kinetic energies in the meV-range would lead to even worse scenarios.

Besides the numerical restriction to a finite box also mapping schemes can be used as *e.g.*  $x = \tan(\frac{y}{2})$  that maps  $[-\infty : \infty]$  to  $[-\pi : \pi]$  [119]. However, using such a mapping directly is infeasible since the oscillations of the wave function would become arbitrarily sharp in the mapped range and hence the representation of the FEF would still be poor.

#### 4.4.1 Complex Absorbing Potential

The complex absorbing potential (CAP) is a method often found in the literature when describing particles with infinite extent[61, 62]. In this scheme, an artificial potential usually of the form

$$W_{\text{CAP}}(\mathbf{r}) = \begin{cases} i\nu(\mathbf{r} - \mathbf{r}_0)^2 & \mathbf{r} > \mathbf{r}_0 \\ 0 & \text{else} \end{cases} \quad (4.3)$$

is added where  $\mathbf{r}_0$  is larger than the bound part of the system. Such a potential damps the reflection of the wave function at the boarders and thus enhances the quality of states obtained on a finite region [177, 178]. However, studies with different shapes of these potentials show that it influences the wave function not only close to the boarders and a proper design of the parameters is non-trivial [179].

To minimise the error due to the CAP, the parameter  $\eta$  can be chosen such that its dependency on the energy vanishes in first order, *i.e.*  $\eta \frac{dE}{d\eta} = 0$  [65, 179]. Moreover, due to non-hermiticity the usual scalar product is not suitable when calculating overlap integrals any more. Too strong  $\eta$  makes reflections, too weak  $\eta$  lead to unstable resonances, making them strongly basis-set dependent [179]

The idea of damping down a function and making sure no reflections are scattered back to the region of interest is common to further approaches as *e.g.* absorbing boundary conditions [180], perfectly matched layer schemes [181–183] or certain variable transformations [91].

- Also read zotero: (In)FEM AbsorbBoundariesVS\_InfFE
- transparent bc!? [184]

#### 4.4.2 Mode-matching Schemes

Consider the solution in an inner and outer region as different variables following the equations

$$\nabla^2 \Psi_1 + V(\mathbf{r}) \Psi_1 - E \Psi_1 = 0 \quad \nabla^2 \Psi_2 - E \Psi_2 = 0 \quad (4.4)$$

whereby the outer function needs to satisfy the Sommerfeld condition  $r^\alpha \left( \frac{\partial \Psi_2}{\partial r} - ik\Psi_2 \right) \rightarrow 0$ . These equations are coupled by the conditions

$$\Psi_1 = \Psi_2 \quad \nabla \Psi_1 \mathbf{n} = \Psi_2 \mathbf{n} \quad (4.5)$$

to ensure continuity of the solution and the gradient normal to the boundary [185]. Thereby, the asymptotic behaviour of the outer function is ensured by taking basis functions that individually fulfil the condition and the boundaries enter the weak formulation since the application of Greens' theorem leads to an extra term.

The mode-matching scheme can be considered as a generalisation of the R-matrix approach in a finite element formulation.

#### 4.4.3 The Boundary Element Method

The boundary element method (BEM) can be used as a self-standing method for solving partial differential equations using the weak formulation [186, 187]. In its pure form, the BEM solves the problems using only conditions given at its boundaries that are connected to the volume properties via Greens' theorem, the Gauss-Ostrogradskii (divergence) theorem as well as Stokes theorem [188]. Even though the BEM procedure suffers strongly from the restriction of being applicable only to linear systems for which a fundamental solution is known as well as the disadvantage of leading to dense, unsymmetric matrix equations [187] it has some popularity until these days [189–191]. But its main advantages come into play when being used with the FEM [192] where the FEM can be used to obtain an accurate solution in the inner region and the boundaries are treated with the BEM. Considering an unbound domain as, *e.g.* the problem of the outgoing electron, the infinite domain  $\Gamma$  can be divided into a finite region  $\Gamma_i$  where the atoms electrostatic potential leads to ... and the remaining domain  $\Gamma_o$  in which the time-independent SE reduces to the Helmholtz-problem whose fundamental solutions are well-known and thus the BEM is applicable [187, 193].

#### 4.4.4 Infinite Elements

The infinite element approach was developed in the 1980-ths for acoustical caluculations and is specificly designed for the Helmholtz equation and can be understood as a advancement of the BEM that is speciallised for the Helmholtz equation. The general idea of the infinite elements is that the solution of the radial Helmholtz equation in spherical symmetry is well-known to be of the form

$$\Psi(r) = \left( \frac{a}{r} + \frac{b}{r^2} + \dots \right) e^{ikr} \quad (4.6)$$

where  $k = |\mathbf{k}|$  is the absolut value of the momentum of the particle and the prefactors thus correspond to different angular momenta of the outgoing electron. In the complete limit any function fulfilling the Sommerfeld radiation condition [42] can be represented.

To use this asymptotic information, in the infinite element region a layer of elements is set onto the outer surface of the finite element region in which the shape functions are of the form (4.6). To fulfill the continuity conditions, the front-face of these elements coincides with the outer face of the respective finite element. while their radial faces have ray-like edges with a common centre in the middle of the finite element region.

Over the decades several formulations of infinite elements were developed which each make an ansatz for the solution of the form

$$\Psi(\mathbf{r}) = \varphi(\mathbf{r}) e^{ik(r-r_0)}, \quad (4.7)$$

where  $\varphi(\mathbf{r}) = f(r)\varphi_2(\mathbf{r})$  is a product of a transversal shape function  $\varphi_2(\mathbf{r})$  which is orthogonal to the radial direction  $r$ . The function  $f(r)$  is a multipole expansion *i.e.*

a polynomial in  $\frac{r_0}{r}$  where Jacobi-polynomials turned out to be most stable [194]. The differences in the formulations are introduced by the test functions applied [40].

## Burnett Elements

In the original Burnett formulation, the test functions are given in the form

$$\Phi(\mathbf{r}) = \varphi(\mathbf{r})e^{-ik(r-r_0)}. \quad (4.8)$$

In the literature, usually the sign in the exponent is changed due to the use of an other convention in the notation of the weak formulation 3.20. Important is, however, that in the equation later on the test and ansatz functions enter both with same oscillatory factor, introducing oscillations with  $e^{2ik(r-r_0)}$  in the Hamiltonian. These oscillations make the Hamiltonian non-hermitian and introduce a strong dependence on the energy  $E = \frac{k^2}{2}$ . Moreover it was found that this formulation can lead to numerical instabilities [40] and a wrong asymptotic behaviour.

## Conjugate Burnett Elements

To get rid of the oscillatory terms, a modification of the test functions applied was introduced such that, in the notation used within this thesis, the test functions coincide with the space of ansatz functions, having the form

$$\Phi(\mathbf{r}) = \varphi(\mathbf{r})e^{ik(r-r_0)}. \quad (4.9)$$

An important disadvantage of this formulation is, however, that in this formulation integrals with infinite contribution arise.

## Astley-Leis Formulation

To regularise the matrix elements, an additional damping is needed. Since the ansatz function (4.7) should not be changed for physical reasons, Astley suggested to chose test functions of the shape

$$\Phi(\mathbf{r}) = D(r)\varphi(\mathbf{r})e^{-ik\mu(r)} \quad (4.10)$$

where in three dimensions  $D(r) = \frac{1}{r^2}$  [39]. This choise of basis functions is done at the expense that the Hamiltonian (3.25) is not hermitian.

Even though the oscillating term  $e^{ikr}$  cancels out, the Hamiltonian is energy dependent and thus the generalised eigenvalue problem (3.24) formulated in chapter 3.5 changes to

$$\mathbb{A}\mathbf{c} + ik\mathbb{B}\mathbf{c} - k^2\mathbb{C}\mathbf{c} = 0 \quad (4.11)$$

with

$$\mathbb{A}_{i,j} = \int \left( V(\mathbf{r}) D(r) \varphi_i(\mathbf{r}) \varphi_j(\mathbf{r}) - \frac{1}{2} D'(r) \varphi_i(\mathbf{r}) \varphi'_j(\mathbf{r}) + \frac{1}{2} D(r) \varphi'_i(\mathbf{r}) \varphi'_j(\mathbf{r}) \right) d\mathbf{r} \quad (4.12)$$

$$\mathbb{B}_{i,j} = \frac{1}{2} \int (-\mu'(r) D' \varphi_i(\mathbf{r}) \varphi_j(\mathbf{r}) + D(r) (\varphi'_i(\mathbf{r}) \varphi_j(\mathbf{r}) - \varphi_i(\mathbf{r}) \varphi'_j(\mathbf{r}))) d\mathbf{r} \quad (4.13)$$

$$\mathbb{C}_{i,j} = \frac{1}{2} \int ((D(r) \mu'(r) \mu'(r) + 1) D(r) \varphi_i(\mathbf{r}) \varphi_j(\mathbf{r})) d\mathbf{r} \quad (4.14)$$

where the relation  $E = \frac{1}{2}k^2$  is used [41].

This reformulation though lead to a quadratic eigenvalue problem instead of the generalised eigenvalue problem from before. However, assuming that the difference between the eigenvalues of (4.11) and the target energy of the outgoing electron is small, the quadratic eigenvalue problem can be approximated by the generalised eigenvalue problem (3.24) by setting  $k$  to the respective target momentum in the Hamiltonian. However, this approximation, which is applied here for simplicity, needs to be verified.

The nonhermiticity of this Hamiltonian is, however, an important drawback of this formulation. The introduced imaginary part of the eigenvalue can be considered as a broadening of the line width and thus as a lifetime as well as an uncertainty of the energy. However, this is not very meaningful since the ansatz is done in particular in such a way that the wave function is not changed in an unphysical sense. To understand the respective results, here a symmetrised ansatz is suggested in the following.

### Symmetrised Formulation

Besides those formulations mentioned above, further infinite element formulations are used [41]. For example, a formulation to the Burnett-elements using a quadratic decay was applied to the harmonic oscillator [51] and DFT calculations [50] successfully. However, the application of infinite elements to FEFs is to my knowledge a novelty in this work.

Since the Astley-Leis elements suffer from their non-symmetric space of ansatz- and test-functions whereas the conjugate Burnett formulation has problems due to infinitely large matrix elements, here a symmetric but damped ansatz of the form

$$\Psi(\mathbf{r}) = D(r)^p \varphi(\mathbf{r}) e^{ik\mu(r)} \quad (4.15)$$

$$\Phi(\mathbf{r}) = D(r)^p \varphi(\mathbf{r}) e^{ik\mu(r)} \quad (4.16)$$

is suggested where  $\varphi(\mathbf{r})$  is of the form as before but  $D(\mathbf{r}) = \frac{1}{r^2}$  with  $p > 0$  ensures that the matrix elements are finite-valued.

## 4.5 Solving Large Eigenvalue Problems

In finite element applications such as those being proposed in this work, matrix equations with hundreds up to hundredthousands of dimensions need to be solved. This requires elaborate strategies, using the sparsity of these matrices.

The focus here is on solving the generalised eigenvalue problem (3.24) and the quadratic problem (4.11) respectively. However, efficient strategies are only known for regular eigen-

value problems of the form

$$\mathbb{A}\mathbf{x} = \lambda\mathbf{x}. \quad (4.17)$$

Hence, the more general forms will be rewritten to become (4.17) as discussed in sections 4.5.1 respectively. Moreover, since the state of interest is a free state, one needs to expect a high density of states for an appropriate mesh. This, however, is a well-known problem in numerical methematics since almost degenerate eigenvalues are very sensitive to small perturbations and their respective eigenvectors even more. Though section 4.5.2 addresses these problems and a way for numerical stabilisation is sketched.

Finally in section 4.5.3 a few methos are presented showing how a small number of approximate eigenpairs can be obtained in a numerically efficient way from the usual eigenproblem (4.17).

#### 4.5.1 Generalised Eigenproblem

The most straight-forward way to reformulate the generalised eigenvalue problem

$$\mathbb{A}\mathbf{x} = \lambda\mathbb{B}\mathbf{x}, \quad (4.18)$$

is to invert the matrix  $\mathbb{B}$ , obtaining the regular Eigenproblem  $\mathbb{B}^{-1}\mathbb{A}\mathbf{x} = \lambda\mathbf{x}$ . This way is possible as long as  $\mathbb{B}$  is invertable and not too large since inversion is a demanding task and the resulting matrix is not sparse anymore [195].

To prevent the use of dense matrices, libraries often do not operate with the matrices themselves but rather with a set of vector on which these matrices act [195]. The most popular scheme of this kind is the Rayleigh-Ritz projection where the initial problem is approximated on a small subspace  $\mathcal{V}_j = \text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_j\}$ , spanned by appropriate vectors  $\mathbf{v}_i$ .

Projecting the original problem (4.18) onto this subspace yields the new system  $\Sigma_j \mathbf{s} = \theta \Theta_j \mathbf{s}$  where  $\Sigma_j = \mathbb{V}_j^T \mathbb{A} \mathbb{V}_j$  and  $\Theta_j = \mathbb{V}_j^T \mathbb{B} \mathbb{V}_j$  respectively which is only of dimensionality  $j$ . The matrix  $\mathbb{V}_j$  is unitary with the rows  $(\mathbb{V}_j)_i = \mathbf{v}_i$ . After solving this dense but small problem, the original eigenvectors can be approximated as  $\mathbf{x}_j = \mathbb{V}_j \mathbf{s}_j$  and  $\lambda = \theta_j$ . The obtained eigenpair is a good approximation to the actual one as long as the subspace  $\mathcal{V}_j$  contains the respective solution or contains a vector which is at least close to it. A commonly used approach the Krylov subspace.

#### 4.5.2 Stabilisation of Eigenproblems

Independent of the efficiency and robustness of the eigensolver in use, seeking solutions of the SE for free particles means that eigenpairs are to be found whose energies are, if the numerical parameters are chosen well, very dense or even degenerate. Unfortunately, dense-lying eigenvalues lead to numerical difficulties and especially the eigenvectors are known to be unreliable in this case. In practice this means that the iterative schemes do not converge anymore, requireing a reformulation of the mathematical probelm.

One way to circumvent the instabilities in the original problem (4.18) is to reformulate it as a minimisation problem [25]. Therefore eq. (4.1) is rewritten as  $(\frac{1}{2}\mathbb{A} + \mathbb{V} - E\mathbb{M}) \mathbf{c}_i = 0$  where  $E$  is the target energy. Since this will have most likely no unambiguous solution,

one minimises the residuum

$$\min_{\|\mathbf{x}\|=1} \left\{ \left\| \left( \frac{1}{2} \mathbb{A} + \mathbb{V} - E \mathbb{M} \right) \mathbf{x} \right\| \right\}. \quad (4.19)$$

Using the  $L_2$ -norm, this equivalent to finding the smallest eigenvalue of

$$\left( \frac{1}{2} \mathbb{A} + \mathbb{V} - E \mathbb{M} \right)^\dagger \left( \frac{1}{2} \mathbb{A} + \mathbb{V} - E \mathbb{M} \right) \mathbf{x}_i = \theta \mathbf{x}_i \quad (4.20)$$

where  $\theta$  is a measure for the error in energy.

Whereas eq (4.20) is only an approximation to the original problem, it has the advantage that it is a quadratic expression. Similar schemes which are, in contrast to that mentioned above, only reformulations of the original problem are often referred to as spectral transformations [195]. One example is the harmonic extraction, where the eigenvalues of original expression (4.18) are first shifted such that the target value is 0, leading to the equation

$$(\mathbb{A} - \tilde{\lambda} \mathbb{M}) \mathbf{x} = (\lambda - \tilde{\lambda}) \mathbf{x} \quad (4.21)$$

where  $\tilde{\lambda}$  is the target value of the original problem. Then, the equation is multiplied by  $\mathbb{A}$ , resulting in the problem

$$(\mathbb{A} - \tilde{\lambda} \mathbb{M}) (\mathbb{A} - \tilde{\lambda} \mathbb{M}) \mathbf{x} = (\lambda - \tilde{\lambda}) (\mathbb{A} - \tilde{\lambda} \mathbb{M}) \mathbb{M} \mathbf{x} \quad (4.22)$$

which is observed to lead to faster convergence for Krylov-subspace methods [195].

Further spectral transformations are the spectral folding [195] where the left and right side of equation (4.21) are squared respectively which often leads to higher stability but the squaring of eigenvalues leads to ambiguities with respect to the eigenvalues  $\lambda$  of the original problem. The scheme that turned out to be most beneficial for the matrices used in this work is the shift and invert scheme where the inverted problem of eq. (4.21), *i.e.*

$$(\mathbb{A} - \tilde{\lambda} \mathbb{M})^{-1} \mathbb{M} \mathbf{x} = \frac{1}{\lambda - \tilde{\lambda}} \mathbf{x}, \quad (4.23)$$

is solved. The formulation (4.23) has the advantage that the transformed eigenvalues  $\frac{1}{\lambda - \tilde{\lambda}}$  are well-separated and on the extrema of the new spectrum, making the convergence faster and more stable [196]. A more generalised transformation is the Cayley-transform which has two parameters  $\tilde{\lambda}$  and  $\tilde{\tilde{\lambda}}$  and yields the usual eigenvalue problem

$$(\mathbb{A} - \tilde{\lambda} \mathbb{M})^{-1} (\mathbb{A} + \tilde{\tilde{\lambda}} \mathbb{M}) \mathbf{x} = \frac{\lambda + \tilde{\tilde{\lambda}}}{\lambda - \tilde{\lambda}} \mathbf{x}. \quad (4.24)$$

Finally, a regularisation of eigenvalue problems can be performed using matrix preconditioners. Many different types of preconditioners are developed [197] but most of them are effective only on a small class of problems as *e.g.* especially well-suited for matrices originating from FEM-problems [198, 199] and can be even specialised to a FEM structure [200, 201] or for a particular physical problem [184]. Often their performance moreover depends on the solver to be used later [202]. This makes their effective use more intricate

and can easily lead to higher costs than their benefit is. In this thesis, the shift-and-invert scheme (4.23) is used.

### 4.5.3 Solving Large Eigenproblems

For the computation of eigenpairs, many classes of solvers have been developed with various numerical properties. Besides direct solvers such as the Gauß-elimination, many iterative solvers have been developed that are especially well-suited for large but spares problems. Besides the famous Jacobi- and Gauß-Seidel algorithms which converge only in certain cases, also the Davidson method and several Krylov subspace methods are commonly used. For finite element problems, the Krylov subspace solvers turned out to be the most efficient class.

As discussed in the sections 4.5.1 and 4.5.2 already, the solution of a generalised eigenvalue problem involves matrix operations which need to be avoided to keep their sparse structure.

As a popular choise for such classes of problems, the Krylov subspace is used. An  $r$ -dimensional Krylov subspace is generated by a vector  $\mathbf{x}$  and a matrix  $\mathbb{A}$  and has the form

$$\mathcal{K}_r(\mathbb{A}, \mathbf{x}) = \text{span} \{ \mathbf{x}, \mathbb{A}\mathbf{x}, \mathbb{A}^2\mathbf{x}, \dots, \mathbb{A}^{r-1}\mathbf{x} \}. \quad (4.25)$$

If  $\mathbb{A}$  is sparse, the evaluation of these expressions is only of order  $\mathcal{O}(d)$  where  $d$  is the dimensionality of  $\mathbf{x}$ . The vectors obtained with large powers of  $\mathbb{A}$  however usually become more and more linearly dependent. To prevent this, the vectors usually are orthonormalised subsequently.

An important issue in this scheme is a good choise for  $\mathbf{x}$  which crucially determines the speed of convergence. If a reasonable start-vector is not given, the space  $\mathcal{K}_r$  needs to be extended by increasing  $r$  iteratively. The orthogonalisation method being used distinguishes different Krylov subspace methods such as the Arnoldi [203] or Lanczos [204]. In the particular implementation, the Krylov-Schur algorithm is used [196, 205]. To keep the dimensionality low, in this algorithm the subspace iteration is restarted after  $r$  reached a certain value, starting with a better guess  $\mathbf{x}$ .

# 5 | Applied Protocol

After describing different approaches to compute PESs in chapter 2 and pointing out why the DO formalism combined with a finite element method for the computation of the photoelectron function is well-suited for our needs, in this chapter the protocol as being used to obtain the PESs is presented.

In the first section the computation of the wave functions of initial (unionised) and final (ionised) states will be described. Thereafter, in section 5.2.1 the setup of the finite element system that is used to compute the free electron function as well as the dipole matrix element will be described. In the final section, the computation of Dyson orbitals will be sketched and it will be described how to transfer it to the finite element setup.

## 5.1 Bound State Functions

The DO formalism as it is described in 2.3 can be used with any self-consistent field method which can predict excited state energies and wave functions. In this work, density functional theory (DFT) is used for ground state calculations and its time dependent counterpart TDDFT to compute excited state energies. The DFT formalism which is introduced in the sections 5.1.1 and 5.1.2 has shown to be accurate and numerically cheap.

### 5.1.1 Ground State Density

The DFT is formally based on the Hohenberg-Kohn theorems [43] which state that the ground-state electron density determines the potential in the SE and with this also the wave-function. Moreover, the second Hohenberg-Kohn theorem states that the energy of the ground state can be found variationally from the density. Thus, the electron density contains all information needed and the computation of the  $N$ -electron wave function can be omitted. Since the Hohenberg-Kohn theorems do not point out a way how to determine the electron density without knowing the wave-function, usually the Kohn-Sham scheme [44] is used where the electrons are described as non-interacting particles in a respective pseudo-potential that is constructed such that the electron density of these Kohn-Sham orbitals corresponds to the real electron density. Since the particles do not interact with each other, the Kohn-Sham orbitals  $\Psi_j(\mathbf{r})$  can be obtained from the one-particle SE

$$\left( -\frac{1}{2} \nabla^2 + V_{\text{eff}}(\mathbf{r}) \right) \Psi_j(\mathbf{r}) = \epsilon_j \Psi_j(\mathbf{r}) \quad (5.1)$$

where  $\epsilon_j$  is the binding energy of the respective Kohn-Sham orbital and the effective potential

$$V_{\text{eff}}(\mathbf{r}) = V_{\text{ext}}(\mathbf{r}) + \int \frac{\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r}' + V_{\text{xc}}(\mathbf{r}) \quad (5.2)$$

can be separated into the external potential  $V_{\text{ext}}(\mathbf{r})$  which consists of the attractive nuclear electrostatic potential and external fields the electrostatic interaction of the particles where  $\rho(\mathbf{r}) = \sum_{i \neq j} |\Psi_i|^2$  and the exchange correlation potential  $V_{\text{xc}}(\mathbf{r})$ . The exchange correlation potential  $V_{\text{xc}}(\mathbf{r})$  contains, besides contributions from exchange and correlation, also the error of kinetic energy in the Kohn-Sham scheme to account for the fact that the Kohn-Sham wave-functions differ from the real orbitals and thus have another kinetic energy as well [206].

Assuming that the Kohn-Sham orbitals coincide with the physical orbitals, the exchange potential acts on an orbital  $\Psi_i(\mathbf{r})$  as

$$V_{x;j}(\mathbf{r})\Psi_i(\mathbf{r}) = \int \frac{\Psi_j^\dagger(\mathbf{r}')\Psi_i(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r}' \Psi_j(\mathbf{r}) \quad (5.3)$$

and thus is non-local [206]. In DFT usually the exchange energy is approximated as a local functional of the density and its derivative, leading to the so-called local density approximation and gradient corrected functionals. The Becke [207] exchange-functional used in this work has the form

$$E_x = \frac{3}{2} \left( \frac{3}{4\pi} \right)^{\frac{1}{3}} \sum_{\sigma} \int \rho_{\sigma}(\mathbf{r})^{\frac{4}{3}} d^3\mathbf{r} - \beta \sum_{\sigma} \int \rho(\mathbf{r})^{\frac{4}{3}} \frac{x_{\sigma}(\mathbf{r})^2}{1 + 6\beta x_{\sigma} \sinh^{-1}(x_{\sigma}(\mathbf{r}))} d^3\mathbf{r} \quad (5.4)$$

where the first summand corresponds to the local density approximation and the second is a semi-empirical gradient correction,  $\sigma$  denotes the spin polarisations and  $x_{\sigma}(\mathbf{r}) = \frac{|\nabla \rho_{\sigma}(\mathbf{r})|}{\rho_{\sigma}^{\frac{4}{3}}}$ . The gradient correction in this functional is constructed such that the asymptotic behaviour of the exchange energy and electron density follow

$$\lim_{r \rightarrow \infty} E_x^{\sigma}(\mathbf{r}) = -\frac{1}{|\mathbf{r}|} \quad (5.5)$$

$$\lim_{r \rightarrow \infty} \rho(\mathbf{r}) = e^{-a_{\sigma}|\mathbf{r}|} \quad (5.6)$$

with  $a_{\sigma}$  is a constant related to the ionisation potential of the system under study [207]. The prefactor of the gradient correction,  $\beta$ , was determined by fitting to several atomic noble-gas systems and is set to be  $\beta = 0.0042$  in atomic units [207].

The correlation potential finally is defined as the functional derivative of the correlation energy with respect to the electron density  $V_c(\mathbf{r}) = \frac{\partial E_c[\rho]}{\partial \rho(\mathbf{r})}$ . In the case of the The LYP-correlation energy functional is based on the Colle-Salvetti formula which estimates the correlation energy in the Hartree-Fock approach and introduces two parameters that are set by fitting to experimental data [208].

### 5.1.2 Excited State Properties

The DFT is not only a well-established method to obtain ground-state properties of many different kinds of systems, but it has also a time-dependent counterpart, formally based on the Runge-Groß theorem [209]. Commonly, when referring to TDDFT, respective the linear-response scheme is meant [210]. It is a first order perturbation expansion and can be used to obtain the energies and occupation numbers of excited states using the so-called Casida-equation [211]

$$\begin{bmatrix} \mathbb{A} & \mathbb{B} \\ \mathbb{B}^\dagger & \mathbb{A}^\dagger \end{bmatrix} \begin{bmatrix} \mathbf{X} \\ \mathbf{Y} \end{bmatrix} = \omega \begin{bmatrix} \mathbb{1} & \mathbb{0} \\ \mathbb{0} & -\mathbb{1} \end{bmatrix} \begin{bmatrix} \mathbf{X} \\ \mathbf{Y} \end{bmatrix} \quad (5.7)$$

where the matrix elements are

$$\mathbb{A}_{ia,jb} = \delta_{ij}\delta_{ab}(\varepsilon_a - \varepsilon_i) + \int \int d\mathbf{r} d\mathbf{r}' \frac{\Psi_i(\mathbf{r})\Psi_a(\mathbf{r})\Psi_j(\mathbf{r}')\Psi_b(\mathbf{r}') - \Psi_i(\mathbf{r})\Psi_j(\mathbf{r})\Psi_a(\mathbf{r}')\Psi_b(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}, \quad (5.8)$$

$$\mathbb{B}_{ia,jb} = \int \int d\mathbf{r} d\mathbf{r}' \frac{\Psi_i(\mathbf{r})\Psi_a(\mathbf{r})\Psi_b(\mathbf{r}')\Psi_j(\mathbf{r}') - \Psi_i(\mathbf{r})\Psi_b(\mathbf{r})\Psi_a(\mathbf{r}')\Psi_j(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} \quad (5.9)$$

where  $i, j$  denote occupied and  $a, b$  virtual states and  $\mathbb{1}$  is the unity matrix,  $\varepsilon_\alpha$  are (Kohn-Sham) orbital binding energies of the Kohn-Sham orbitals  $\Psi_\alpha(\mathbf{r})$ . The indices  $i, j$  and  $a, b$  denote occupied and virtual orbitals with binding energy  $\varepsilon_i, \varepsilon_a$  respectively [210]. The Casida equation (5.7) is a non-hermitian eigenvalue problem whose eigenvalues  $\omega$  correspond to the transition energies and the eigenvectors denote the transition coefficients. The different blocks in equation (5.7) have different physical interpretation. As the negative sign of the transition energy in the lower block indicates, lower half of the eigen vector, denoted as  $\mathbf{Y}$ , corresponds to negative excitation energies  $\varepsilon_i$  and can be assigned to de-excitations from virtual orbitals which makes no physical sense if the reference state is a ground state [210]. Since the matrix  $\mathbb{B}$  contains usually only small elements, it is often set to zero which is referred to as Tamm-Dancoff approximation and reduces the original equation to a hermitian equation of half dimensionality [211].

### 5.1.3 OTRSH-scheme

One main issue concerned with the DFT-formalism is that the approximate exchange functional decays exponentially instead of  $\frac{1}{r}$  and  $\frac{1}{r^4}$  for the exchange and correlation terms respectively [45]. **But the Becke functional should have fixed this!?** This wrong behaviour affects the obtained wave functions and thus the orbital energies [17].

To reduce this error, so-called range-separated hybrid (RSH) functionals are used, in which the DFT-exchange term as *e.g.* the Becke functional (5.4) is used for small interelectronic distances only, while at larger distances, where correlation effects are not that important, the Hartree-Fock exact-exchange (5.3) is used. The interchange between the schemes is done by the scheme

$$\frac{1}{r} = \underbrace{\frac{\alpha + \beta \text{erf}(\omega r)}{r}}_{\text{exact exchange}} + \underbrace{\frac{1 - \alpha - \beta \text{erf}(\omega r)}{r}}_{\text{DFT exchange}} \quad (5.10)$$

where  $\alpha + \beta = 1$  and  $\omega$  are parameters to be chosen. Besides taking the standard parameters that are fitted to a set of testing-systems, ab-initio schemes for choosing  $\alpha$  and  $\omega$  are available which are referred to as optimally-tuned RSH (OTRSH) schemes. Using such a scheme, the degrees of freedom obtained are used to optimise a particular system-property such as Koopmans' theorem by minimising the functional [45]

$$J(\alpha_{\text{opt}}, \omega_{\text{opt}}) = \min_{\alpha, \omega} \{ |E_N(\alpha, \omega) - E_{N-1}(\alpha, \omega) - \varepsilon_{\text{HOMO}}| \} \quad (5.11)$$

or a similar one which can also account for lower-lying orbitals or the electron affinities which should resemble the binding energies of the lowest unoccupied molecular orbital (LUMO).

Another quantity to optimise is the derivative discontinuity [212].

For this optimization procedure the Gaussian package **G09** [53] is used with the 6-31G(d) [213, 214] basis set and the functional LC-BLYP [215]. The ground state DFT calculation, determination of geometries and the linear-response TDDFT calculations have been conducted with a locally modified version of **NWChem** [52], employing the basis set def2-tzvp [216] without symmetry restrictions. The Kohn-Sham orbitals (obtained by ground state DFT) and CI-coefficients (obtained by linear-response TDDFT which yields the configuration interaction singles expansion for a number of excited states) as well as the atomic overlap matrix are interfaced to the in-house software **DYSON** [31] that computes the DOs. The integration of the transition dipole matrix elements in eq. (2.16) finally is performed using **ezDyson** [33].

### 5.1.4 Computing the Dyson orbital

The DO is computed in the framework of this thesis with the in-house code **DYSON** [31].

## 5.2 Free electron function

The FEF is computed with a finite element scheme, using the program **Free Willy** developed in the framework of this thesis. It is based on the library **Libmesh** [54] which itself uses several libraries for the required linear algebra and the mesh-setup.

### 5.2.1 Setup of the Grid

The most crucial part of the FEM is the mesh under usage. In the program **FreeWilly** it is obtained from a set of points via Delaunay triangulation (see A.2 for details) using the library **tetgen** [217]. The distribution of these points is responsible for the quality of the obtained solution and depends on many parameters in a non-trivial way. The dependence is in particular on

**The molecular geometry** Close to the cores it should be denser, at larger distances a coarser grid is possible.

**The kinetic energy** of the photo electron which gives the maximum distance between two points

**The largest angular momentum of the Dyson orbital** determines the angular momentum of the photoelectron to be resembled.




Figure 5.1: Example of a mesh for the water-geometry. It consists of 5 spheres with constant number of points per sphere. The overlapping regions are cut off. Left: cut through the nuclear plane.

In the general scheme the generation of the point-distribution follows the suggestion of Son and Chu [116]: The grid is set up of atomic grids, *e.g.* spherical distributions, each centred at atoms where the overlapping regions are cut off. The radii of these spheres needs to be much larger than the bond lengths, holes or interatomic gaps in the molecule to (*e.g.* holes in aromatic rings or the space between ligands of larger molecules). Using this scheme, the grid-parameters are determined by the size of the largest sphere  $r_{\max}$ , number  $N$  of spheres used as well as the radial distribution of them and the angular distribution of points on these spheres.

For the radial distribution Son and Chu [112] suggested the scheme

$$r_i = \frac{il}{N - i + \frac{lN}{r_{\max}}} \quad i = 1, \dots, N \quad (5.12)$$

where  $l$  is a parameter to chose. As an alternative distribution, here the formula

$$r_i = \frac{il}{\left(\frac{N}{i}\right)^p \left(\frac{Nl}{r_{\max}} - 1\right) + 1} \quad i = 1, \dots, N \quad (5.13)$$

is to be tested where  $l$  and  $p \geq 1$  are parameters to chose whereby the condition  $N < \frac{r_{\max}}{l}$  should be fulfilled to prevent the singularity.

While the distribution of spheres follows, at least on a qualitative level, a clear scheme since it should always resemble the local kinetic energy, the distribution on the surface of each sphere is not that clear. Qualitatively it is preferable to chose a regular grid but this more challenging. Since the ... grid, which is known from the ... is clearly a bad choice due to its high density close to the poles, in geological applications often so-called geodesic grids are chosen.

Some explanations are given here:

- Lebedev-grid -*j* try to get spherical harmonics with highest L accurate.
- If also required: all weights should be equal: spherical t-designs.
- Leads directly to the question of uniform distribution.-*j* not directly by algorithms
- other approach: close packings
- energy minimisation: Consider nodes as interacting particles -*j* minimise E for some model potential.

see <http://people.maths.ox.ac.uk/beentjes/Essays/QuadratureSphere.pdf>. - crude but fast algorithms Since in finite element theory the sphere neither needs to be really round nor is there any global functional defined on it, the complicated distributions described above may be not even needed. An other approach therefore is to use an algorithm that gives just a more or less uniform distribution.

Approach used for climate models: so-called geodesic grids: subdivision of polyhedra, projected onto the sphere[218, 219], see also for further information.

The point sets I took are from here: [220, 221].

The above mentioned techniques show the large variety of different approaches and it is not very clear which one will meet our needs best. Moreover, here the problem is not only two dimensional but the whole sphere (not only its surface) needs to be subdivided. Hence, besides the question of a radial density of different spherical surfaces, also the number of points per sphere as function of the radial distance needs to be considered.

Whether the approach of subdividing the atomic meshes into radial and angular parts as opposed to another volume tessellation can be questioned and may turn out to be inefficient. Application of Geodesic grid in calculating surface charges: [222], also mentioning Connolly algorithm (refs 26, 29 therein).

The second scheme has a divergence around  $Nl = r_{\max}$ . It can be shown that the condition  $N \leq \frac{r}{l}$  is enough here to stabilise it.

To account for the molecular geometry and the general tendency of the photo electron to oscillate stronger in the vicinity of the nuclei, the mesh is built out of spheres, centred at the nuclear positions. Thereby, a study of Son [112] had shown that it is numerically most efficient when the overlapping regions of these spheres are cut out. Figure 5.1 shows an example for the water molecule.

For the radial distribution we will use the the function

$$r_i = \frac{1 + x_i}{1 - x_i + \frac{2L}{r_{\max}}} L \quad x_i = \frac{2i}{N_r} - 1$$

suggested by Son *et. al.*[112, 116]. The parameters  $N_r$ ,  $L$  specifying the number of spheres and their distribution; the larger  $L$  is, the denser are the spheres close to the centre.

The optimal choice of these parameters as well as the angular distribution of points on the spheres is still an open question. Finally, after putting the points together as described above, they are connected to a Delaunay triangulation using `tetgen` [217]. Here additional points may be introduced to guarantee well-shaped elements (*i.e.* no sharp peaks).

One scheme suggested by Son and Chu [112]:

$$r_i = \frac{il}{N - i + \frac{lN}{r_{\max}}} \quad i = 1, \dots, N$$

where  $l$  is a parameter to chose. Own scheme:

$$r_i = \frac{il}{\left(\frac{N}{i}\right)^p \left(\frac{Nl}{r_{\max}} - 1\right) + 1} \quad i = 1, \dots, N$$

where  $l$  and  $p \leq 1$  are parameters to chose. Thereby it is important to mention that in this scheme (in contrast to the above one) the (asymptotic for  $r \rightarrow \infty$ ) maximum distance between two spheres is  $l$  and hence could be physically chosen to  $l \approx \frac{\lambda}{2}$ .

The second scheme has a divergence around  $Nl = r_{\max}$ . It can be shown that the condition  $N \leq \frac{r}{l}$  is enough here to stabilise it.

Besides the tricky question about this mapping, also the number of points per sphere depending on the radial distance should be considered. In [112] this is chosen to be constant without further discussion but of course there are several possible design criteria as well. Besides a constant number, also a constant spherical density (hence  $N \propto r^2$ ) are possible; However, it might be sensible to follow a similar idea than in the radial mapping: Being fine close to the nuclei and get coarser with increasing distance. In particular, I followed the design rule of keeping the space between points in radial direction similar to space between points in angular distribution. Hence,  $d_{\text{spheric}} = \sqrt{\frac{4\pi r_i^2}{N_i}} \approx r_i - r_{i-1}$ . For the above described radial schemes, this results in

$$N_i = \frac{4\pi}{\left(1 - \frac{i-1}{i} \frac{\frac{N-i+\frac{lN}{r_{\max}}}{N-i+1+\frac{lN}{r_{\max}}}}{\right)^2}$$

for the first scheme and

$$N_i = \frac{4\pi}{\left(1 - \frac{i-1}{i} \frac{\left(\frac{N}{i}\right)^p \left(\frac{lN}{r_{\max}} - 1\right) + 1}{\left(\frac{N}{i-1}\right)^p \left(\frac{Nl}{r_{\max}} - 1\right) + 1}\right)^2}$$

for the latter.

The respective radii and number of points per sphere are shown in figure 5.2. Special mapping schemes are the constant radial mapping  $r_i = ai$  which corresponds to a constant spherical grid density  $N_i = 4\pi a^2 i^2$  and an exponential map  $r_i = q^i r_0$  which would require a constant number of spherical grid points  $N_i = \frac{4\pi q^2}{(1-q)^2}$  according to the rule derived above.

Since some of the spherical schemes described above allow only for certain numbers of points each, here, the best approximation is used respectively.

### 5.2.2 Obtaining ESP

- obtained from NWChem - interpolation of that grid

## 5.3 Obtaining the DOs

- from coefficients to values




Figure 5.2: the radius and number of points of the spheres as for  $N = 15$ ,  $r_{\max} = 5$  and  $l = 2$ .

# 6 | Results

Showing PES: - benzene - near-threshold PES - Find systems where PES is known for different photon energies

There is a paper[223] with vibronically resolved PES from experiment.  
PES of N<sub>2</sub> with 'angular resolution' (latter not directly) [224].

purine and pyrimidine: comparison to Green's function methods via the [76]-paper.

For AlO<sup>-</sup> there is a paper with 2 spectra with one and two transitions each, having vibrational structure[225].

Atomic Systems have some experimental data as well: For Xe and Kr the spectra at photon energies of 150 eV are shown in reference [226].

A combined theoretical and experimental study on CH<sub>2</sub>F<sub>2</sub> is in ref. [227]. Here, theory is quite bad and experiment is also angular resolved, thus may be interesting.

## 6.1 Benchmarks and Tests on the Formalism

### 6.1.1 Comparison of Formulations

While the unconjugated Burnett-formulation was not very successfull, the conjugated formulations of Burnett and Astley-Leis might both be interesting to compare for a quantum mechanical problem. Since the conjugated Burnett elements lead to infinitely large matrix elements, here the Astley-Leis elements (4.10) will be compared with a symmetrised form where the shape functions are of the form

$$\Phi(\mathbf{r}) = \sqrt{D(r)}\varphi(\mathbf{r})e^{-ik\mu(r)} \quad (6.1)$$

for both, the test- and ansatz functions. Using this formulation, only the kinetic energy expression changes to

$$\begin{aligned} \int (\nabla\Phi_i(\mathbf{r})) (\nabla\Psi_j(\mathbf{r})) d\mathbf{r} &= \int \frac{1}{4} \frac{D'(\mathbf{r})D'(\mathbf{r})}{D(\mathbf{r})} \varphi_i(\mathbf{r})\varphi_j(\mathbf{r}) + \frac{1}{2} D'(\mathbf{r}) (\varphi_i(\mathbf{r})\varphi'_j(\mathbf{r}) + \varphi'_i(\mathbf{r})\varphi_j(\mathbf{r})) \\ &+ D (\varphi'_i(\mathbf{r})\varphi'_j(\mathbf{r}) + ik\mu'(r)(\varphi_i(\mathbf{r})\varphi'_j(\mathbf{r}) - \varphi'_i(\mathbf{r})\varphi_j(\mathbf{r})) - (ik\mu'(r))^2 \varphi_i(\mathbf{r})\varphi_j(\mathbf{r})) d\mathbf{r}. \end{aligned} \quad (6.2)$$

Discussion of spectrum:

- What is the meaning of imaginary eigen-values? How to interprete results?
- Should one sort the 'initial' energies to make it comparable to 'symmetrised'?
- Which conclusions to draw from this study?




Figure 6.2: Decomposition of the first 31 solutions into spherical waves with angular momenta up to  $l = 7$ . Left: Astley-Leis-formulation, right: symmetrised form.

Figure 6.3: Figure to be placed

### Discussion of Spherical Waves:

- In both cases: very strong mixing of quantum numbers
- Is the difference in nature more different than that obtained from slightly differing parameters?
- The symmetrised form indeed seems to damp the  $s$ - and  $p$ -waves out.
- Can one understand these expansions/differences in terms of damping for large  $r$ ?
- Which conclusions can be drawn from this study?

Investigation of the spectrum as a function of the power  $p$  for the symmetric and non-symmetric formulation of the Hamiltonian.




Figure 6.1: The first 50 eigenvalues obtained with the original Astley-Leis formulation (imaginary part not shown) and with the symmetrised form (4.15).

### 6.1.2 Numerical Convergence

Tested on Lithium with a sphere with  $r_{\max} = 2.8$  1: Using the mapping of Son with a naive approach, setting the number of circles by hand and growing quadratically with the size ( $N_{\text{circle}} = 20$ ,  $l = 2.0$ ,  $N = 10$ ) 2: using the Son scheme with  $N$  according to above formula 3: using the tm scheme as described above,  $p=1$  4: using the tm scheme as described above,  $p=2$

scheme	#elements	error in energy [a.u.]
1	2370	0.496220, 0.520199, 0.572279
2	2022	0.038136, 0.046119, 0.055691
3	2166	-0.003638, 0.027219, 0.050588
4	2338	none converged

ber of elements were kept as similar as possible.

In the FEM scheme used to compute the FEF, several parameters governing numerical

In these calculations the num-




Figure 6.5: The eigen energies for different box sizes (coloured dots in the upper pannel) where the box size is denoted by the colour in atomic units. The lower pannel shows the decomposition of the solutions into spherical waves for three box sizes, marked by a star in the color bar respecively.

and physical characteristics are used. The size of the Krylov-subspace does not play a role for the quality of the solution as shown in Figure 6.4 for the symmetric formulation with a power  $p = \frac{1}{2}$  of the damping function  $D$ . Such a good behaviour can, however, only be observed for the symmetric case because in case of the non-hermitian formulation the ordering of the eigenvalues can not be done unambiguously.

More important than the subspace used numerically is obviously the size of the sphere used for the FEM description.

- ‘Convergence’ speed depends on power of  $D$ , chose  $p = \frac{1}{2}$  here.
- Increasing box-size leads to contributions of larger  $l$
- Non-symmetric formulation leads to stronger mixing of angular momenta
- Spectra converged for  $p = 1/8$ , see Figure A.1 in supplement.




Figure 6.4: The energies of the 20 states that fit the target energy best, depending on the size of the Krylov-suspace, indicated by the colour.

## 6.2 atomic Lithium

As the smallest system that has multiple electrons in its anionic state as well, Lithium is an important test object that has theoretical as well as experimental reference data available[66, 67, 228]. Experiment from Moore, cited in [228]; original not found. Experiment 2 from [30].




Figure 6.6: The functional  $J(\alpha, \omega)$  described in eqation (5.11) for Lithium

Cross-section: [229].

The appearance of  $2p$ -states are accessed by a dipole-transition in the bound part and can not be described within the theory applied here due to the neglect of the conjugate DO term in equation (2.14) [63].

## 6.3 Triatomic Linear Molecules

There are several studies on  $\text{CO}_2$  [230–233],  $\text{CS}_2$  [232, 233],  $\text{COS}$  [232, 233] and  $\text{N}_2\text{O}$  [233]

$\text{CS}_2$  is known to have strong correlation effects [72]

## 6.4 water

The PES and threshold PES, where the kinetic energy of the photoelectron is kept constant instead of the energy of the photon, of water is experimentally well studied for different photon energies. As example, in Ref. [234] the PES of water and several TPES of water and heavy water are investigated with high resolution.

Similar high accuracy is achieved in Ref. [235] where the PES of  $\text{H}_2\text{O}^+$  and  $\text{D}_2\text{O}^+$  are studied with a He1 source, radiating at 537 Angs.

Further PES are available at higher photon energies: In [236], photons at energies of 1200 eV are used. A further study of Winter *et. al.* measured spectra at 60, 75 80, 100 and 120 eV in gas and liquid phase [1].

These studies are of interest here since they yield a variety of different test-cases on a single system that can be used to prove the method being used to be valid.




Figure 6.7: The functional  $J(\alpha, \omega)$  described in equation (5.11) for  $\text{CO}_2$

## 7 | Resumee

- is it worth the effort? - What energy-range is achievable? - Which effects are still missing?

- How to proceed with it?

# Bibliography

- [1] B. Winter et al. “Full Valence Band Photoemission from Liquid Water Using EUV Synchrotron Radiation”. In: *The Journal of Physical Chemistry A* 108.14 (Apr. 2004), pp. 2625–2632. DOI: [10.1021/jp030263q](https://doi.org/10.1021/jp030263q).
- [2] Kathrin M. Lange and Emad F. Aziz. “The Hydrogen Bond of Water from the Perspective of Soft X-Ray Spectroscopy”. In: *Chemistry – An Asian Journal* 8.2 (2013), pp. 318–327. DOI: [10.1002/asia.201200533](https://doi.org/10.1002/asia.201200533).
- [3] Sophie E. Canton et al. “Visualizing the non-equilibrium dynamics of photoinduced intramolecular electron transfer with femtosecond X-ray pulses”. In: *Nature Communications* 6.6359 (Mar. 2015). DOI: [10.1038/ncomms7359](https://doi.org/10.1038/ncomms7359).
- [4] Benjamin E. Van Kuiken and Munira Khalil. “Simulating Picosecond Iron K-Edge X-ray Absorption Spectra by ab Initio Methods To Study Photoinduced Changes in the Electronic Structure of Fe(II) Spin Crossover Complexes”. In: *The Journal of Physical Chemistry A* 115.39 (2011). PMID: 21846088, pp. 10749–10761. DOI: [10.1021/jp2056333](https://doi.org/10.1021/jp2056333). eprint: <http://dx.doi.org/10.1021/jp2056333>.
- [5] Xin Yang et al. “Solvation of the Azide Anion (N<sub>3</sub><sup>-</sup>) in Water Clusters and Aqueous Interfaces: A Combined Investigation by Photoelectron Spectroscopy, Density Functional Calculations, and Molecular Dynamics Simulations”. In: *The Journal of Physical Chemistry A* 108.39 (2004), pp. 7820–7826. DOI: [10.1021/jp0496396](https://doi.org/10.1021/jp0496396). eprint: <http://dx.doi.org/10.1021/jp0496396>.
- [6] Babak Minofar et al. “Interior and Interfacial Aqueous Solvation of Benzene Di-carboxylate Dianions and Their Methylated Analogues: A Combined Molecular Dynamics and Photoelectron Spectroscopy Study”. In: *The Journal of Physical Chemistry A* 109.23 (2005), pp. 5042–5049. DOI: [10.1021/jp050836u](https://doi.org/10.1021/jp050836u). eprint: <http://dx.doi.org/10.1021/jp050836u>.
- [7] Stephan Thürmer et al. “Flexible H<sub>2</sub>O<sub>2</sub> in Water: Electronic Structure from Photoelectron Spectroscopy and Ab Initio Calculations”. In: *The Journal of Physical Chemistry A* 115.23 (2011). PMID: 21332235, pp. 6239–6249. DOI: [10.1021/jp111674s](https://doi.org/10.1021/jp111674s). eprint: <http://dx.doi.org/10.1021/jp111674s>.
- [8] Mihajlo Etinski and Christel M. Marian. “Ab initio investigation of the methylation and hydration effects on the electronic spectra of uracil and thymine”. In: *Phys. Chem. Chem. Phys.* 12 (19 2010), pp. 4915–4923. DOI: [10.1039/B925677F](https://doi.org/10.1039/B925677F).
- [9] Franck Lépine, Giuseppe Sansone, and Marc J.J. Vrakking. “Molecular applications of attosecond laser pulses”. In: *Chem. Phys. Lett.* 578 (July 18, 2013), pp. 1–14.

- [10] Eleftherios Goulielmakis et al. “Real time observation of valence electron motion”. In: *Nature* 466 (Aug. 5, 2010), pp. 739–744. DOI: [10.1038/nature09212](https://doi.org/10.1038/nature09212).
- [11] Dr. Giuseppe Sansone et al. “Electron Correlation in Real Time”. In: *Chem. Phys. Chem.* 13.3 (2012), pp. 661–680. DOI: [10.1002/cphc.201100528](https://doi.org/10.1002/cphc.201100528).
- [12] Matthias F. Kling and Marc J.J. Vrakking. “Attosecond Electron Dynamics”. In: *Annual Review of Physical Chemistry* 59.1 (May 2008), pp. 463–492. DOI: [10.1146/annurev.physchem.59.032607.093532](https://doi.org/10.1146/annurev.physchem.59.032607.093532).
- [13] M. Uiberacker et al. “Attosecond real-time observation of electron tunnelling in atoms”. In: *Nature* 446.7136 (Apr. 5, 2007), pp. 627–632. DOI: [10.1038/nature05648](https://doi.org/10.1038/nature05648).
- [14] Ferenc Krausz and Misha Ivanov. “Attosecond physics”. In: *Reviews of Modern Physics* 81.1 (Feb. 2, 2009), pp. 163–234. DOI: [10.1103/RevModPhys.81.163](https://doi.org/10.1103/RevModPhys.81.163).
- [15] T Koopmans. “Über die Zuordnung von Wellenfunktionen und Eigenwerten zu den Einzelnen Elektronen Eines Atoms”. In: *Physica* 1.1 (1934), pp. 104–113. DOI: [http://dx.doi.org/10.1016/S0031-8914\(34\)90011-2](http://dx.doi.org/10.1016/S0031-8914(34)90011-2).
- [16] C.-O. Almbladh and U. von Barth. “Exact results for the charge and spin densities, exchange-correlation potentials, and density-functional eigenvalues”. In: *Phys. Rev. B* 31 (6 Mar. 1985), pp. 3231–3244. DOI: [10.1103/PhysRevB.31.3231](https://doi.org/10.1103/PhysRevB.31.3231).
- [17] David A. Egger et al. “Outer-valence Electron Spectra of Prototypical Aromatic Heterocycles from an Optimally Tuned Range-Separated Hybrid Functional”. In: *JCTC* 10.5 (2014), pp. 1934–1952. DOI: [10.1021/ct400956h](https://doi.org/10.1021/ct400956h).
- [18] Thomas Körzdörfer et al. “Assessment of the performance of tuned range-separated hybrid density functionals in predicting accurate quasiparticle spectra”. In: *Phys. Rev. B* 86 (20 Nov. 2012), p. 205110. DOI: [10.1103/PhysRevB.86.205110](https://doi.org/10.1103/PhysRevB.86.205110).
- [19] T. Körzdörfer et al. “When to trust photoelectron spectra from Kohn-Sham eigenvalues: The case of organic semiconductors”. In: *Phys. Rev. B* 79 (20 May 2009), p. 201205. DOI: [10.1103/PhysRevB.79.201205](https://doi.org/10.1103/PhysRevB.79.201205).
- [20] C-Z Gao et al. “On the dynamics of photo-electrons in C 60”. In: *J Phys. B-A. Mol. Opt.* 48.10 (2015), p. 105102.
- [21] Yu. N. Zhuravlev and D. V. Korabelnikov. “The nature of the electronic states and photoelectron spectra of oxyanion crystals”. In: *Journal of Structural Chemistry* 50.6 (2009), pp. 1021–1028. DOI: [10.1007/s10947-009-0151-7](https://doi.org/10.1007/s10947-009-0151-7).
- [22] R. Leckey. “Ultraviolet Photoelectron Spectroscopy of Solids”. In: *Surface Analysis Methods in Materials Science*. Ed. by D. John O’Connor, Brett A. Sexton, and Roger St. C. Smart. Berlin, Heidelberg: Springer Berlin Heidelberg, 1992, pp. 291–300. DOI: [10.1007/978-3-662-02767-7\\_14](https://doi.org/10.1007/978-3-662-02767-7_14).
- [23] J Schirmer et al. “Break-down of the molecular-orbital picture of ionization: CS, PN and P 2”. In: *Journal of Physics B: Atomic and Molecular Physics* 11.11 (1978), p. 1901.
- [24] J. Schirmer et al. “Strong Correlation Effects in inner Valence Ionization of N<sub>2</sub> AND CO”. In: *Chemical Physics* 26.1 (1977), pp. 149–153. DOI: [http://dx.doi.org/10.1016/0301-0104\(77\)87101-2](http://dx.doi.org/10.1016/0301-0104(77)87101-2).

- [25] M. Brosolo and P. Decleva. “Variational approach to continuum orbitals in a spline basis: An application to H<sub>2</sub>+ photoionization”. In: *Chemical Physics* 159.2 (Jan. 1992), pp. 185–196. DOI: 10.1016/0301-0104(92)80069-8.
- [26] A. Abdurrouf and F. H. M. Faisal. “Theory of intense-field dynamic alignment and high-order harmonic generation from coherently rotating molecules and interpretation of intense-field ultrafast pump-probe experiments”. In: *Phys. Rev. A* 79 (2 Feb. 2009), p. 023405. DOI: 10.1103/PhysRevA.79.023405.
- [27] Peng Zhang and A. G. R. Thomas. “Enhancement of high-order harmonic generation in intense laser interactions with solid density plasma by multiple reflections and harmonic amplification”. In: *Applied Physics Letters* 106.13 (Mar. 30, 2015), p. 131102. DOI: 10.1063/1.4916739.
- [28] B. Dromey et al. “High harmonic generation in the relativistic limit”. In: *Nature Physics* 2.7 (July 2006), pp. 456–459. DOI: 10.1038/nphys338.
- [29] Enrico Fermi. *Nuclear Physics*. Chicago: University of Chicago Press, 1949.
- [30] B. Sonntag and P. Zimmermann. “XUV spectroscopy of metal atoms”. In: *Reports on Progress in Physics* 55.7 (1992), p. 911.
- [31] Gilbert Grell. “Theoretical Study of X-Ray Photoelectron Spectroscopy of Transition Metal Complexes”. University of Rostock, 2015.
- [32] L. D. Landau and E.M. Lifschitz. *Quantenmechanik*. 9th ed. Berlin: Akademie-Verlag, 1979.
- [33] C. Melania Oana and Anna I. Krylov. “Cross sections and photoelectron angular distributions in photodetachment from negative ions using equation-of-motion coupled-cluster Dyson orbitals”. In: *J. Chem. Phys.* 131.12, 124114 (2009). DOI: <http://dx.doi.org/10.1063/1.3231143>.
- [34] Yuan Liu and Chuangang Ning. “Calculation of photodetachment cross sections and photoelectron angular distributions of negative ions using density functional theory”. In: *The Journal of Chemical Physics* 143.14 (Oct. 14, 2015), p. 144310. DOI: 10.1063/1.4932978.
- [35] Dietrich Braess. *Finite Elemente*. 4th ed. Heidelberg: Springer, 2007.
- [36] David Gilbarg. *Elliptic Partial Differential Equations of Second Order*. Berlin: 1983, 1983.
- [37] Philippe G. Ciarlet. *The Finite Element Method for Elliptic Problems*. Amsterdam et al.: SIAM, 2002.
- [38] R. J. Astley. “Wave envelope and infinite elements for acoustical radiation”. In: *International Journal for Numerical Methods in Fluids* 3.5 (1983), pp. 507–526.
- [39] R. J. Astley et al. “Three-dimensional wave-envelope elements of variable order for acoustic radiation and scattering. Part I. Formulation in the frequency domain”. In: *The Journal of the Acoustical Society of America* 103.1 (1998), pp. 49–63.
- [40] R. J. Astley. “Infinite elements for wave problems: a review of current formulations and an assessment of accuracy”. In: *International Journal for Numerical Methods in Engineering* 49.7 (2000), pp. 951–976.

- [41] Daniel Dreyer. “Efficient Infinite Elements for Exterior Acoustics”. PhD thesis. Germany: TU Hamburg-Harburg, 2004.
- [42] A. Sommerfeld. *Partial differential equation in physics*. 1949.
- [43] P. Hohenberg and W. Kohn. “Inhomogeneous Electron Gas”. In: *Phys. Rev.* 136 (3B Nov. 1964), B864–B871. DOI: [10.1103/PhysRev.136.B864](https://doi.org/10.1103/PhysRev.136.B864).
- [44] W. Kohn and L. J. Sham. “Self-Consistent Equations Including Exchange and Correlation Effects”. In: *Phys. Rev.* 140 (4A Nov. 1965), A1133–A1138. DOI: [10.1103/PhysRev.140.A1133](https://doi.org/10.1103/PhysRev.140.A1133).
- [45] Olga S. Bokareva et al. “Tuning Range-Separated Density Functional Theory for Photocatalytic Water Splitting Systems”. In: *Journal of Chemical Theory and Computation* 11.4 (Apr. 14, 2015), pp. 1700–1709. DOI: [10.1021/acs.jctc.5b00068](https://doi.org/10.1021/acs.jctc.5b00068).
- [46] Gilbert Grell et al. “Multi-reference approach to the calculation of photoelectron spectra including spin-orbit coupling”. In: *J. Chem. Phys.* 143.7, 074104 (2015). DOI: <http://dx.doi.org/10.1063/1.4928511>.
- [47] Iann C. Gerber and János G. Ángyán. “Potential curves for alkaline-earth dimers by density functional theory with long-range correlation corrections”. In: *Chemical Physics Letters* 416.4–6 (2005), pp. 370–375. DOI: <http://dx.doi.org/10.1016/j.cplett.2005.09.059>.
- [48] Iann C. Gerber and János G. Ángyán. “Hybrid functional with separated range”. In: *Chemical Physics Letters* 415.1–3 (2005), pp. 100–105. DOI: <http://dx.doi.org/10.1016/j.cplett.2005.08.060>.
- [49] Saumitra Saha et al. “Coexistence of 1,3-butadiene conformers in ionization energies and Dyson orbitals”. In: *The Journal of Chemical Physics* 123.12, 124315 (2005). DOI: <http://dx.doi.org/10.1063/1.2034467>.
- [50] Alejandro Soba et al. “Real-space density functional theory and time dependent density functional theory using finite/infinite element methods”. In: *Computer Physics Communications* 183.12 (Dec. 2012), pp. 2581–2588. DOI: [10.1016/j.cpc.2012.07.007](https://doi.org/10.1016/j.cpc.2012.07.007).
- [51] Peter Bettess and Richard A. Abram. “Finite and infinite elements for a simple problem in quantum mechanics”. In: *Communications in numerical methods in engineering* 18.5 (2002), pp. 325–334.
- [52] M. Valiev et al. “NWChem: A comprehensive and scalable open-source solution for large scale molecular simulations”. In: *Comput. Phys. Commun.* 181.9 (2010), pp. 1477–1489. DOI: <http://dx.doi.org/10.1016/j.cpc.2010.04.018>.
- [53] M. J. Frisch et al. *Gaussian 09 Revision E.01*. Gaussian Inc. Wallingford CT 2009.
- [54] B. S. Kirk et al. “libMesh: A C++ Library for Parallel Adaptive Mesh Refinement/Coarsening Simulations”. In: *Engineering with Computers* 22.3–4 (2006). <http://dx.doi.org/10.1007/s00366-006-0049-3>, pp. 237–254.
- [55] Vicente Hernandez, Jose E. Roman, and Vicente Vidal. “SLEPc: A scalable and flexible toolkit for the solution of eigenvalue problems”. In: *ACM Trans. Math. Software* 31.3 (2005), pp. 351–362.

- [56] V. Hernandez, J. E. Roman, and V. Vidal. “SLEPc: Scalable Library for Eigenvalue Problem Computations”. In: *Lect. Notes Comput. Sci.* 2565 (2003), pp. 377–391.
- [57] Satish Balay et al. *PETSc Web page*. <http://www.mcs.anl.gov/petsc>. 2016.
- [58] Stephan Hüfner. *Photoelectron Spectroscopy*. 3rd ed. Berlin: Springer Verlag, 2003.
- [59] C.J. Milne, T.J. Penfold, and M. Chergui. “Recent experimental and theoretical developments in time-resolved X-ray spectroscopies”. In: *Coordination Chemistry Reviews* 277-278 (Oct. 2014), pp. 44–68. DOI: [10.1016/j.ccr.2014.02.013](https://doi.org/10.1016/j.ccr.2014.02.013).
- [60] O. V. Gritsenko and E. J. Baerends. “Time-dependent Dyson orbital theory”. In: *Phys. Chem. Chem. Phys.* (2016). DOI: [10.1039/C6CP00888G](https://doi.org/10.1039/C6CP00888G).
- [61] S. Bauch, L. K. Sørensen, and L. B. Madsen. “Time-dependent generalized-active-space configuration-interaction approach to photoionization dynamics of atoms and molecules”. In: *Phys. Rev. A* 90.6 (Dec. 9, 2014). DOI: [10.1103/PhysRevA.90.062508](https://doi.org/10.1103/PhysRevA.90.062508).
- [62] H. R. Larsson et al. “Correlation effects in strong-field ionization of heteronuclear diatomic molecules”. In: *Phys. Rev. A* 93 (1 Jan. 2016), p. 013426. DOI: [10.1103/PhysRevA.93.013426](https://doi.org/10.1103/PhysRevA.93.013426).
- [63] Aurora Ponzi et al. “Dynamical photoionization observables of the CS molecule: The role of electron correlation”. In: *The Journal of Chemical Physics* 140.20, 204304 (2014). DOI: <http://dx.doi.org/10.1063/1.4876495>.
- [64] Zhigang Sun, Nanquan Lou, and Gunnar Nyman. “Time-Dependent Wave Packet Split Operator Calculations on a Three-Dimensional Fourier Grid in Radau Coordinates Applied to the OCLO Photoelectron Spectrum”. In: *The Journal of Physical Chemistry A* 108.42 (2004), pp. 9226–9232. DOI: [10.1021/jp0477203](https://doi.org/10.1021/jp0477203). eprint: <http://dx.doi.org/10.1021/jp0477203>.
- [65] Thomas-C. Jagau and Anna I. Krylov. “Complex Absorbing Potential Equation-of-Motion Coupled-Cluster Method Yields Smooth and Internally Consistent Potential Energy Surfaces and Lifetimes for Molecular Resonances”. In: *JPC* 5.17 (Sept. 4, 2014), pp. 3078–3085. DOI: [10.1021/jz501515j](https://doi.org/10.1021/jz501515j).
- [66] L. Journal et al. “First experimental determination and theoretical calculation of partial photoionization cross sections of lithium over the energy region of hollow Atomic States”. In: *Physical review letters* 76.1 (1996), p. 30.
- [67] S. Diehl et al. “Angle-Resolved Photoelectron Spectrometry Studies of the Autoionization of the 2 s 2 2 p 2 P Triply Excited State of Atomic Lithium: Experimental Results and R-Matrix Calculations”. In: *Physical review letters* 84.8 (2000), p. 1677.
- [68] A.G. Sunderland et al. “A parallel R-matrix program PRMAT for electron–atom and electron–ion scattering calculations”. In: *Comput. Phys. Commun.* 145.3 (2002), pp. 311–340. DOI: [http://dx.doi.org/10.1016/S0010-4655\(02\)00140-6](http://dx.doi.org/10.1016/S0010-4655(02)00140-6).
- [69] Yu Yan and M. J. Seaton. “R-matrix theory of the hydrogen atom”. In: *Journal of Physics B: Atomic and Molecular Physics* 18.13 (1985), p. 2577.
- [70] B.D. Buckley, P.G. Burke, and Vo Ky Lan. “R-matrix calculations for electron–molecule scattering”. In: *Computer Physics Communications* 17.1 (1979), pp. 175–179. DOI: [http://dx.doi.org/10.1016/0010-4655\(79\)90080-8](http://dx.doi.org/10.1016/0010-4655(79)90080-8).

- [71] Craig A. Bayse and Kristine N. Ortwine. “Calculation of Photoelectron Spectra of Molybdenum and Tungsten Complexes Using Green’s Functions Methods”. In: *The Journal of Physical Chemistry A* 111.32 (2007). PMID: 17636964, pp. 7841–7847. DOI: 10.1021/jp063029j. eprint: <http://dx.doi.org/10.1021/jp063029j>.
- [72] J. Schirmer et al. “Strong correlation effects in the ionisation of CS<sub>2</sub>”. In: *Chem. Phys. Lett.* 61.1 (1979), pp. 30–35. DOI: [http://dx.doi.org/10.1016/0009-2614\(79\)85078-2](http://dx.doi.org/10.1016/0009-2614(79)85078-2).
- [73] W.G. Richards. “The use of Koopmans’ Theorem in the interpretation of photoelectron spectra”. In: *International Journal of Mass Spectrometry and Ion Physics* 2.6 (1969), pp. 419–424. DOI: [http://dx.doi.org/10.1016/0020-7381\(69\)80040-9](http://dx.doi.org/10.1016/0020-7381(69)80040-9).
- [74] Anthony R. Schwager, Kevin H. Bell, and Ellak I. von Nagy-Felsobuki. “PHOTO-ELECTRON SPECTRA OF THE ALKYL PHENYL DISULFIDES”. In: *Phosphorus, Sulfur, and Silicon and the Related Elements* 70.1 (1992), pp. 287–295. DOI: 10.1080/10426509208049177. eprint: <http://dx.doi.org/10.1080/10426509208049177>.
- [75] K.C. Prince et al. “Study of complex molecules of biological interest with synchrotron radiation”. In: *Journal of Electron Spectroscopy and Related Phenomena* 204, Part B (2015). Gas phase spectroscopic and dynamical studies at Free-Electron Lasers and other short wavelength sources, pp. 335–344. DOI: <http://dx.doi.org/10.1016/j.elspec.2015.08.010>.
- [76] A W Potts et al. “An experimental and theoretical study of the valence shell photoelectron spectra of purine and pyrimidine molecules”. In: *J Phys. B-A. Mol. Opt.* 36.14 (2003), p. 3129.
- [77] David A. Egger et al. “Outer-valence Electron Spectra of Prototypical Aromatic Heterocycles from an Optimally Tuned Range-Separated Hybrid Functional”. In: *JCTC* 10.5 (2014). PMID: 24839410, pp. 1934–1952. DOI: 10.1021/ct400956h.
- [78] Diana Yepes et al. “Photoemission Spectra and Density Functional Theory Calculations of 3d Transition Metal–Aqua Complexes (Ti–Cu) in Aqueous Solution”. In: *The Journal of Physical Chemistry B* 118.24 (2014). PMID: 24902000, pp. 6850–6863. DOI: 10.1021/jp5012389.
- [79] Leo P. Kadanoff and Gordon Baym. *Quantum Statistical Mechanics: Green’s Function Method in Equilibrium and Nonequilibrium Problems*. Addison-Wesley Publishing Co. Inc., 1989.
- [80] W. Von Niessen, J. Schirmer, and L. S. Cederbaum. “Computational methods for the one-particle green’s function”. In: *Computer Physics Reports* 1.2 (1984), pp. 57–125.
- [81] J. Schirmer, L. S. Cederbaum, and O. Walter. “New approach to the one-particle Green’s function for finite Fermi systems”. In: *Phys. Rev. A* 28 (3 Sept. 1983), pp. 1237–1259. DOI: 10.1103/PhysRevA.28.1237.
- [82] M J Seaton. “Use of the R matrix method for bound-state calculations. I. General theory”. In: *Journal of Physics B: Atomic and Molecular Physics* 18.11 (1985), p. 2111.

- [83] L. M. Kiernan et al. “Resonant photoionization of atomic lithium in the region of the first and second inner-shell thresholds”. In: *J Phys. B-A. Mol. Opt.* 29.6 (1996), p. L181.
- [84] B M McLaughlin and R C Forrey. “Low energy electron collisions with the N 2 H radical”. In: *Journal of Physics: Conference Series* 388.5 (2012), p. 052027.
- [85] Keith H. Johnson. “Scattered-Wave Theory of the Chemical Bond\*”. In: Advances in Quantum Chemistry 7 (1973). Ed. by Per-Olov Löwdin, pp. 143–185. DOI: [http://dx.doi.org/10.1016/S0065-3276\(08\)60561-4](http://dx.doi.org/10.1016/S0065-3276(08)60561-4).
- [86] J. C. Slater and K. H. Johnson. “Self-Consistent-Field  $X\alpha$  Cluster Method for Polyatomic Molecules and Solids”. In: *Phys. Rev. B* 5 (3 Feb. 1972), pp. 844–853. DOI: [10.1103/PhysRevB.5.844](https://doi.org/10.1103/PhysRevB.5.844).
- [87] F Beleznay and M J Lawrence. “The ‘muffin-tin’ approximation in the calculation of electronic band structure”. In: *Journal of Physics C: Solid State Physics* 1.5 (1968), p. 1288.
- [88] D. D. Koelling, A. J. Freeman, and F. M. Mueller. “Shifts in the Electronic Band Structure of Metals Due to Non-Muffin-Tin Potentials”. In: *Phys. Rev. B* 1 (4 Feb. 1970), pp. 1318–1324. DOI: [10.1103/PhysRevB.1.1318](https://doi.org/10.1103/PhysRevB.1.1318).
- [89] M. Venut, M. Stener, and P. Decleva. “Valence photoionization of C<sub>6</sub>H<sub>6</sub> by the B-spline one-centre expansion density functional method”. In: *Chemical Physics* 234.1 (1998), pp. 95–109. DOI: [http://dx.doi.org/10.1016/S0301-0104\(98\)00179-7](http://dx.doi.org/10.1016/S0301-0104(98)00179-7).
- [90] J. A. Richards and F. P. Larkins. “Molecular photoionisation calculations with numerical continuum wavefunctions: application to the hydrogen molecule”. In: *Journal of Physics B: Atomic and Molecular Physics* 17.6 (1984), p. 1015.
- [91] Liang Tao, C. W. McCurdy, and T. N. Rescigno. “Grid-based methods for diatomic quantum scattering problems: A finite-element discrete-variable representation in prolate spheroidal coordinates”. In: *Phys. Rev. A* 79 (1 Jan. 2009), p. 012719. DOI: [10.1103/PhysRevA.79.012719](https://doi.org/10.1103/PhysRevA.79.012719).
- [92] Gireesh M. Krishnan, Subhas Ghosal, and Susanta Mahapatra. “Theoretical Study of the Electronic Nonadiabatic Transitions in the Photoelectron Spectroscopy of F<sub>2</sub>O”. In: *The Journal of Physical Chemistry A* 110.3 (2006). PMID: 16420003, pp. 1022–1030. DOI: [10.1021/jp0543882](https://doi.org/10.1021/jp0543882). eprint: <http://dx.doi.org/10.1021/jp0543882>.
- [93] Russell T Pack and Gregory A. Parker. “Quantum reactive scattering in three dimensions using hyperspherical (APH) coordinates. Theory”. In: *The Journal of Chemical Physics* 87.7 (1987), pp. 3888–3921. DOI: <http://dx.doi.org/10.1063/1.452944>.
- [94] F. L. Yip, C. W. McCurdy, and T. N. Rescigno. “Hybrid orbital and numerical grid representation for electronic continuum processes: Double photoionization of atomic beryllium”. In: *Phys. Rev. A* 81 (5 May 2010), p. 053407. DOI: [10.1103/PhysRevA.81.053407](https://doi.org/10.1103/PhysRevA.81.053407).

- [95] Matthias Ruckenbauer, Sebastian Mai, and Leticia Marquetand Philipp and González. “Revealing Deactivation Pathways Hidden in Time-Resolved Photoelectron Spectra”. In: *Scientific Reports* (2016), p. 35522. DOI: 10.1038/srep35522.
- [96] Teijo Åberg. “Theory of X-Ray Satellites”. In: *Phys. Rev.* 156 (1 Apr. 1967), pp. 35–41. DOI: 10.1103/PhysRev.156.35.
- [97] Eric L. Michelsen. *Quirky Quantum Concepts*. Springer, 2014. DOI: 10.1007/978-1-4614-9305-1.
- [98] K. Gokhberg et al. “Molecular photoionization cross sections by Stieltjes–Chebyshev moment theory applied to Lanczos pseudospectra”. In: *J. Chem. Phys.* 130.6, 064104 (2009). DOI: <http://dx.doi.org/10.1063/1.3073821>.
- [99] Samer Gozem et al. “Photoelectron Wave Function in Photoionization: Plane Wave or Coulomb Wave?” In: *The Journal of Physical Chemistry Letters* 6.22 (2015). PMID: 26509428, pp. 4532–4540. DOI: 10.1021/acs.jpclett.5b01891. eprint: <http://dx.doi.org/10.1021/acs.jpclett.5b01891>.
- [100] L. C. Green, S. Matsushima, and E. K. Kolchin. “Tables of the Continuum Wave Functions for Hydrogen.” In: 3 (Nov. 1958), p. 459. DOI: 10.1086/190041.
- [101] Alexander Humeniuk et al. “Time-resolved photoelectron imaging spectra from non-adiabatic molecular dynamics simulations”. In: *The Journal of Chemical Physics* 139.13, 134104 (2013). DOI: <http://dx.doi.org/10.1063/1.4820238>.
- [102] Gaia Tomasello, Alexander Humeniuk, and Roland Mitrić. “Exploring Ultrafast Dynamics of Pyrazine by Time-Resolved Photoelectron Imaging”. In: *The Journal of Physical Chemistry A* 118.37 (2014). PMID: 24927218, pp. 8437–8445. DOI: 10.1021/jp5017328. eprint: <http://dx.doi.org/10.1021/jp5017328>.
- [103] Benoît Mignolet et al. “Probing Rapidly-Ionizing Super-Atom Molecular Orbitals in C<sub>60</sub>: A Computational and Femtosecond Photoelectron Spectroscopy Study”. In: *ChemPhysChem* 14.14 (2013), pp. 3332–3340. DOI: 10.1002/cphc.201300585.
- [104] P.W. Langhoff. “Bounds for second-order optical properties from quantum-mechanical sum rules and the theory of moments”. In: *Chemical Physics Letters* 9.2 (1971), pp. 89–94. DOI: [http://dx.doi.org/10.1016/0009-2614\(71\)80193-8](http://dx.doi.org/10.1016/0009-2614(71)80193-8).
- [105] Janusz Cukras et al. “Photoionization cross section by Stieltjes imaging applied to coupled cluster Lanczos pseudo-spectra”. In: *J. Chem. Phys.* 139.9, 094103 (2013). DOI: <http://dx.doi.org/10.1063/1.4819126>.
- [106] P.W. Langhoff. “Stieltjes imaging of atomic and molecular photoabsorption profiles”. In: *Chem. Phys. Lett.* 22.1 (1973), pp. 60–64. DOI: [http://dx.doi.org/10.1016/0009-2614\(73\)80534-2](http://dx.doi.org/10.1016/0009-2614(73)80534-2).
- [107] P. W. Langhoff et al. “Moment-theory investigations of photoabsorption and dispersion profiles in atoms and ions”. In: *Phys. Rev. A* 14 (3 Sept. 1976), pp. 1042–1056. DOI: 10.1103/PhysRevA.14.1042.
- [108] M. Stener, P. Decleva, and A. Lisini. “Molecular photoionization cross sections by the local density LCAO Stieltjes imaging approach”. In: *Journal of Electron Spectroscopy and Related Phenomena* 74.1 (1995), pp. 29–43. DOI: [http://dx.doi.org/10.1016/0368-2048\(95\)02357-7](http://dx.doi.org/10.1016/0368-2048(95)02357-7).

- [109] Liang Gong et al. “Numerical analysis on quantum dots-in-a-well structures by finite difference method”. In: *Superlattices and Microstructures* 60 (2013), pp. 311–319. DOI: <http://dx.doi.org/10.1016/j.spmi.2013.05.012>.
- [110] Liang Gong et al. “Numerical computation of pyramidal quantum dots with band non-parabolicity”. In: *Superlattices and Microstructures* 61 (2013), pp. 81–90. DOI: <http://dx.doi.org/10.1016/j.spmi.2013.06.011>.
- [111] I.K. Gainullin and M.A. Sonkin. “High-performance parallel solver for 3D time-dependent Schrodinger equation for large-scale nanosystems”. In: *Computer Physics Communications* 188 (2015), pp. 68–75. DOI: <http://dx.doi.org/10.1016/j.cpc.2014.11.005>.
- [112] Sang-Kil Son. “Voronoi-cell finite difference method for accurate electronic structure calculation of polyatomic molecules on unstructured grids”. In: *Journal of Computational Physics* 230.5 (2011), pp. 2160–2173. DOI: <http://dx.doi.org/10.1016/j.jcp.2010.12.012>.
- [113] Franz Aurenhammer. “Voronoi Diagrams& Mdash;a Survey of a Fundamental Geometric Data Structure”. In: *ACM Comput. Surv.* 23.3 (Sept. 1991), pp. 345–405. DOI: [10.1145/116873.116880](https://doi.org/10.1145/116873.116880).
- [114] N. Sukumar. “Voronoi cell finite difference method for the diffusion operator on arbitrary unstructured grids”. In: *International Journal for Numerical Methods in Engineering* 57.1 (2003), pp. 1–34.
- [115] Qiang Du and Max Gunzburger. “Grid generation and optimization based on centroidal Voronoi tessellations”. In: *Applied Mathematics and Computation* 133.2–3 (2002), pp. 591–607. DOI: [http://dx.doi.org/10.1016/S0096-3003\(01\)00260-0](http://dx.doi.org/10.1016/S0096-3003(01)00260-0).
- [116] Sang-Kil Son and Shih-I Chu. “Theoretical study of orientation-dependent multi-photon ionization of polyatomic molecules in intense ultrashort laser fields: A new time-dependent Voronoi-cell finite difference method”. In: *Chem. Phys.* 366.1–3 (2009), pp. 91–102. DOI: <http://dx.doi.org/10.1016/j.chemphys.2009.09.006>.
- [117] B. Shizgal. *Spectral Methods in Chemistry and Physics. Applications to Kinetic Theory and Quantum Mechanics*. 2nd ed. Springer Verlag, 2015.
- [118] Joseph B. Fourier. *Theorie Analytique de la Chaleur*. Firmin Didot, 1822.
- [119] Bengt Fornberg. *A Practical Guide to Pseudospectral Methods*. Ed. by P. G. Ciarlet et al. Vol. 1. Cambridge University Press, 1996.
- [120] David J. Tannor. *Introduction to Quantum Mechanics. A Time-dependent Perspective*. Sausalito, California: University Science Book, 2007.
- [121] Dequan Yu, Shu-Lin Cong, and Zhigang Sun. “An improved Lobatto discrete variable representation by a phase optimisation and variable mapping method”. In: *Chem. Phys.* 458 (2015), pp. 41–51. DOI: <http://dx.doi.org/10.1016/j.chemphys.2015.07.009>.

- [122] Liang-You Peng and Anthony F. Starace. “Application of Coulomb wave function discrete variable representation to atomic systems in strong laser fields”. In: *JCP* 125.15, 154311 (2006).
- [123] A. R. Edmonds. *Angular Momentum in Quantum Mechanics*. 4th ed. Princeton, New Jersey: Princeton University Press, 1996.
- [124] Akbar Mohebbi, Mostafa Abbaszadeh, and Mehdi Dehghan. “The use of a meshless technique based on collocation and radial basis functions for solving the time fractional nonlinear Schrödinger equation arising in quantum mechanics”. In: *Engineering Analysis with Boundary Elements* 37.2 (2013), pp. 475–485. DOI: <http://dx.doi.org/10.1016/j.enganabound.2012.12.002>.
- [125] Lihua Wang, Fuyun Chu, and Zheng Zhong. “Study of radial basis collocation method for wave propagation”. In: *Engineering Analysis with Boundary Elements* 37.2 (2013), pp. 453–463. DOI: <http://dx.doi.org/10.1016/j.enganabound.2012.12.001>.
- [126] Richard Franke. “Scattered data interpolation: tests of some methods”. In: *Math. Comp.* 38.157 (1982), pp. 181–200. DOI: 10.2307/2007474.
- [127] J. C. Carr, W. R. Fright, and R. K. Beatson. “Surface interpolation with radial basis functions for medical imaging”. In: *IEEE Transactions on Medical Imaging* 16.1 (Feb. 1997), pp. 96–107. DOI: 10.1109/42.552059.
- [128] P. F. Batcho. “Computational method for general multicenter electronic structure calculations”. In: *Phys. Rev. E* 61 (6 June 2000), pp. 7169–7183. DOI: 10.1103/PhysRevE.61.7169.
- [129] S. Petersen, D. Dreyer, and O. von Estorff. “Practical considerations of the acoustic FEM for higher frequencies.” In: *30. Tagung der Arbeitsgemeinschaft für Akustik DAGA 2004*. mubve. Straßburg, 2004, pp. 307–308.
- [130] David A. Mazzotti. “Spectral difference methods for solving differential equations”. In: *Chem. Phys. Lett.* 299.5 (1999), pp. 473–480. DOI: [http://dx.doi.org/10.1016/S0009-2614\(98\)01324-4](http://dx.doi.org/10.1016/S0009-2614(98)01324-4).
- [131] C. Liang, R. Kannan, and Z.J. Wang. “A p-multigrid spectral difference method with explicit and implicit smoothers on unstructured triangular grids”. In: *Computers & Fluids* 38.2 (2009), pp. 254–265. DOI: <http://dx.doi.org/10.1016/j.compfluid.2008.02.004>.
- [132] Yen Liu, Marcel Vinokur, and Z.J. Wang. “Spectral difference method for unstructured grids I: Basic formulation”. In: *Journal of Computational Physics* 216.2 (2006), pp. 780–801. DOI: <http://dx.doi.org/10.1016/j.jcp.2006.01.024>.
- [133] Michael W. Frazier. *An Introduction to Wavelets Through Linear Algebra*. New York: Springer, 1999.
- [134] Stephan Dahlke. “Wavelets: Construction Principles and Applications to the Numerical Treatment of Operator Equations”. PhD thesis. Germany: RWTH Aachen, 1996.

- [135] T. Torsti et al. “Three real-space discretization techniques in electronic structure calculation”. In: *Physica Status Solidi (B)* 243.5 (2006), pp. 1016–1053. DOI: 10.1002/pssb.200541348.
- [136] *Information technology – JPEG 2000 image coding system: Conformance testing*. Standard. International Organization for Standardization, Dec. 2004.
- [137] Gerlind Plonka and Manfred Tasche. “On the Computation of Periodic Spline Wavelets”. In: *Appl. Comput. Harmonic Anal.* 2 2.1 (1995), pp. 1–14. DOI: <http://dx.doi.org/10.1006/acha.1995.1001>.
- [138] “6. Orthonormal Bases of Compactly Supported Wavelets”. In: *Ten Lectures on Wavelets*, pp. 167–213. DOI: 10.1137/1.9781611970104.ch6. eprint: <http://epubs.siam.org/doi/pdf/10.1137/1.9781611970104.ch6>.
- [139] *Wavelet Applications in Chemical Engineering*. 1st ed. Springer US, 1994. DOI: 10.1007/978-1-4615-2708-4.
- [140] Angela Kunoth. *Wavelet Methods- Ellipting Boundary Value Problems and Control Problems*. 1st ed. Teubner, 2001.
- [141] Marcílio N Guimarães and Frederico V Prudente. “A study of the confined hydrogen atom using the finite element method”. In: *J Phys. B-A. Mol. Opt.* 38.15 (2005), p. 2811.
- [142] Dong Xu, Jernej Stare, and Andrew L. Cooksy. “Solving the vibrational Schrödinger equation on an arbitrary multidimensional potential energy surface by the finite element method”. In: *Comput. Phys. Commun.* 180.11 (2009), pp. 2079–2094. DOI: <http://dx.doi.org/10.1016/j.cpc.2009.06.010>.
- [143] R. Alizadegan, K. J. Hsia, and T. J. Martinez. “A divide and conquer real space finite-element Hartree–Fock method”. In: *J. Chem. Phys.* 132.3, 034101 (2010). DOI: <http://dx.doi.org/10.1063/1.3290949>.
- [144] Jun Fang, Xingyu Gao, and Aihui Zhou. “A Kohn–Sham equation solver based on hexahedral finite elements”. In: *Journal of Computational Physics* 231.8 (2012), pp. 3166–3180. DOI: <http://dx.doi.org/10.1016/j.jcp.2011.12.043>.
- [145] W. Bangerth et al. “The deal.II Library, Version 8.4”. In: *Journal of Numerical Mathematics* 24 (2016).
- [146] F. Hecht. “New development in FreeFem++”. In: *J. Numer. Math.* 20.3-4 (2012), pp. 251–265.
- [147] Pavel Karban et al. “Numerical solution of coupled problems using code Agros2D”. In: *Computing* 95.1 (2013), pp. 381–408. DOI: 10.1007/s00607-013-0294-4.
- [148] B. Patzák. “OOFEM - an object-oriented simulation tool for advanced modeling of materials and structures”. In: *Acta Polytechnica* 52.6 (2012), pp. 59–66.
- [149] C. Perdigou and B. Audoly. “The viscous curtain: General formulation and finite-element solution for the stability of flowing viscous sheets”. In: *Journal of the Mechanics and Physics of Solids* 96 (2016), pp. 291–311. DOI: <http://dx.doi.org/10.1016/j.jmps.2016.07.015>.

- [150] F. Ballarin et al. “Fast simulations of patient-specific haemodynamics of coronary artery bypass grafts based on a POD-Galerkin method and a vascular shape parametrization”. In: *Journal of Computational Physics* 315 (June 2016), pp. 609–628.
- [151] H. Huang and B. W. Spencer. “Grizzly model of fully coupled heat transfer, moisture diffusion, alkali-silica reaction and fracturing processes in concrete”. In: *Proceedings of the 9th International Conference on Fracture Mechanics of Concrete and Concrete Structures (FraMCoS-9), Berkeley, CA*. Ed. by V. Saouma, J. Bolander, and E. Landis. Paper 194, <http://dx.doi.org/10.21012/FC9.194>. May 2016.
- [152] M. P. Lüthi et al. “Heat sources within the Greenland Ice Sheet: dissipation, temperate paleo-firn and cryo-hydrologic warming”. In: *The Cryosphere* 9.1 (2015), pp. 245–253. DOI: [10.5194/tc-9-245-2015](https://doi.org/10.5194/tc-9-245-2015).
- [153] P. Bauman, J. Jagodzinski, and B. S. Kirk. “Statistical calibration of thermocouple gauges used for inferring heat flux”. In: *42nd AIAA Thermophysics Conference, Fluid Dynamics and Co-located Conferences*. <http://dx.doi.org/10.2514/6.2011-3779>. June 2011.
- [154] Kyle A. Gamble et al. “A feasibility study on the use of the MOOSE computational framework to simulate three-dimensional deformation of CANDU reactor fuel elements”. In: *Nuclear Engineering and Design* 293 (2015), pp. 385–394. DOI: <http://dx.doi.org/10.1016/j.nucengdes.2015.07.028>.
- [155] Xianfeng Yang et al. “Fluid–structure interaction analysis of the drop impact test for helicopter fuel tank”. In: *SpringerPlus* 5.1 (2016), p. 1573. DOI: [10.1186/s40064-016-3040-5](https://doi.org/10.1186/s40064-016-3040-5).
- [156] Jörg Ackermann. “Finite-element-method expectation values for correlated two-electron wave functions”. In: *Phys. Rev. A* 52 (3 Sept. 1995), pp. 1968–1975. DOI: [10.1103/PhysRevA.52.1968](https://doi.org/10.1103/PhysRevA.52.1968).
- [157] Armin Scrinzi. “A 3-dimensional finite elements procedure for quantum mechanical applications”. In: *Comput. Phys. Commun.* 86.1 (1995), pp. 67–80. DOI: [http://dx.doi.org/10.1016/0010-4655\(94\)00160-4](http://dx.doi.org/10.1016/0010-4655(94)00160-4).
- [158] J.J. Soares Neto and Jan Linderberg. “A numerical study of various finite-element method schemes applied to quantum mechanical calculations”. In: *Comput. Phys. Commun.* 66.1 (1991), pp. 55–65. DOI: [http://dx.doi.org/10.1016/0010-4655\(91\)90007-8](http://dx.doi.org/10.1016/0010-4655(91)90007-8).
- [159] Weiying Zheng, Lung-an Ying, and Peizhu Ding. “Numerical solutions of the Schrödinger equation for the ground lithium by the finite element method”. In: *Appl. Math. Comput.* 153.3 (2004), pp. 685–695. DOI: [http://dx.doi.org/10.1016/S0096-3003\(03\)00664-7](http://dx.doi.org/10.1016/S0096-3003(03)00664-7).
- [160] J. Ackermann, B. Erdmann, and R. Roitzsch. “A self-adaptive multilevel finite element method for the stationary Schrödinger equation in three space dimensions”. In: *J. Chem. Phys.* 101.9 (1994), pp. 7643–7650. DOI: <http://dx.doi.org/10.1063/1.468257>.

- [161] J. E. Pask et al. “Real-space local polynomial basis for solid-state electronic-structure calculations: A finite-element approach”. In: *Phys. Rev. B* 59 (19 May 1999), pp. 12352–12358. DOI: [10.1103/PhysRevB.59.12352](https://doi.org/10.1103/PhysRevB.59.12352).
- [162] Jan Linderberg. “Finite element methods in quantum mechanics”. In: *Computer Physics Reports* 6.1 (1987), pp. 209–242. DOI: [http://dx.doi.org/10.1016/0167-7977\(87\)90013-X](http://dx.doi.org/10.1016/0167-7977(87)90013-X).
- [163] Lauri Lehtovaara, Ville Havu, and Martti Puska. “All-electron density functional theory and time-dependent density functional theory with high-order finite elements”. In: *J. Chem. Phys.* 131.5, 054103 (2009). DOI: <http://dx.doi.org/10.1063/1.3176508>.
- [164] P. Motamarri et al. “Higher-order adaptive finite-element methods for Kohn–Sham density functional theory”. In: *Journal of Computational Physics* 253 (2013), pp. 308–343. DOI: <http://dx.doi.org/10.1016/j.jcp.2013.06.042>.
- [165] E. Romero and J. E. Roman. “A parallel implementation of Davidson methods for large-scale eigenvalue problems in SLEPc”. In: *ACM Trans. Math. Software* 40.2 (2014), 13:1–13:29. DOI: [10.1145/2543696](https://doi.org/10.1145/2543696).
- [166] V. Hernandez, J.E. Roman, and A. Tomas. “Parallel Arnoldi eigensolvers with enhanced scalability via global communications rearrangement”. In: *Parallel Computing* 33.7–8 (2007), pp. 521–540. DOI: <http://dx.doi.org/10.1016/j.parco.2007.04.004>.
- [167] Joseph M. Elble, Nikolaos V. Sahinidis, and Panagiotis Vouzis. “GPU computing with Kaczmarz’s and other iterative algorithms for linear systems”. In: *Parallel Computing* 36.5–6 (2010). Parallel Matrix Algorithms and Applications, pp. 215–231. DOI: <http://dx.doi.org/10.1016/j.parco.2009.12.003>.
- [168] Filipp Furche et al. “Accelerating molecular property calculations with nonorthonormal Krylov space methods”. In: *The Journal of Chemical Physics* 144.17, 174105 (2016). DOI: <http://dx.doi.org/10.1063/1.4947245>.
- [169] I. Babuska, B. A. Szabo, and I. N. Katz. “The p-Version of the Finite Element Method”. In: *SIAM Journal on Numerical Analysis* 18.3 (1981), pp. 515–545.
- [170] W. Rachowicz, D. Pardo, and L. Demkowicz. “Fully automatic hp-adaptivity in three dimensions”. In: *Computer Methods in Applied Mechanics and Engineering* 195.37–40 (2006). John H. Argyris Memorial Issue. Part I, pp. 4816–4842. DOI: <http://dx.doi.org/10.1016/j.cma.2005.08.022>.
- [171] Andreas Meinken. *Finite Elemente in der Praxis*. 1st ed. Expert Verlag, 2001.
- [172] Zhong-Ci Shi. “Nonconforming finite element methods”. In: *Journal of Computational and Applied Mathematics* 149.1 (2002). Scientific and Engineering Computations for the 21st Century - Methodologies and Applications Proceedings of the 15th Toyota Conference, pp. 221–225. DOI: [http://dx.doi.org/10.1016/S0377-0427\(02\)00531-9](http://dx.doi.org/10.1016/S0377-0427(02)00531-9).
- [173] Siu-Wing Cheng and Jonathan Dey Tamal K. and Shewchuk. *Delaunay Mesh Generation*. 1st ed. Taylor & Francis, 2012.

- [174] V.I. Lebedev and D.N. Laikov. “A quadrature formula for the sphere of the 131st-algebraic order of accuracy”. In: *Doklady Mathematics* 59.3 (1999), pp. 477–481.
- [175] Robert S. Womersley and Ian H. Sloan. “How good can polynomial interpolation on the sphere be?” In: *Advances in Computational Mathematics* 14.3 (2001), pp. 195–226. DOI: [10.1023/A:1016630227163](https://doi.org/10.1023/A:1016630227163).
- [176] Ian H. Sloan and Robert S. Womersley. “Good approximation on the sphere, with application to geodesy and the scattering of sound”. In: *Journal of Computational and Applied Mathematics* 149.1 (2002). Scientific and Engineering Computations for the 21st Century - Methodologies and Applications Proceedings of the 15th Toyota Conference, pp. 227–237. DOI: [http://dx.doi.org/10.1016/S0377-0427\(02\)00532-0](http://dx.doi.org/10.1016/S0377-0427(02)00532-0).
- [177] J.G. Muga et al. “Complex absorbing potentials”. In: *Physics Reports* 395.6 (2004), pp. 357–426. DOI: <http://dx.doi.org/10.1016/j.physrep.2004.03.002>.
- [178] U V Riss and H-D Meyer. “The transformative complex absorbing potential method: a bridge between complex absorbing potentials and smooth exterior scaling”. In: *Journal of Physics B: Atomic, Molecular and Optical Physics* 31.10 (1998), p. 2279.
- [179] Thomas-C. Jagau et al. “A Fresh Look at Resonances and Complex Absorbing Potentials: Density Matrix-Based Approach”. In: *JPCL* 5.2 (Jan. 16, 2014), pp. 310–315. DOI: [10.1021/jz402482a](https://doi.org/10.1021/jz402482a).
- [180] Bjorn Engquist and Andrew Majda. “Absorbing boundary conditions for the numerical simulation of waves”. In: *Math. Comp.* 31.0436612 (1977), pp. 629–651.
- [181] Steffen Marburg and Bodo Nolte, eds. *Computational Acoustics of Noise Propagation in Fluids - Finite and Boundary Element Methods*. Springer, 2008.
- [182] Francis Collino. “Perfectly Matched Absorbing Layers for the Paraxial Equations”. In: *Journal of Computational Physics* 131.1 (1997), pp. 164–180. DOI: <http://dx.doi.org/10.1006/jcph.1996.5594>.
- [183] Chunxiong Zheng. “A perfectly matched layer approach to the nonlinear Schrödinger wave equations”. In: *Journal of Computational Physics* 227.1 (2007), pp. 537–556. DOI: <http://dx.doi.org/10.1016/j.jcp.2007.08.004>.
- [184] Riyad Kechroud et al. “Preconditioning techniques for the solution of the Helmholtz equation by the finite element method”. In: *Mathematics and Computers in Simulation* 65.4–5 (2004). Wave Phenomena in Physics and Engineering: New Models, Algorithms, and Applications, pp. 303–321. DOI: <http://dx.doi.org/10.1016/j.matcom.2004.01.004>.
- [185] R. J. Astley. “FE mode-matching schemes for the exterior Helmholtz problem and their relationship to the FE-DtN approach”. In: *Communications in Numerical Methods in Engineering* 12.4 (1996), pp. 257–267. DOI: [10.1002/\(SICI\)1099-0887\(199604\)12:4<257::AID-CNM977>3.0.CO;2-8](https://doi.org/10.1002/(SICI)1099-0887(199604)12:4<257::AID-CNM977>3.0.CO;2-8).
- [186] D.N. Dai. “An improved boundary element formulation for wave propagation problems”. In: *Engineering Analysis with Boundary Elements* 10.4 (1992), pp. 277–281. DOI: [http://dx.doi.org/10.1016/0955-7997\(92\)90140-3](http://dx.doi.org/10.1016/0955-7997(92)90140-3).

- [187] Martin Costabel. “Principles of boundary element methods”. In: *Computer Physics Reports* 6.1 (1987), pp. 243–274. DOI: [http://dx.doi.org/10.1016/0167-7977\(87\)90014-1](http://dx.doi.org/10.1016/0167-7977(87)90014-1).
- [188] Whye-Teong Ang. *A Beginner’s Course in Boundary Element Methods*. Boca Raton, Florida: Universal Publishers, 2007.
- [189] Santosh Kumar Rana and Amit Jena. “A BEM formulation of two dimensional steady state heat conduction in exchanger tubes of arbitrary cross sections”. In: *International Journal of Heat and Mass Transfer* 106 (2017), pp. 195–211. DOI: <http://dx.doi.org/10.1016/j.ijheatmasstransfer.2016.10.055>.
- [190] Y.P. Gong, C.Y. Dong, and X.C. Qin. “An isogeometric boundary element method for three dimensional potential problems”. In: *Journal of Computational and Applied Mathematics* 313 (2017), pp. 454–468. DOI: <http://dx.doi.org/10.1016/j.cam.2016.10.003>.
- [191] Haijun Wu et al. “A Fast Multipole Boundary Element Method for Three-Dimensional Half-Space Acoustic Wave Problems Over an Impedance Plane”. In: *International Journal of Computational Methods* 12.01 (2015), p. 1350090. DOI: [10.1142/S0219876213500904](https://doi.org/10.1142/S0219876213500904).
- [192] F. Guerracino, V. Minutolo, and L. Nunziante. “A simple analysis of soil-structure interaction by BEM-FEM coupling”. In: *Engineering Analysis with Boundary Elements* 10.4 (1992), pp. 283–289. DOI: [http://dx.doi.org/10.1016/0955-7997\(92\)90141-S](http://dx.doi.org/10.1016/0955-7997(92)90141-S).
- [193] O. C. Zienkiewicz, D. W. Kelly, and P. Bettess. “The coupling of the finite element method and boundary solution procedures”. In: *International Journal for Numerical Methods in Engineering* 11.2 (1977), pp. 355–375. DOI: [10.1002/nme.1620110210](https://doi.org/10.1002/nme.1620110210).
- [194] D. Dreyer and O. Von Estorff. “Improved conditioning of infinite elements for exterior acoustics”. In: *International Journal for Numerical Methods in Engineering* 58.6 (2003). 00049, pp. 933–953.
- [195] J. E. Roman et al. *SLEPc Users Manual*. Tech. rep. DSIC-II/24/02 - Revision 3.7. D. Sistemes Informàtics i Computació, Universitat Politècnica de València, 2016.
- [196] V. Hernandez et al. *A survey of software for sparse eigenvalue problems*. Tech. rep. STR-7. Available at <http://slepc.upv.es>. Universitat Politècnica de València, 2009.
- [197] Uwe Helmke and Jens Jordan. “Control and Stabilization of Linear Equation Solvers”. In: *Perspectives in Mathematical System Theory, Control, and Signal Processing: A Festschrift in Honor of Yutaka Yamamoto on the Occasion of his 60th Birthday*. Ed. by Jan C. Willems et al. Berlin, Heidelberg: Springer Berlin Heidelberg, 2010, pp. 73–82. DOI: [10.1007/978-3-540-93918-4\\_7](https://doi.org/10.1007/978-3-540-93918-4_7).
- [198] R. Shamasundar and W.A. Mulder. “Improving the accuracy of mass-lumped finite-elements in the first-order formulation of the wave equation by defect correction”. In: *Journal of Computational Physics* 322 (2016), pp. 689–707. DOI: <http://dx.doi.org/10.1016/j.jcp.2016.07.006>.

- [199] AmirHossein Aminfar and Eric Darve. “A fast, memory efficient and robust sparse preconditioner based on a multifrontal approach with applications to finite-element matrices”. In: *International Journal for Numerical Methods in Engineering* 107.6 (2016), pp. 520–540. DOI: [10.1002/nme.5196](https://doi.org/10.1002/nme.5196).
- [200] Wolfgang Dahmen and Angela Kunoth. “Multilevel preconditioning”. In: *Numerische Mathematik* 63.1 (1992), pp. 315–344. DOI: [10.1007/BF01385864](https://doi.org/10.1007/BF01385864).
- [201] Harry Yserentant. “On the multi-level splitting of finite element spaces”. In: *Numerische Mathematik* 49.4 (1986), pp. 379–412. DOI: [10.1007/BF01389538](https://doi.org/10.1007/BF01389538).
- [202] Noriyuki Kushida. “Newton–Raphson preconditioner for Krylov type solvers on GPU devices”. In: *SpringerPlus* 5.1 (2016), p. 788. DOI: [10.1186/s40064-016-2346-7](https://doi.org/10.1186/s40064-016-2346-7).
- [203] V. Hernandez et al. *A survey of software for sparse eigenvalue problems*. Tech. rep. STR-4. Available at <http://slepc.upv.es>. Universitat Politècnica de València, 2009.
- [204] V. Hernandez et al. *A survey of software for sparse eigenvalue problems*. Tech. rep. STR-5. Available at <http://slepc.upv.es>. Universitat Politècnica de València, 2009.
- [205] G. W. Stewart. “A Krylov–Schur Algorithm for Large Eigenproblems”. In: *SIAM Journal on Matrix Analysis and Applications* 23.3 (2002), pp. 601–614. DOI: [10.1137/S0895479800371529](https://doi.org/10.1137/S0895479800371529). eprint: <http://dx.doi.org/10.1137/S0895479800371529>.
- [206] Wolfram Koch and Max C. Holthausen. *A Chemist’s Guide to Density Functional Theory*. 2nd ed. Wiley-VCH Verlag GmbH, 2001.
- [207] A. D. Becke. “Density-functional exchange-energy approximation with correct asymptotic behavior”. In: *Phys. Rev. A* 38 (6 Sept. 1988), pp. 3098–3100. DOI: [10.1103/PhysRevA.38.3098](https://doi.org/10.1103/PhysRevA.38.3098).
- [208] Chengteh Lee, Weitao Yang, and Robert G. Parr. “Development of the Colle-Salvetti correlation-energy formula into a functional of the electron density”. In: *Phys. Rev. B* 37 (2 Jan. 1988), pp. 785–789. DOI: [10.1103/PhysRevB.37.785](https://doi.org/10.1103/PhysRevB.37.785).
- [209] Erich Runge and E. K. U. Gross. “Density-Functional Theory for Time-Dependent Systems”. In: *Phys. Rev. Lett.* 52 (12 Mar. 1984), pp. 997–1000. DOI: [10.1103/PhysRevLett.52.997](https://doi.org/10.1103/PhysRevLett.52.997).
- [210] A. Dreuw and M. Head-Gordon. “Single-Reference ab Initio Methods for the Calculation of Excited States of Large Molecules”. In: *Chem. Rev.* 105.11 (2005), pp. 4009–4035.
- [211] M.E. Casida, A. Ipatov, and F. Cordova. “Linear-Response Time-Dependent Density Functional Theory for Open-Shell Molecules”. In: *Time-Dependent Density Functional Theory*. Ed. by Miguel A.L. Marques et al. Berlin, Heidelberg: Springer Berlin Heidelberg, 2006, pp. 243–257. DOI: [10.1007/3-540-35426-3\\_16](https://doi.org/10.1007/3-540-35426-3_16).
- [212] Paula Mori-Sanchez and Aron J. Cohen. “The derivative discontinuity of the exchange-correlation functional”. In: *Phys. Chem. Chem. Phys.* 16 (28 2014), pp. 14378–14387. DOI: [10.1039/C4CP01170H](https://doi.org/10.1039/C4CP01170H).

- [213] R. Ditchfield, W. J. Hehre, and J. A. Pople. “Self-Consistent Molecular-Orbital Methods. IX. An Extended Gaussian-Type Basis for Molecular-Orbital Studies of Organic Molecules”. In: *The Journal of Chemical Physics* 54.2 (1971), pp. 724–728. DOI: <http://dx.doi.org/10.1063/1.1674902>.
- [214] G. A. Petersson et al. “A complete basis set model chemistry. I. The total energies of closed-shell atoms and hydrides of the first-row elements”. In: *The Journal of Chemical Physics* 89.4 (1988), pp. 2193–2218. DOI: <http://dx.doi.org/10.1063/1.455064>.
- [215] Hisayoshi Iikura et al. “A long-range correction scheme for generalized-gradient-approximation exchange functionals”. In: *The Journal of Chemical Physics* 115.8 (2001), pp. 3540–3544. DOI: <http://dx.doi.org/10.1063/1.1383587>.
- [216] Florian Weigend and Reinhart Ahlrichs. “Balanced basis sets of split valence, triple zeta valence and quadruple zeta valence quality for H to Rn: Design and assessment of accuracy”. In: *Phys. Chem. Chem. Phys.* 7 (18 2005), pp. 3297–3305. DOI: 10.1039/B508541A.
- [217] Hang Si. “TetGen, a Delaunay-Based Quality Tetrahedral Mesh Generator”. In: *ACM Trans. Math. Softw.* 41.2 (Feb. 2015), 11:1–11:36. DOI: 10.1145/2629697.
- [218] Hirofumi Tomita et al. “Shallow Water Model on a Modified Icosahedral Geodesic Grid by Using Spring Dynamics”. In: *Journal of Computational Physics* 174.2 (2001), pp. 579–613. DOI: <http://dx.doi.org/10.1006/jcph.2001.6897>.
- [219] D. A. Randall et al. “Climate modeling with spherical geodesic grids”. In: *Computing in Science Engineering* 4.5 (Sept. 2002), pp. 32–41. DOI: 10.1109/MCISE.2002.1032427.
- [220] Robert S. Womersley and Ian H. Sloan. “How good can polynomial interpolation on the sphere be?” In: *Advances in Computational Mathematics* 14.3 (2001), pp. 195–226. DOI: 10.1023/A:1016630227163.
- [221] J Fliege and U Maier. “The distribution of points on the sphere and corresponding cubature formulae”. In: *IMA Journal of Numerical Analysis* 19.2 (1999), pp. 317–334. DOI: 10.1093/imanum/19.2.317. eprint: <http://imajna.oxfordjournals.org/content/19/2/317.full.pdf+html>.
- [222] Mark A. Spackman. “Potential derived charges using a geodesic point selection scheme”. In: *Journal of Computational Chemistry* 17.1 (1996), pp. 1–18. DOI: 10.1002/(SICI)1096-987X(19960115)17:1<1::AID-JCC1>3.0.CO;2-V.
- [223] Guo-Zhu Zhu and Lai-Sheng Wang. “Communication: Vibrationally resolved photoelectron spectroscopy of the tetracyanoquinodimethane (TCNQ) anion and accurate determination of the electron affinity of TCNQ”. In: *J. Chem. Phys.* 143.22, 221102 (2015). DOI: <http://dx.doi.org/10.1063/1.4937761>.
- [224] J B West et al. “A high resolution angle resolved photoelectron spectroscopy study of N<sub>2</sub>”. In: *Physica Scripta* 41.4 (1990), p. 487.

- [225] Sunil R. Desai, Hongbin Wu, and Lai-Sheng Wang. “Vibrationally resolved photoelectron spectroscopy of AlO- and AlO<sub>2</sub><sup>-</sup>”. In: *International Journal of Mass Spectrometry and Ion Processes* 159.1 (1996), pp. 75–80. DOI: [http://dx.doi.org/10.1016/S0168-1176\(96\)04443-6](http://dx.doi.org/10.1016/S0168-1176(96)04443-6).
- [226] S. Alitalo et al. “The valence photoelectron satellite spectra of Kr and Xe”. In: *Journal of Electron Spectroscopy and Related Phenomena* 114-116 (2001). Proceeding of the Eight International Conference on Electronic Spectroscopy and Structure, pp. 141–146. DOI: [http://dx.doi.org/10.1016/S0368-2048\(00\)00277-2](http://dx.doi.org/10.1016/S0368-2048(00)00277-2).
- [227] C.G. Ning et al. “Probing Dyson orbitals with Green’s Function theory and Electron Momentum Spectroscopy”. In: *Chem. Phys. Lett.* 421.1–3 (2006), pp. 52–57. DOI: <http://dx.doi.org/10.1016/j.cplett.2006.01.040>.
- [228] S Krummacher et al. “Inner-shell photoemission studies of lithium and sodium vapour”. In: *Journal of Physics B: Atomic and Molecular Physics* 15.23 (1982), p. 4363.
- [229] Saha H. P. and Lin Dong. “Theoretical study of the valence and core photoionization of a three-electron system from lithium to neon”. In: *Journal of Physics B: Atomic, Molecular and Optical Physics* 30.7 (1997), p. 1651.
- [230] I. Reineck et al. “High-resolution UV photoelectron spectrum of CO<sub>2</sub>”. In: *Chemical Physics* 78.3 (1983), pp. 311–318. DOI: [http://dx.doi.org/10.1016/0301-0104\(83\)85116-7](http://dx.doi.org/10.1016/0301-0104(83)85116-7).
- [231] H. Veenhuizen et al. “High resolution angle-resolved photoelectron spectrum of CO<sub>2</sub>, excited with polarized resonance radiation”. In: *Journal of Electron Spectroscopy and Related Phenomena* 41.2 (1986), pp. 205–223. DOI: [http://dx.doi.org/10.1016/0368-2048\(86\)85004-6](http://dx.doi.org/10.1016/0368-2048(86)85004-6).
- [232] Lai-Sheng Wang et al. “High resolution UV photoelectron spectroscopy of CO+2, COS+ and CS+2 using supersonic molecular beams”. In: *Journal of Electron Spectroscopy and Related Phenomena* 47 (1988), pp. 167–186. DOI: [http://dx.doi.org/10.1016/0368-2048\(88\)85010-2](http://dx.doi.org/10.1016/0368-2048(88)85010-2).
- [233] C.R. Brundle and D.W. Turner. “Studies on the photoionisation of the linear triatomic molecules: N<sub>2</sub>O, COS, CS<sub>2</sub> and CO<sub>2</sub> using high-resolution photoelectron spectroscopy”. In: *International Journal of Mass Spectrometry and Ion Physics* 2.3 (1969), pp. 195–220. DOI: [http://dx.doi.org/10.1016/0020-7381\(69\)80018-5](http://dx.doi.org/10.1016/0020-7381(69)80018-5).
- [234] S.Y. Truong et al. “Threshold photoelectron spectroscopy of H<sub>2</sub>O and D<sub>2</sub>O over the photon energy range 12–40eV”. In: *Chemical Physics* 355.2 (Jan. 2009), pp. 183–193. DOI: [10.1016/j.chemphys.2008.12.009](https://doi.org/10.1016/j.chemphys.2008.12.009).
- [235] L. Karlsson. “Isotopic and vibronic coupling effects in the valence electron spectra of H<sub>2</sub> 16O, H<sub>2</sub> 18O, and D<sub>2</sub> 16O”. In: *J. Chem. Phys.* 62.12 (1975), p. 4745. DOI: [10.1063/1.430423](https://doi.org/10.1063/1.430423).
- [236] C.G. Ning et al. “High resolution electron momentum spectroscopy of the valence orbitals of water”. In: *Chemical Physics* 343.1 (Jan. 2008), pp. 19–30. DOI: [10.1016/j.chemphys.2007.09.030](https://doi.org/10.1016/j.chemphys.2007.09.030).

# A | Appendix

## A.1 Additional Data

## A.2 Delaunay Triangulation

The explicit setup of non-regular meshes for the use in finite element schemes is not trivial at all. There are however schemes available, among which are the Delaunay and Voronoi tessellations which are their respective dual schemes. Since in this thesis, only Delaunay tessellation is used, I will

focus on this method and its properties only. In mathematics one understands under a triangulation a connection of points to simplices. Thereby a structure is denoted as a simplex in  $n$  dimensions, if has as few vertices as possible in this dimension. A simplex in  $2D$  e.g. is a triangle while it is in  $3D$  a tetrahedron. Moreover, a simplex is denoted as Delaunay simplex if there is a circumsphere such that no vertex is inside of this sphere.

A bit more intuitive access to this scheme can be obtained via the Voronoi diagram: A Voronoi diagram splits a given volume (in  $3D$ ) into elements using a set of points  $p_i$  in this volume by assigning each point to the element of his nearest point  $p_i$ . Having this tesselation, one can transform it to a set of Delaunay simplices by connecting the points  $p_i$  each with their direct neighbours [217].




Figure A.1: The eigen energies of the first 20 solutions (circles) for different powers  $p$  of the damping function  $D(r)$  which determines the colours. The spectra do not change significantly for  $p < \frac{1}{8}$ .