

Finding Mean Absolute Deviation (MAD)

1. The mean is given. Find the MAD.

a. 5, 7, 9, 11 (mean 8) _____

b. 49, 50, 50, 51 (mean 50) _____

2. Tavon found distances of -4 , -1 , 2 and 3 from the mean and said the MAD is 0. Find the correct MAD.

Show your work.

MAD _____

3. A data set has a mean of 72 and a MAD of 4. Which is true?

A Every value is 72.

B A typical value is about 4 from 72.

C The range is 4.

D Every value is 4 from 72.

HOW DID IT GO? ☐ I got it ☐ Almost ☐ I need help

cut here

Finding Mean Absolute Deviation (MAD)

1. The mean is given. Find the MAD.

a. 5, 7, 9, 11 (mean 8) _____

b. 49, 50, 50, 51 (mean 50) _____

2. Tavon found distances of -4 , -1 , 2 and 3 from the mean and said the MAD is 0. Find the correct MAD.

Show your work.

MAD _____

3. A data set has a mean of 72 and a MAD of 4. Which is true?

A Every value is 72.

B A typical value is about 4 from 72.

C The range is 4.

D Every value is 4 from 72.

HOW DID IT GO? ☐ I got it ☐ Almost ☐ I need help

Comparing Data Sets Using MAD

1. Both sets have a mean of 13. Find each MAD.

a. X: 10, 12, 14, 16 _____

b. Y: 4, 12, 14, 22 _____

2. Set M has mean 25 and MAD 1.5. Set N has mean 25 and MAD 7. Which set is steadier?

Show your work.

Set _____

3. Two classes both have a mean score of 75. Which statement must be true?

A Their MADs are equal.

B Their spreads may differ.

C Their scores are identical.

D Their ranges are equal.

HOW DID IT GO? ☐ I got it ☐ Almost ☐ I need help

cut here

Comparing Data Sets Using MAD

1. Both sets have a mean of 13. Find each MAD.

a. X: 10, 12, 14, 16 _____

b. Y: 4, 12, 14, 22 _____

2. Set M has mean 25 and MAD 1.5. Set N has mean 25 and MAD 7. Which set is steadier?

Show your work.

Set _____

3. Two classes both have a mean score of 75. Which statement must be true?

A Their MADs are equal.

B Their spreads may differ.

C Their scores are identical.

D Their ranges are equal.

HOW DID IT GO? ☐ I got it ☐ Almost ☐ I need help