Lesson 6.6 Teaching Support Guide

Teach-tomorrow route

Print: current Guided Notes and Practice A for the class; Exit Tickets; optional Language Support and supported homework. Keep this guide and answer keys teacher-facing. Use the full-size model pages in the updated notes.

Phase	Planned minutes	Teaching move
Retrieve	5	Warm-Up items 1–4; note the missing prerequisite before launch.
Launch	7	Name the session target and invite a prediction from the conceptual task.
Model	10	Link a visual model to the worked example; rehearse the mathematical explanation.
Practice	18	Core: A #1, #2, #4, #6 and the error task. Choose remaining items from evidence.
Check	5	Exit Ticket; use the diagnostic page before assigning reteaching.

This is a planned 45-minute route, not a measured classroom duration. For a shorter lesson, keep the model and exit; reduce repeated practice. Form B is parallel practice or reassessment only when its items are unseen.

S1: Simplifying to Show Equivalence · I can simplify an expression by combining like terms to test whether two expressions are equivalent.

S2: Testing and Justifying Equivalence · I can test whether two expressions are equivalent by substituting values and by reasoning.

Spaced retrieval: revisit one item after 2 lessons and after approximately 1 week. Mix it with the next unit’s prerequisite check; do not announce the operation before students choose a method.

Session 1 · Simplifying to Show Equivalence

I can simplify an expression by combining like terms to test whether two expressions are equivalent. · 6.EE.A.3

Observe and respond

Exit item	Expected evidence	Next prompt/model
1	a. 13x b. 5y + 5	Make one group of variable tiles and one group of units before writing the sum.
2	7k + 6	Make one group of variable tiles and one group of units before writing the sum.
3	A — 9n + 2	Ask the learner to show why A — 9n + 2 is correct and reject a specific distractor using a model.

Distinguish the error

Conceptual signal: Combines variable terms and constants into one coefficient..

Response: Make one group of variable tiles and one group of units before writing the sum.

Also check: Combining a constant with a variable term (12x + 3 + 4x as 19x).

Also check: Treating 5n + 5n as n^2 or $10n^2$ instead of 10n.

If calculation alone is wrong, use Session 1 Prerequisite Practice A for the matching skill, then return to the model. If the interpretation is wrong, label the quantities before reteaching the operation. Independent re-entry requires a correct setup and a connected explanation.

Session 1 · Fresh check and scoring

Reserve this check until after revision; it is distinct from the exit ticket. Do not show this teacher page to learners beforehand.

Simplify 7a + 2 + 3a + 4 and explain the two groups.

Teacher solution: 10a + 6; variable terms combine separately from constants.

Score the evidence

Score	Evidence	Next action
3	Valid representation, correct result/units, connected explanation.	Continue; fade a support next time.
2	Correct setup and meaning with a calculation/notation slip, or an incomplete explanation.	Brief targeted feedback, then another unseen check.
1	Some relevant facts but a model or interpretation error.	Use the named model and the corresponding prerequisite page.
0	No interpretable evidence yet.	Oral/diagram response to separate access needs from mathematics.

Complete response anchor: Six x-tiles and eight unit tiles give 6x + 8. Like terms refer to the same variable part; unit constants cannot be added to the coefficient.

Misconception response anchor: Combines variable terms and constants into one coefficient.. Use make one group of variable tiles and one group of units before writing the sum.

Language access: point/draw and label → oral rehearsal → relationship sentence → independent explanation. Offer the support needed for this task and fade it based on evidence. Suggested frame: My model or expression represents ___ because ___. The structure shows ___, so the result is ___.

Session 2 · Testing and Justifying Equivalence

I can test whether two expressions are equivalent by substituting values and by reasoning. · 6.EE.A.4

Observe and respond

Exit item	Expected evidence	Next prompt/model
1	a. 12 b. 9	Use properties for an all-values argument and one differing value to reject a claim.
2	m = 1: 6 and 6; m = 4: 15 and 24. Not equivalent.	Use properties for an all-values argument and one differing value to reject a claim.
3	C — Distributive	Ask the learner to show why C — Distributive is correct and reject a specific distractor using a model.

Distinguish the error

Conceptual signal: Treats one matching substitution as proof of equivalence..

Response: Use properties for an all-values argument and one differing value to reject a claim.

Also check: Deciding two expressions are equivalent after one matching test value.

Also check: Distributing to only the first term (3(k + 2) as 3k + 2).

If calculation alone is wrong, use Session 2 Prerequisite Practice A for the matching skill, then return to the model. If the interpretation is wrong, label the quantities before reteaching the operation. Independent re-entry requires a correct setup and a connected explanation.

Session 2 · Fresh check and scoring

Reserve this check until after revision; it is distinct from the exit ticket. Do not show this teacher page to learners beforehand.

Compare 2(y+5) and 2y+10, then compare 2y+10 and 4y.

Teacher solution: First equivalent by distribution; second fails at y=1: 12 versus 4.

Score the evidence

Score	Evidence	Next action
3	Valid representation, correct result/units, connected explanation.	Continue; fade a support next time.
2	Correct setup and meaning with a calculation/notation slip, or an incomplete explanation.	Brief targeted feedback, then another unseen check.
1	Some relevant facts but a model or interpretation error.	Use the named model and the corresponding prerequisite page.
0	No interpretable evidence yet.	Oral/diagram response to separate access needs from mathematics.

Complete response anchor: First pair equivalent by distribution. Second not equivalent: x=1 gives 9 versus 6. One counterexample disproves equivalence; one matching test does not prove it.

Misconception response anchor: Treats one matching substitution as proof of equivalence.. Use use properties for an all-values argument and one differing value to reject a claim.

Language access: point/draw and label → oral rehearsal → relationship sentence → independent explanation. Offer the support needed for this task and fade it based on evidence. Suggested frame: My model or expression represents ___ because ___. The structure shows ___, so the result is ___.