



Project Presentation:



Bar Amit






Random Hypothesis Research

Research question: Does weight affect blood sugar levels?

Research findings:

- There is a significant jump in weight when moving from level 1 to level 2, but levels 2 and 3 show an almost identical weight range.
- Above a certain weight, blood sugar tends to increase, but weight alone cannot distinguish between high and very high blood sugar levels.
- It is possible to find people with very high weight and normal blood sugar levels, as well as people with low weight and high blood sugar levels.

Research conclusions:



High weight affects the transition to abnormal blood sugar levels, but it does not distinguish between high and very high levels; at these stages, other variables have a greater influence. The existence of exceptions—people with high weight and normal blood sugar—proves that obesity is a significant risk factor, but it is not the only factor determining blood sugar levels.

(1)


Research question: Is there a relationship between high cholesterol and weight?

Research findings:

1. As a person's weight increases, their cholesterol level also increases.
2. People with healthy cholesterol levels are usually within a weight range of around 75 kg.

Research conclusions:

1. We discovered that people with normal cholesterol levels tend to weigh less compared to people with high (bad) cholesterol levels.





Conclusion

The results of my research show weight as a significant risk factor that increases the likelihood of developing heart disease. My research showed that weight has a positive effect on blood sugar levels (as weight increases, blood sugar levels also increase), and high blood sugar is one of the main and most common factors contributing to heart disease and diabetes.

In addition, the research showed that as cholesterol levels increase, weight also tends to increase. Combining these findings can help indicate the level of risk for certain individuals. For example, an older person with excess weight may be at a “double risk” compared to a person with only one of these risk factors. This can help determine the urgency of treatment, identify which individuals are at higher risk than others, and decide who should receive treatment first.

