

## Question 1

Correct

Mark 1.00 out of 1.00

Find Duplicate in Array.

Given a read only array of n integers between 1 and n, find one number that repeats.

Input Format:

First Line - Number of elements

n Lines - n Elements

Output Format:

Element x - That is repeated

**For example:**

Input	Result
5 1 1 2 3 4	1

**Answer:** (penalty regime: 0 %)

```

1  #include <stdio.h>
2  int main(){
3      int n;
4      scanf("%d",&n);
5      int arr[n];
6      for(int i=0;i<n;i++){
7          scanf("%d",&arr[i]);
8      }
9      int k=0;
10     for(int i=0;i<n-1;i++){
11         for(int j=i+1;j<n;j++){
12             if(arr[i]==arr[j]){
13                 printf("%d",arr[i]);
14                 k=1;
15                 break;
16             }
17         }
18         if(k==1){
19             break;
20         }
21     }
22 }
```

	Input	Expected	Got	
✓	11 10 9 7 6 5 1 2 3 8 4 7	7	7	✓
✓	5 1 2 3 4 4	4	4	✓
✓	5 1 1 2 3 4	1	1	✓

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.

## Question 1

Correct

Mark 1.00 out of 1.00

Find Duplicate in Array.

Given a read only array of n integers between 1 and n, find one number that repeats.

Input Format:

First Line - Number of elements

n Lines - n Elements

Output Format:

Element x - That is repeated

**For example:**

Input	Result
5 1 1 2 3 4	1

**Answer:** (penalty regime: 0 %)

```

1  #include <stdio.h>
2  int main(){
3      int n;
4      scanf("%d",&n);
5      int arr[n];
6      for(int i=0;i<n;i++){
7          scanf("%d",&arr[i]);
8      }
9      int k=0;
10     for(int i=0;i<n-1;i++){
11         for(int j=i+1;j<n;j++){
12             if(arr[i]==arr[j]){
13                 printf("%d",arr[i]);
14                 k=1;
15                 break;
16             }
17         }
18         if(k==1){
19             break;
20         }
21     }
22 }
```

	Input	Expected	Got	
✓	11 10 9 7 6 5 1 2 3 8 4 7	7	7	✓
✓	5 1 2 3 4 4	4	4	✓
✓	5 1 1 2 3 4	1	1	✓

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.

## Question 1

Correct

Mark 1.00 out of 1.00

Find the intersection of two sorted arrays.

OR in other words,

Given 2 sorted arrays, find all the elements which occur in both the arrays.

Input Format

· The first line contains T, the number of test cases. Following T lines contain:

1. Line 1 contains N1, followed by N1 integers of the first array
2. Line 2 contains N2, followed by N2 integers of the second array

Output Format

The intersection of the arrays in a single line

Example

Input:

```
1
3 10 17 57
6 2 7 10 15 57 246
```

Output:

```
10 57
```

Input:

```
1
6 1 2 3 4 5 6
2 1 6
```

Output:

```
1 6
```

**For example:**

Input	Result
1 3 10 17 57 6 2 7 10 15 57 246	10 57

**Answer:** (penalty regime: 0 %)

```
1 #include <stdio.h>
2 int main(){
3     int t;
4     scanf("%d",&t);
5     int n;
6     scanf("%d",&n);
7     int arr[n];
8     for(int i=0;i<n;i++){
9         scanf("%d",&arr[i]);
10    }
11    int n1;
12    scanf("%d",&n1);
13    int arr1[n1];
14    for(int i=0;i<n1;i++){
15        scanf("%d",&arr1[i]);
```

```
16     }
17     for(int i=0;i<n;i++){
18         for(int j=0;j<n1;j++){
19             if(arr[i]==arr1[j]){
20                 // printf("Dfg");
21                 printf("%d ",arr[i]);
22             }
23         }
24     }
25 }
```

	Input	Expected	Got	
✓	1 3 10 17 57 6 2 7 10 15 57 246	10 57	10 57	✓
✓	1 6 1 2 3 4 5 6 2 1 6	1 6	1 6	✓

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.

◀ 2-Finding Duplicates-O(n) Time Complexity,O(1) Space Complexity

Jump to...

4-Print Intersection of 2 sorted arrays-O(m+n)Time Complexity,O(1) Space Complexity ▶

## Question 1

Correct

Mark 1.00 out of 1.00

Find the intersection of two sorted arrays.

OR in other words,

Given 2 sorted arrays, find all the elements which occur in both the arrays.

Input Format

· The first line contains T, the number of test cases. Following T lines contain:

1. Line 1 contains N1, followed by N1 integers of the first array
2. Line 2 contains N2, followed by N2 integers of the second array

Output Format

The intersection of the arrays in a single line

Example

Input:

```
1
3 10 17 57
6 2 7 10 15 57 246
```

Output:

```
10 57
```

Input:

```
1
6 1 2 3 4 5 6
2 1 6
```

Output:

```
1 6
```

**For example:**

Input	Result
1 3 10 17 57 6 2 7 10 15 57 246	10 57

**Answer:** (penalty regime: 0 %)

```
1 #include <stdio.h>
2 int main(){
3     int t;
4     scanf("%d",&t);
5     int n;
6     scanf("%d",&n);
7     int arr[n];
8     for(int i=0;i<n;i++){
9         scanf("%d",&arr[i]);
10    }
11    int n1;
12    scanf("%d",&n1);
13    int arr1[n1];
14    for(int i=0;i<n1;i++){
15        scanf("%d",&arr1[i]);
```

```
16     }
17     for(int i=0;i<n;i++){
18         for(int j=0;j<n1;j++){
19             if(arr[i]==arr1[j]){
20                 // printf("Dfg");
21                 printf("%d ",arr[i]);
22             }
23         }
24     }
25 }
```

	Input	Expected	Got	
✓	1 3 10 17 57 6 2 7 10 15 57 246	10 57	10 57	✓
✓	1 6 1 2 3 4 5 6 2 1 6	1 6	1 6	✓

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.

◀ 3-Print Intersection of 2 sorted arrays-O(m\*n)Time Complexity,O(1) Space Complexity

Jump to...

5-Pair with Difference-O(n^2)Time Complexity,O(1) Space Complexity ▶

## Question 1

Correct

Mark 1.00 out of 1.00

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that  $A[j] - A[i] = k$ ,  $i \neq j$ .

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as  $5 - 1 = 4$

So Return 1.

For example:

Input	Result
3 1 3 5 4	1

Answer: (penalty regime: 0 %)

```

1  #include <stdio.h>
2  int main(){
3      int n;
4      scanf("%d",&n);
5      int arr[n];
6      for(int i=0;i<n;i++){
7          scanf("%d",&arr[i]);
8      }
9      int k;
10     scanf("%d",&k);
11     int key=0;
12     for(int i=0;i<n;i++){
13         for(int j=0;j<n;j++){
14             if((i!=j)&&(arr[j]-arr[i]==k)){
15                 key=1;
16                 break;
17             }
18         }
19         if(key==1){
20             printf("1");
21             break;
22         }
23     }
24     if(k==0){
25         printf("0");
26     }
27 }
```

	Input	Expected	Got	
✓	3 1 3 5 4	1	1	✓
✓	10 1 4 6 8 12 14 15 20 21 25 1	1	1	✓
✓	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✓
✓	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✓

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.

[◀ 4-Print Intersection of 2 sorted arrays- \$O\(m+n\)\$ Time Complexity, \$O\(1\)\$  Space Complexity](#)

Jump to...

[6-Pair with Difference - \$O\(n\)\$  Time Complexity, \$O\(1\)\$  Space Complexity ▶](#)



## Question 1

Correct

Mark 1.00 out of 1.00

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that  $A[j] - A[i] = k$ ,  $i \neq j$ .

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as  $5 - 1 = 4$

So Return 1.

**For example:**

Input	Result
3 1 3 5 4	1

**Answer:** (penalty regime: 0 %)

```

1  #include <stdio.h>
2  int main(){
3      int n;
4      scanf("%d",&n);
5      int arr[n];
6      for(int i=0;i<n;i++){
7          scanf("%d",&arr[i]);
8      }
9      int k;
10     scanf("%d",&k);
11     int key=0;
12     for(int i=0;i<n;i++){
13         for(int j=0;j<n;j++){
14             if((i!=j)&&(arr[j]-arr[i]==k)){
15                 key=1;
16                 break;
17             }
18         }
19         if(key==1){
20             printf("1");
21             break;
22         }
23     }
24     if(k==0){
25         printf("0");
26     }
27 }
```

	Input	Expected	Got	
✓	3 1 3 5 4	1	1	✓
✓	10 1 4 6 8 12 14 15 20 21 25 1	1	1	✓
✓	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✓
✓	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✓

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.

◀ 5-Pair with Difference-O(n^2)Time Complexity,O(1) Space Complexity

Jump to...

1-Finding Duplicates-O(n^2) Time Complexity,O(1) Space Complexity ▶