

2.1 Combinatorics Applied Structures

3) 4 shirt's sizes, 6 shirt colors, 3 emblem choices
 $\{XL, L, M, S\} * \{\text{red, blue, green, white, orange, yellow}\} * \{\text{dragon, none, dog}\}$
 $(4)(6)(3) = 72$ different combinations of shirts.

5) 6 officers, 2 rows, 3 people different arrangements
 $6! = 6 * 5 * 4 * 3 * 2 * 1 = 720$

7) 2 blouses, 2 ^{3 bottoms} pants, 1 skirt, 1 blazer

$2 * 3 * 1 = 6$ different outfits

Blazer optional (yes, ²no)

$2 * 3 * 2 = 12$ different outfits

1 sweater $2 * 3 * 2 * 1 = 12$ different outfits

Sweater optional (yes, ²no)

$2 * 3 * 2 * 2 = 24$ different outfits

9) a) 2 choices {0, 1} 8 bits/pack

$2^8 = 256$ different bit patterns

b) first 4 = last 4

$2^4 = 16$

c) half even \rightarrow half odd

$\frac{2^8}{2} = 128$

11) a) $A = \{1, 2, 3, 4\}$ $P(A) = 2^4 = 16$ ^{different subsets}

b) $A = \{1, 2, 3, 4, 5\}$ proper subsets of A $2^5 - 1 = 31$

$$13) \begin{matrix} 1 \\ A & B & C \\ 1 & 2 & 3 \end{matrix} \quad \begin{matrix} 2 \\ C & A & B \\ 1 & 2 & 3 \end{matrix} \quad \begin{matrix} 3 \\ B & C & A \\ 1 & 2 & 3 \end{matrix} \quad 3 \text{ different ways}$$

$$15) \{fish, lamb, beef\}, \{peas, carrots\}, \{pie, ice cream, cake\}$$

$$3 * 2 * 3 = 18 \text{ different dinners}$$

2.2

$$1) 1000 \cdot 999 \cdot 998 = 997,002,000 \text{ different ways to distribute}$$

$$\text{or } P(1000, 3) = \frac{1000!}{(1000-3)!} = \frac{1000 \cdot 999 \cdot 998 \cdot \cancel{997!}}{\cancel{997!}} = 997,002,000$$

$$5) 15! = 15 \cdot 14 \cdot \dots \cdot 1 = 1,307,674,400,000$$

$$\text{or } P(15, 15) = \frac{15!}{(15-15)!} = \frac{15!}{1}$$

$$7) a) P(15, 5) = \frac{15!}{(15-5)!} = \frac{15 \cdot 14 \cdot 13 \cdot 12 \cdot 11 \cdot \cancel{10!}}{\cancel{10!}} = 360,360$$

one spot second spot

$$b) 15 \cdot 14 \cdot 13 \cdot 12 \cdot 11$$

two players
for one spot

$$\rightarrow 2 \cdot 14 \cdot 13 \cdot 12 \cdot 11 = 48,048$$

2.3

$$1) A = \{a, b, c\} \text{ Partition}$$

$$\{\{a\}, \{b\}, \{c\}\}, \{\{a, b\}, \{c\}\}, \{\{a\}, \{b, c\}\}, \{\{a, c\}, \{b\}\},$$

$$\{\{a, b, c\}\}$$

2.4

1) 3 faculty 4 students 10 faculty 25 students

$$\binom{10}{3} \cdot \binom{25}{4} = \frac{10!}{(10-3)!3!} \cdot \frac{25!}{(25-4)!4!}$$

$$= \frac{10!}{7!3!} \cdot \frac{25!}{21!4!} = \frac{10 \cdot 9 \cdot 8 \cdot 7}{7!3!} \cdot \frac{25 \cdot 24 \cdot 23 \cdot 2 \cdot 21}{21!4!}$$

$$= \frac{720}{6} \cdot \frac{303600}{24} = 1,518,000$$

3) $\{1, 2, 3, \dots, 10\}$ How many subsets contain 7 elements

$$\binom{10}{7} + \binom{10}{8} + \binom{10}{9} + \binom{10}{10} = \frac{10!}{(10-7)!7!} + \frac{10!}{(10-8)!8!} + \frac{10!}{(10-9)!9!} + \frac{10!}{(10-10)!10!}$$

$$= \frac{10 \cdot 9 \cdot 8 \cdot 7}{3!2!} + \frac{10 \cdot 9 \cdot 8}{2!1!} + \frac{10 \cdot 9}{1!1!} + \frac{10!}{10!} = \frac{720}{6} + \frac{90}{2} + 10 + 1 = 176$$

7) 52 cards each player gets 5 cards

a) How many hands possible?

combination of 5 cards are chose from 52 cards

$$\binom{52}{5} = \frac{52!}{(52-5)!5!} = \frac{52 \cdot 51 \cdot 50 \cdot 49 \cdot 48 \cdot 47}{47! \cdot 5!} = 2,598,960$$

b) 1st player 2nd 3rd 4th

$$\binom{52}{5} \cdot \binom{47}{5} \cdot \binom{42}{5} \cdot \binom{37}{5} = 2,598,960 \cdot \frac{47!}{(47-5)!5!} \cdot \frac{42!}{(42-5)!5!} \cdot \frac{37!}{(37-5)!5!}$$

$$= 2,598,960 + 1,533,939 + 850,668 + 435,897 = 5,419,464$$

4 aces in
 9) 5 hand cards 52 cards 2 aces in 5 hand card

$$\binom{4}{2} \cdot \binom{48}{3} = \frac{4!}{(4-2)!2!} \cdot \frac{48!}{(48-3)!3!} = 6 \cdot \frac{103776}{6} = 103776$$

11) $\binom{12}{5} \cdot \binom{7}{4} \cdot \binom{3}{3}$

group of 5 from 12 group of 4 from 7 group of 3 from 3

$$\frac{12!}{(12-5)!5!} \cdot \frac{7!}{(7-4)!4!} \cdot \frac{3!}{(3-3)!3!} = \frac{12 \cdot 11 \cdot 10 \cdot 9 \cdot 8 \cdot 7}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5} \cdot \frac{7 \cdot 6 \cdot 5 \cdot 4}{3 \cdot 4} \cdot \frac{3}{1} = 792 \cdot 35 \cdot 1 = 27720$$

Book of Proof

3.2

1) (T, H, E, O, R, Y) 6 Elements repetition allowed
 a) length 4 list How many?

$$6 \cdot 6 \cdot 6 \cdot 6 = 1296$$

b) length 4 list begin with T

$$1 \cdot 6 \cdot 6 \cdot 6 = 216$$

c) length 4 list don't begin with T

$$5 \cdot 6 \cdot 6 \cdot 6 = 1080$$

3) (A, B, C, D, E, F) length 3 list rep allowed from 6

$$6 \cdot 6 \cdot 6 = 216$$

b) rep not allowed $6 \cdot 5 \cdot 4 = 120$

c) rep not allowed contain A $(A \cdot 5 \cdot 4) + (5 \cdot A \cdot 4) + (5 \cdot 4 \cdot A) = 60$

d) rep allowed contain A $6 \cdot 6 \cdot 6 - 5 \cdot 5 \cdot 5 = 91$

8 bits - 2 choices

5) a) $2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 = 2^8 = 256$ reps allowed

b) $2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 1 = 2^7 = 128$ end in zero

c) $2 \cdot 1 \cdot 2 \cdot 1 \cdot 2 \cdot 2 \cdot 2 \cdot 2 = 2^6 = 64$ second and fourth = 1

d) $2^6 + 2^6 + 2^6 = 192$

$(n \cdot 1 \cdot n \cdot 0 \cdot n \cdot n \cdot n \cdot n) (n \cdot 0 \cdot n \cdot 1 \cdot n \cdot n \cdot n \cdot n) (n \cdot 1 \cdot n \cdot 1 \cdot n \cdot n \cdot n \cdot n)$

7) 4 letter codes from $\overbrace{ABCD \dots}^{26}$

$26^4 = 456,976$

4 letter codes no 2 consecutive letters

$26 \cdot 25 \cdot 25 \cdot 25 = 406,250$

any $\hat{\leftarrow}$ can't be $\hat{\leftarrow}$ can't be $\hat{\leftarrow}$ can't be

9) 5 colors \cdot 3 engines \cdot 2 transmissions = 30 different combinations

3.3

1) 52 cards Five hands 1 red card in five hands

$52 \cdot 51 \cdot 50 \cdot 49 \cdot 48 = 311,875,200$ Possible 5 hand combinations

26 red cards 26 black cards

$26 \cdot 25 \cdot 24 \cdot 23 \cdot 22 = 7,893,600$ black cards

$311,875,200 - 7,893,600 = \boxed{303,981,600}$

All black or All Hearts

$26 \cdot 25 \cdot 24 \cdot 23 \cdot 22 + 13 \cdot 12 \cdot 11 \cdot 10 \cdot 9 = \boxed{8,048,040}$

All black

All red

$$3) (26 \cdot 25 \cdot 24 \cdot 23 \cdot 22) + (26 \cdot 25 \cdot 24 \cdot 23 \cdot 22) = 15,787,200$$

$$5) 1 - 9999 \text{ no rep's}$$

$$\begin{array}{ccccccc} \text{one digit \#s} & & 2 \text{ digit \#s} & & 3 \text{ digit \#s} & & 4 \text{ digit \#s} \\ 9 & + & 9 \cdot 9 & + & 9 \cdot 9 \cdot 8 & + & 9 \cdot 9 \cdot 8 \cdot 7 \end{array}$$

$$= 5274$$

One repeated digit

$$9999 - 5274 = 4725$$

total # of integers total # of no repeated

$$7) \text{ five long } 26 \text{ letters } 1 \text{ upper case}$$

upper case and lower case

$$52 \cdot 52 \cdot 52 \cdot 52 \cdot 52 = 380,204,032$$

all lower case

$$26 \cdot 26 \cdot 26 \cdot 26 \cdot 26 = 11,881,376$$

$$380,204,032 - 11,881,376 = \boxed{368,322,656}$$

all combinations - all upper/lower case combinations

must mix

$$52^5 - 26^5 - 26^5 = \boxed{356,441,280}$$

$$9) \text{ list of 6 no rep's A, B, C, D, E, F, G, H}$$

$$AE**** + *AE*** + **AE** + ***AE*$$

$$+ ****AE = (6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1) 5 = 3600$$

11) integers divisible by 5: 1-1000

$$1000/5 = 200 \text{ integers} \quad 1000 - 200 = 800 \text{ not divisible by 5}$$

3.4

1) smallest n for $n!$ has 10 digits

$$13! = 13 \cdot 12 \cdot \dots \cdot 1 = 6,227,020,800$$

3) 5 digit positive, odd integers no rep

$$(1, 3, 5, 7, 9) \quad 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 5! = 120$$

$$5) \frac{120!}{118!} = \frac{120 \cdot 119 \cdot 118!}{118!} = 120 \cdot 119 = 14,280$$

7) How many 9 digit #'s can be made: 1-9

all odd start on the left no rep.

odd even
1 3 5 7 9 2 4 6 8

$$5!4! = (5 \cdot 4 \cdot 3 \cdot 2 \cdot 1)(4 \cdot 3 \cdot 2 \cdot 1) = 2880$$

9) ABCDEFG permutations? ABC consecutively and

$$\text{order } \boxed{ABC}^{*} * * * * = 5! = 120$$

11) 7 hands 52 cards How many are not all red cards

$$P(52, 7) = \frac{52!}{(52-7)!} = 674,274,182,400$$

$$P(26, 7) = \frac{26!}{(26-7)!} = 3,315,312,000$$

$$P(52, 7) - P(26, 7) = 670,958,870,400$$

13) 6 list no rep 26 letters

$$P(26, 6) = \frac{26!}{(26-6)!} = \frac{26 \cdot 25 \cdot 24 \cdot 23 \cdot 22 \cdot 21 \cdot \cancel{20!}}{\cancel{20!}} = 165,765,600$$

15) 15 people 4 list different ways

$$P(15, 4) = \frac{15!}{(15-4)!} = \frac{15 \cdot 14 \cdot 13 \cdot 12 \cdot \cancel{11!}}{\cancel{11!}} = 32,760$$

17) 10 people 3 list different ways

$$P(10, 3) = \frac{10!}{(10-3)!} = \frac{10 \cdot 9 \cdot 8 \cdot \cancel{7!}}{\cancel{7!}} = 720$$

3.5

1) A has 37 elements How many subsets have 10 elements

$$\binom{37}{10} = \frac{37!}{(37-10)!10!} = \frac{37 \cdot 36 \cdot \dots \cdot 27!}{\cancel{27!}10!} = 348,330,136$$

subsets 30 elements

$$\binom{37}{30} = \frac{37 \cdot 36 \cdot \dots \cdot 30!}{(37-30)!30!} = \frac{37 \cdot 36 \cdot 35 \cdot 34 \cdot 33 \cdot 32 \cdot 31 \cdot \cancel{30!}}{\cancel{30!}} = 10,295,472$$

subsets 0 elements

$$\binom{37}{0} = \frac{37!}{(37-0)!0!} = \frac{37!}{37! \cdot 1} = 1$$

3) 56 subsets with 3 elements $|X| = ?$

$$\binom{z}{3} = 56 \quad \frac{z!}{(z-3)!3!} = \frac{z \cdot (z-1) \cdot (z-2) \cdot \cancel{(z-3)!}}{\cancel{(z-3)!}3!} = 20$$

$$\frac{8!}{(8-3)!3!} = \frac{8 \cdot 7 \cdot 6 \cdot \cancel{5!}}{\cancel{5!}3!} = 56 \quad z=8 \quad \binom{8}{3} = 56 \quad |X|=8$$

5) 16 digit choose seven 1's

$$\binom{16}{7} = \frac{16!}{(16-7)!7!} = \frac{16 \cdot 15 \cdot \dots \cdot 9!}{9! \cdot 7!} = 11,440$$

7) $|\{X \in \mathcal{P}(\{0,1,2,3,4,\dots,9\}) : |X| < 4\}|$

$$= \binom{10}{0} + \binom{10}{1} + \binom{10}{2} + \binom{10}{3} = \frac{10!}{(10-0)!0!} + \frac{10!}{(10-1)!1!} + \frac{10!}{(10-2)!2!} + \frac{10!}{(10-3)!3!}$$

$$= 1 + 10 + 45 + 120 = 176$$

9) 6 list ABCDEF no rep, D occurs before A

DA**** $\binom{6}{2} \cdot 4! = \frac{6!}{(6-2)!2!} \cdot (4 \cdot 3 \cdot 2 \cdot 1) = 360$

11) 10 digits integers no 0's three 6's

$$\binom{10}{3} \cdot 8 \cdot 8 \cdot 8 \cdot 8 \cdot 8 \cdot 8 \cdot 8 = \frac{10!}{(10-3)!3!} \cdot 8^7 = 251,658,240$$

13) $n, k \in \mathbb{Z} \quad 0 \leq k \leq n \quad \binom{n}{k} = \binom{n}{n-k}$

$$\binom{n}{k} = \frac{n!}{(n-k)!k!} \quad \binom{n}{n-k} = \frac{n!}{(n-(n-k))!(n-k)!}$$

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} = \frac{n!}{(n-(n-k))!(n-k)!} = \binom{n}{n-k}$$

15) 10 digits no #1, four's

total 10 digits binaries

$$2^{10} = 1024$$

total of 10 digits with four 1's

$$\binom{10}{4} = \frac{10!}{(10-4)!4!} = 210$$

$$1024 - 210 = 814$$

17) 10 digit binary (four 1's or five 1's) or (not have)

$$\text{have } \binom{10}{4} + \binom{10}{5} = \frac{10!}{(10-4)!4!} + \frac{10!}{(10-5)!5!} = 210 + 252 = 462$$

$$\text{not have } 2^{10} - \binom{10}{4} - \binom{10}{5} = 1024 - 462 = 562$$

19) 5 hand cards 52 deck 4 suit each 13 cards

$$\binom{13}{5} + \binom{13}{5} + \binom{13}{5} + \binom{13}{5} = (1287)4 = 5148$$

hearts aces

3.6

1) Row 11 Pascal's triangle

$$(x+y)^{11} = \binom{11}{0} \binom{11}{1} \binom{11}{2} \binom{11}{3} \binom{11}{4} \dots \binom{11}{11}$$

$$= 1, 11, 55, 165, 330, 462, 462, 330, 165, 55, 11, 1$$

3) coefficient x^8 in $(x+2)^{13}$

$$(x+y)^{13} \rightarrow x^8 y^5 = \binom{13}{5} x^8 y^5 = 1287 x^8 y^5$$

$$y=2 \rightarrow 1287 x^8 (2)^5 = 41,184 x^8$$

$$5) \sum_{k=0}^n \binom{n}{k} = 2^n \quad 2^n = (1+1)^n$$

$$= \binom{n}{0} 1^n 1^0 + \binom{n}{1} 1^{n-1} 1^1 + \dots + \binom{n}{n-1} 1^1 1^{n-1} + \binom{n}{n} 1^0 1^n$$

$$\Rightarrow \sum_{k=0}^n \binom{n}{k} = 2^n$$

$$\Rightarrow \binom{n}{0} + \binom{n}{1} + \binom{n}{2} + \dots + \binom{n}{n}$$

$$\begin{aligned}
 7) \sum_{k=0}^n 3^k \binom{n}{k} &= 4^n & 4^n &= (1+3)^n \\
 &= \binom{n}{0} 1^n 3^0 + \binom{n}{1} 1^{n-1} 3^1 + \dots + \binom{n}{n} 1^0 3^n \\
 &= \sum_{k=0}^n 3^k \binom{n}{k}
 \end{aligned}$$

$$9) \binom{n}{0} - \binom{n}{1} + \binom{n}{2} - \binom{n}{3} + \dots \pm \binom{n}{n} = 0$$

$$0^n = (1 + (-1))^n$$

$$= \binom{n}{0} 1^n (-1)^0 + \binom{n}{1} 1^{n-1} (-1)^1 + \dots + \binom{n}{n} 1^0 (-1)^n$$

$$= \binom{n}{0} - \binom{n}{1} + \binom{n}{2} - \dots \pm \binom{n}{n} = 0$$

$$11) 9^n = \sum_{k=0}^n (-1)^k \binom{n}{k} (10^{n-k})$$

$$9^n = (10 + (-1))^n$$

$$= \binom{n}{0} 10^n (-1)^0 + \binom{n}{1} 10^{n-1} (-1)^1 + \dots + \binom{n}{n} 10^0 (-1)^n$$

$$3) \binom{n}{3} = \binom{2}{2} + \binom{3}{2} + \binom{4}{2} + \binom{5}{2} + \dots + \binom{n-1}{2}$$

$$n \geq 3 \quad \binom{n}{3} = \binom{n-1}{3} + \binom{n-1}{2} = \binom{n-2}{3} + \binom{n-2}{2} + \binom{n-1}{2} = \binom{n-3}{3} + \binom{n-3}{2} + \binom{n-2}{2} + \binom{n-1}{2}$$

$$= \dots = \binom{2}{2} + \binom{3}{2} + \binom{4}{2} + \dots + \binom{n-1}{2}$$

3.7

$$1) |A| = 100 \text{ senior mm} \quad |B| = \text{senior nm} \quad |A \cap B| = 33 \text{ senior bmm}$$

$$|A \cup B| = 523 \text{ seniors}$$

$$|A \cup B| = |A| + |B| - |A \cap B|$$

$$523 = 100 + |B| - 33$$

$$|B| = 523 + 33 - 100 = 456$$

3) 4 digit positive integers even or zeros

A = even B = no zeros

0 0 0 0
↑ ↑ ↑ ↑

1 to 9 0-9 0-9 0,2,4,6,8

$$|A| = 9 \cdot 10 \cdot 10 \cdot 5 = 4500$$

0 0 0 0
↑ ↑ ↑ ↑

1 to 9 1 to 9 1-9 1,3,5,7,9

$$|B| = 9 \cdot 9 \cdot 9 \cdot 5 = 3645$$

$$|A \cup B| = |A| + |B| = 8145$$

5) 7 digits begin 1 or end 1 or four 1's

$$2^6 + 2^6(4) - 2^5 - \binom{6}{3} - \binom{6}{3} + \binom{5}{2} = 101$$

$$|A \cup B \cup C| = |A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C|$$

7) 4 hands 52 card deck

all 4 same suit or all red.

$$|A| = \binom{13}{4} 4 \quad |B| = \binom{26}{4} \quad |A \cap B| = 2 \binom{13}{4}$$

$$|A \cup B| = |A| + |B| - |A \cap B| = 4 \binom{13}{4} + \binom{26}{4} - 2 \binom{13}{4} = 16,380$$

9) 4 list LISTED rep allowed all vowels, ends D

$$|A| = 2 \cdot 2 \cdot 6 \cdot 6 = 144 \quad |B| = 6 \cdot 6 \cdot 6 \cdot 1 = 216$$

$$|A \cap B| = 2 \cdot 2 \cdot 6 \cdot 1 = 24$$

$$|A \cup B| = 144 + 216 - 24 = 336$$

11) 7 digits even or three 0's

$$|A| = 9 \cdot 10 \cdot 10 \cdot 10 \cdot 10 \cdot 10 \cdot 5 = 4,500,000$$

$$|B| = 9 \cdot \binom{6}{3} \cdot 9 \cdot 9 \cdot 9 = 131,220$$

$$|A \cap B| = 9 \cdot \binom{5}{3} \cdot 9 \cdot 9 \cdot 9 + 9 \cdot \binom{5}{2} \cdot 9 \cdot 9 \cdot 9$$

$$|A \cup B| = 4,500,000 + 131,220 - 29160 = 4,569,560$$

13) 8 digits end 1 or four 1's

$$|A| = 2^7 = 128 \quad |B| = \binom{8}{4} = 70$$

$$|A \cap B| = \binom{7}{4} = 35$$

$$|A \cup B| = 128 + 70 - 35 = 163$$

15) 10 digit begin in 1 or end 1

$$|A| = 2^9$$

$$|B| = 2^9$$

$$|A \cap B| = 2^8$$

$$|A \cup B| = 2^9 + 2^9 - 2^8 = 768$$