INTERVIEW NOTES — BRENDA'S POULTRY FARM

PCSE211 Group Project: Egg Production Tracking System

Organization	Brenda's Poultry Farm
Location	Muyenga, Kampala, Uganda
Date and method	19 October 2026 — Questionnaire
Respondent	Farm worker (name not recorded)

Note: This draft is based on information provided by the project group. Please check that the details match what the worker said before submitting it.

1. Purpose of the questionnaire

The questionnaire was used to learn how the farm keeps egg production records, what problems workers face when using the records, and what information the farm needs. The answers will help the group plan a simple Egg Production Tracking System.

2. Questions and responses

Question	Response
1. What does the farm mainly do?	The farm keeps chickens for egg production.
2. About how many laying hens does the farm have?	The farm has about 70 laying hens.
3. How many eggs does the farm collect?	The farm collects about 50 to 60 eggs a day. This is around 350 to 420 eggs in a week.
4. Who collects and records the eggs?	Farm workers collect the eggs and record the production.
5. How do you record the eggs collected each day?	The farm uses a large counter book meant for recording egg production.
6. What is the main problem with the current record book?	Workers have to calculate the totals by hand. Sometimes they make mistakes when adding up the figures.
7. How do these mistakes affect the farm?	When figures are added wrongly, it becomes hard to know the correct production total and compare production over time.
8. Have you had problems with missing or wrong records?	The problem reported for this project is mistakes during manual calculations. Whether records have also gone missing should be checked if needed.
9. How do you work out weekly or monthly production?	Production figures are taken from the counter book and added by hand, so calculation mistakes can happen.
10. What happens to eggs that are cracked or damaged?	If an egg is cracked but the inner membrane is still intact, the family may eat it at home or sell it cheaply to local people for immediate use. If the shell is badly cracked, the egg is thrown away.
11. What reports would be useful to the farm?	Daily and weekly egg production reports are useful. Flock health, sales and inventory reports were also mentioned, but this project will focus on egg production records.
12. Would a simple computerised system help? Why?	A system could add up production figures automatically and make daily and weekly totals easier to check. This benefit should be confirmed with the worker.
13. What should the new system be able to do?	It should record the number of eggs collected each day and show daily and weekly totals. It could also record damaged eggs separately if needed.

14. Is there anything else we should know about the current process?	The main point for the project is that production is written in a counter book and totals are calculated by hand. Any other details should be checked with the farm worker.

3. Main findings

- The farm has about 70 laying hens and collects around 50–60 eggs per day, or about 350–420 eggs per week.
- Egg production is recorded in a large counter book.
- Workers add up the figures by hand. Mistakes can make it hard to track production correctly over time.
- Cracked eggs with the inner membrane still intact may be eaten at home or sold cheaply for immediate use. Eggs with badly cracked shells are thrown away.

4. Proposed system and scope

The proposed system will focus on egg production records only. It should let a worker enter the number of eggs collected each day and calculate daily and weekly totals automatically. A separate count of damaged eggs can be included if the farm needs it. Sales management, general inventory and flock health will not be separate functions because the project is meant to solve one main problem.

5. Information confirmed for the project

- About 70 laying hens are kept, with around 50–60 eggs collected daily and 350–420 weekly.
- One worker records production in the counter book, and Brenda checks the entries and confirms the totals.
- Damaged eggs are counted separately. Eggs with an intact inner membrane may be eaten at home or sold cheaply for immediate use; badly cracked eggs are thrown away.