

EE 5323- Take Home Exam 2

Fall 2021

This exam has 6 pages in all. There are 4 problems.

.....

Almost all questions require numerical calculations to arrive at the answers. To obtain full credit, show all your work. No partial credit will be given without the supporting work.

.....

Name: Bardha Maja

Pledge of honor:

"On my honor I have neither given nor received aid on this examination."

Signature: [Signature]

# 1. Lyapunov Function

Use Lyapunov function to examine the stability of the following systems. Be clear and show all steps.

a.

$$\dot{x}_1 = x_2 \sin x_1 - x_1$$

$$\dot{x}_2 = -x_1 \sin x_1 - x_2$$

$$V(x_1, x_2) = \frac{1}{2} (x_1^2 + x_2^2) > 0$$

$$V' = x_1 \dot{x}_1 + x_2 \dot{x}_2 = 0 \Rightarrow$$

$$x_1 (x_2 \sin x_1 - x_1) + x_2 (-x_1 \sin x_1 - x_2) = 0 \Rightarrow$$

$$x_1 x_2 \sin x_1 - x_1^2 - x_1 x_2 \sin x_1 - x_2^2 = 0 \Rightarrow -(x_1^2 + x_2^2) = 0$$

eq (0,0) only

$$V' \leq 0 \text{ then } \underline{\text{GAS}}$$

b.

$$\dot{x}_1 = x_2 \sin x_1 - x_1$$

$$\dot{x}_2 = -x_1 \sin x_1$$

$$V' = x_1 \dot{x}_1 + x_2 \dot{x}_2 = 0 \Rightarrow$$

$$x_1 (x_2 \sin x_1 - x_1) + x_2 (-x_1 \sin x_1) = 0$$

$$x_1 x_2 \sin x_1 - x_1^2 - x_1 x_2 \sin x_1 = 0$$

$$V' = -x_1^2 ; \quad V' = 0, x_1 = 0$$

$$V' \leq 0 \checkmark$$

SIPL

marginally stable  
because it only depends  
on  $x_1$

## 2. LaSalle's Extension

$$\ddot{x} + k_1 \dot{x} + k_2 \dot{x}^3 + k_3 x^5 = 0 \Rightarrow \ddot{x} = -k_1 \dot{x} - k_2 \dot{x}^3 - k_3 x^5$$

- a. Use Lyapunov to check the stability. Hint: Use the energy as the Lyapunov function.  
Take the potential energy as

$$PE = \int_0^x (k_2 \dot{x}^3 + k_3 x^5) dx \quad V = \frac{1}{2} \dot{x}^2 + \int_0^x C(y) dy = K + U;$$

$$V = \frac{1}{2} \dot{x}^2 + \int_0^x C(y) dy \Rightarrow \text{Per Leibniz's theorem: } \frac{d}{dt} \int_{\alpha}^{\beta} F(x,t) dx = \int_{\alpha}^{\beta} \frac{\partial F(x,t)}{\partial t} dx = \beta \dot{F}(\beta,t) - \alpha \dot{F}(\alpha,t) + \int_{\alpha}^{\beta} \frac{\partial F(x,t)}{\partial t} dx$$

$$\Rightarrow \dot{V} = \dot{x}(-k_1 \dot{x} - k_2 \dot{x}^3 - k_3 x^5) + \dot{x}(k_2 \dot{x}^2 + k_3 x^4) - (0)(1) + (0+0) \Big|_{\alpha=0}^{\beta=x}$$

$$\Rightarrow \dot{V} = -k_1 \dot{x}^2 \leq 0 \text{ if } k_1 \geq 0,$$

$$\dot{V} = 0 \Rightarrow \dot{x} \rightarrow 0 \text{ or } k_1 = 0 \quad \underline{\text{DISH}}$$

↳ Hints at use of LaSalle's extension  
we plug this in dynamics equation.

- b. Use LaSalle's extension to find a stronger type of stability for the system.

if  $\dot{x} \rightarrow 0 \Rightarrow \ddot{x} \rightarrow 0$ ; plug in sys dynamic

$$\begin{matrix} \ddot{x} & + & k_1 \dot{x} & + & k_2 \dot{x}^3 & + & k_3 x^5 & = & 0 \\ 0 & & 0 & & 0 & & & & \end{matrix} \Rightarrow k_3 x^5 = 0 \Rightarrow \underline{x=0}$$

$\Rightarrow$  thus the system is dissipative  
& reaches equilibrium at  $x=0$ ; thus it is  
Considered Global Asymptotic Stable (GAS)

### 3. Lyapunov Equation for Linear Systems

Use Lyapunov Equation to check the stability of the linear systems

a.  $\dot{x} = Ax = \begin{bmatrix} 0 & 1 \\ 0 & -6 \end{bmatrix} x$

*not stable  
unstable*

$$A^T P + P A = -Q \Rightarrow \begin{bmatrix} a_1 & a_3 \\ a_2 & a_4 \end{bmatrix} \begin{bmatrix} p_1 & p_2 \\ p_3 & p_4 \end{bmatrix} + \begin{bmatrix} p_1 & p_2 \\ p_3 & p_4 \end{bmatrix} \begin{bmatrix} a_1 & a_3 \\ a_2 & a_4 \end{bmatrix} = -Q$$

$$\begin{bmatrix} a_1 p_1 + a_3 p_3 + a_1 p_1 + a_3 p_3 & a_1 p_2 + a_3 p_4 + a_2 p_1 + a_4 p_3 \\ a_2 p_1 + a_4 p_3 + a_1 p_3 + a_3 p_1 & a_2 p_2 + a_4 p_4 + a_2 p_2 + a_4 p_4 \end{bmatrix} = -Q,$$

where I test the following  
Q matrices:

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \dots$$

I a unique solution  
DNE

b.  $\dot{x} = Ax = \begin{bmatrix} 0 & 1 \\ -3 & -4 \end{bmatrix} x$

stable w/

$$Q = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$m_{11} = .83$$

$$m_{22} = .1667$$

$$\begin{bmatrix} .83 & -.5 \\ -.5 & .5 \end{bmatrix}$$

$\left[ \begin{array}{c} \text{Same} \\ \text{a.p.g. matrix} \\ \text{from 3.a.} \end{array} \right] = -Q$

$$m_{22} = (.83 \cdot .5) - (.25)$$

#### 4. UUB

Use Lyapunov to show that the system

$$\dot{x}_1 = x_1 x_2^2 - x_1 (x_1^2 + x_2^2 - 3)$$

$$\dot{x}_2 = -x_1^2 x_2 - x_2 (x_1^2 + x_2^2 - 3)$$

is uniformly ultimately bounded UUB. That is, show that the Lyapunov derivative is **NEGATIVE OUTSIDE A BOUNDED REGION**. Find the radius of the bounded region outside which  $\dot{V} < 0$ . Any states outside this region are attracted towards the origin.

$$V(x_1, x_2) = \frac{1}{2} (x_1^2 + x_2^2 - 3)^2 > 0 \Rightarrow \dot{V} = \frac{d}{dt} (x_1^2 + x_2^2 - 3) (2x_1 \dot{x}_1 + 2x_2 \dot{x}_2)$$

$$\begin{aligned} \Rightarrow \dot{V} &= (x_1^2 + x_2^2 - 3) \left( 2x_1 \left( x_1 x_2^2 - x_1 (x_1^2 + x_2^2 - 3) \right) + 2x_2 \left( -x_1^2 x_2 - x_2 (x_1^2 + x_2^2 - 3) \right) \right) \\ &= (x_1^2 + x_2^2 - 3) \left( \underbrace{2x_1 (-x_1^3 - x_1 x_2^2 + 3x_1)}_{2x_1(-x_1^3 - x_1 x_2^2 + 3x_1)} + \underbrace{2x_2 (-x_1^2 x_2 - x_2^3 + 3x_2)}_{2x_2(-x_1^2 x_2 - x_2^3 + 3x_2)} \right) \\ &= (x_1^2 + x_2^2 - 3) \left( -2x_1^2 (x_1^2 + x_2^2 - 3) - 2x_2^2 (x_1^2 + x_2^2 - 3) \right) \end{aligned}$$

$$\Rightarrow \dot{V} = \underbrace{(x_1^2 + x_2^2 - 3)^2}_{>0} \underbrace{(-2x_1^2 - 2x_2^2)}_{\leq 0} = 0$$

$$\Rightarrow \|\dot{V}\| \leq \underbrace{\|x_1^2 + x_2^2 - 3\|}_{\downarrow \text{creates boundary}} \underbrace{\|(-2)\|}_{\times} \underbrace{\|x_1^2 + x_2^2\|}_{\geq 0}$$

Per Cauchy Schwarz  
 $\|x \cdot y\| \leq \|x\| \cdot \|y\|$   
 given an orthonormal space

where  
the radius  
is  $\sqrt{3}$

