

The Modified General Version of the Master Theorem

Douglas R. Stinson
David R. Cheriton School of Computer Science
University of Waterloo
Waterloo, Ontario, N2L 3G1, Canada

September 28, 2015

Our goal is to prove the following generalization of the Master Theorem stated on slide 60. This is similar to the version given in [CLRS], except that the condition in the third case has been replaced by a simpler condition¹.

Theorem 1. *Suppose that $a \geq 1$ and $b > 1$. Denote $x = \log_b a$. Suppose $T(n)$ is defined by the recurrence*

$$T(n) = \begin{cases} a T\left(\frac{n}{b}\right) + f(n) & \text{if } n > 1 \\ d & \text{if } n = 1. \end{cases}$$

Then, for $n = b^j$, it holds that

$$T(n) \in \begin{cases} \Theta(n^x) & \text{if } f(n) \in O(n^{x-\epsilon}) \text{ for some } \epsilon > 0 \\ \Theta(n^x \log n) & \text{if } f(n) \in \Theta(n^x) \\ \Theta(f(n)) & \text{if } f(n)/n^{x+\epsilon} \text{ is an increasing function of } n \\ & \text{for some } \epsilon > 0. \end{cases}$$

We begin by using the recursion tree method to obtain a formula for $T(n)$. This is very similar to what was done in class. The resulting formula is:

$$T(n) = a^j f(1) + \sum_{i=0}^{j-1} a^i f(n/b^i) = d n^x + \sum_{i=0}^{j-1} a^i f(n/b^i), \quad (1)$$

where we use the fact that $a^j = (b^x)^j = (b^j)^x = n^x$.

¹This condition was pointed out to me by Timothy Chan.

case 1

Here we have $f(n) \in O(n^{x-\epsilon})$ for some $\epsilon > 0$. Denote $y = x - \epsilon$; then $f(n) \in O(n^y)$. Note that $x - y = \epsilon > 0$, so $b^{x-y} > 1$.

First, from (1), we have $T(n) \geq d n^x$, so $T(n) \in \Omega(n^x)$.

For the upper bound, we use the fact that $f(n) \in O(n^y)$, which implies that there is a constant $c > 0$ such that $f(n) \leq c n^y$ for all $n > 1$. Now, from (1), we have

$$\begin{aligned}
T(n) &\leq d n^x + \sum_{i=0}^{j-1} c a^i (n/b^i)^y \\
&= d n^x + c n^y \sum_{i=0}^{j-1} (a/b^y)^i \\
&= d n^x + c n^y \sum_{i=0}^{j-1} (b^{x-y})^i \\
&\in O(n^x + n^y (b^{x-y})^j), \text{ since } \sum_{i=0}^{j-1} (b^{x-y})^i \in O((b^{x-y})^j) \text{ (slide 42)} \\
&= O(n^x + n^y (b^j)^{x-y}) \\
&= O(n^x + n^y n^{x-y}) \\
&= O(n^x).
\end{aligned}$$

Because $T(n) \in \Omega(n^x)$ and $T(n) \in O(n^x)$, we have that $T(n) \in \Theta(n^x)$.

case 2

In this case, we have that $f(n) \in \Theta(n^x)$. The proof of this case is the same as the proof of the corresponding case of the simplified version of the Master Theorem.

case 3

Here, $f(n)/n^{x+\epsilon}$ is an increasing function of n . Denote $y = x + \epsilon$. Then $x - y = \epsilon < 0$, so $b^{x-y} < 1$. Note also that $n^y \in O(f(n))$.

First, from (1), we have $T(n) \geq f(n)$, so $T(n) \in \Omega(f(n))$.

For the upper bound, we use the fact that $f(n)/n^y$ is an increasing function of n . Therefore,

$$\frac{f(n)}{n^y} > \frac{f\left(\frac{n}{b}\right)}{\left(\frac{n}{b}\right)^y} > \frac{f\left(\frac{n}{b^2}\right)}{\left(\frac{n}{b^2}\right)^y} > \dots,$$

which implies that

$$f(n) > b^y f\left(\frac{n}{b}\right) > b^{2y} f\left(\frac{n}{b^2}\right) > \dots > b^{(j-1)y} f\left(\frac{n}{b^{j-1}}\right). \quad (2)$$

Now, from (1) and (2), we obtain

$$\begin{aligned}
T(n) &= d n^x + \sum_{i=0}^{j-1} a^i f(n/b^i) \\
&< d n^x + \sum_{i=0}^{j-1} (a/b^y)^i f(n) \\
&= d n^x + f(n) \sum_{i=0}^{j-1} (a/b^y)^i \\
&\in O(n^x + f(n)), \text{ since } \sum_{i=0}^{j-1} (a/b^y)^i \in O(1) \text{ (slide 42)} \\
&= O(f(n)),
\end{aligned}$$

since $x < y$ and $n^y \in O(f(n))$.