Non-steroidal anti-inflammatory Drugs and Organ Damage

May / 2023

Abstract

Non-steroidal mitigating drugs (NSAIDs) are ordinarily utilized prescriptions to ease agony and aggravation. Regardless of their boundless use, they are known to make different unfriendly impacts, including gastrointestinal, cardiovascular, and renal harmfulness. These aftereffects can prompt huge grimness and mortality, making it fundamental to assess the security of these medications, especially with long haul use.

A few examinations have recommended a relationship between NSAID use and organ harm. Gastrointestinal difficulties like dying, ulcers, and holes have been indisputable and are the most normally detailed aftereffects related with NSAID use. Essentially, cardiovascular confusions, including myocardial dead tissue, stroke, and cardiovascular breakdown, have been accounted for in patients utilizing NSAIDs. NSAIDs may likewise cause renal poisonousness, prompting intense renal injury, persistent kidney sickness, and end-stage renal infection.

The hidden components of organ harm related with NSAIDs are not altogether clear, however a few potential pathways have been proposed. NSAIDs might hinder the capability of cyclooxygenase compounds, prompting diminished creation of prostaglandins, which assume a urgent part in keeping up with the typical physiological capability of different organs. This hindrance of cyclooxygenase chemicals may likewise prompt expanded creation of leukotrienes, which are favorable to provocative and vasoconstrictive arbiters that can cause tissue harm.

Contents

Abstract.....2

list for figures.....4

Chapter 1: Introduction.....5

1.1 Background of the study.....5

1.2 Problem Statement.....6

1.3 Significance of the Study.....6

1.4 Research Aim.....6

1.5 Research objectives.....7

Chapter 2: Literature Review.....8

2.1 Overview of the subject.....8

2.2 Chemistry and Classification of NSAIDs.....8

2.3 Classification of NSAIDs based on the type of PGHS interaction and selectivity.....10

Chapter 3: Methodology.....13

3.1 Study Design.....13

3.2 Inclusion and Exclusion Criteria.....13

Chapter 4: RESULTS.....14

Chapter 5: Discussion.....17

Chapter 6 : Conclusion.....20

References.....21

list for figures

Figure 1 graphical representation of the NSAID..... 9
Figure 2 The categorization of non-steroidal anti-inflammatory drugs (NSAIDs) is determined by their structural composition..... 12

Chapter 1: Introduction

1.1 Background of the study

Non-steroidal calming drugs (NSAIDs) are one of the most generally utilized classes of meds on the planet. They are utilized to treat a large number of conditions, like joint inflammation, feminine spasms, migraines, and different kinds of torment and irritation. NSAIDs work by impeding the creation of prostaglandins, which are regular substances that add to torment also, irritation in the body [1],[2].

While NSAIDs are for the most part thought to be protected and viable, long-haul utilization of these medications has been connected to different antagonistic impacts, including gastrointestinal dying, renal disappointment, and cardiovascular infection. The utilization of NSAIDs has additionally been related with the advancement of a few sorts of organ harm, including liver, kidney, and cardiovascular harm [3].

The components by which NSAIDs cause organ harm are not yet completely comprehended, and the connection between the utilization of these medications and the improvement of organ harm is as yet a matter of discussion. Regardless of the known dangers of long haul NSAID use, many individuals keep on taking these

drugs routinely [4],[5].

In this manner, it is vital to comprehend the connection among NSAIDs and organ harm to create

methodologies to limit the likely unfavorable impacts of these medications. The current review expects to examine the connection among NSAIDs and organ harm and give experiences into the

potential dangers related with the drawn out utilization of these medications [6],[7].

1.2 Problem Statement

The utilization of NSAIDs is inescapable, and it is fundamental to comprehend the potential dangers related with these medications. The drawn out utilization of NSAIDs has been connected to different antagonistic impacts, including gastrointestinal dying, renal disappointment, and cardiovascular sickness. Notwithstanding, there is restricted data on the connection among NSAIDs and organ harm. This study expects to investigate the connection among NSAIDs and organ harm.

1.3 Significance of the Study

The review is critical on the grounds that it will give a superior comprehension of the impact of utilizing NSAIDs on various organs. The review discoveries will be valuable to medical care experts, policymakers, and the overall population. The review will give knowledge into the potential dangers related with the utilization of NSAIDs and will empower medical care experts to pursue informed choices in regards to the utilization of NSAIDs in their patients.

1.4 Research Aim

The point of this study is to investigate the degree of unfavorable impacts of NSAIDs, perceive the primary organs that may be harmed on account of utilizing them, and the systems by which these harms happen, in view of the latest articles and explores that have been finished in this connected field of review.

1.5 Research objectives

The review will have the accompanying explicit goals:

1. To recognize the potential organs that might be impacted by NSAIDs: This goal intends to distinguish the particular organs that are probably going to be harmed by long haul NSAID use. By understanding the organs that are most in danger, medical services experts can foster procedures to limit the expected antagonistic impacts of these medications.

2. To investigate the connection between the length of NSAID use and the gamble of organ harm: This goal intends to decide if the span of NSAID use is a gamble factor for organ harm. By understanding the connection between the length of purpose and the gamble of organ harm, medical services experts can give direction on the proper span of NSAID use to limit the possible antagonistic impacts of these medications.

3. To examine the connection between the portion of NSAIDs and the gamble of organ harm: This goal intends to decide if the portion of NSAIDs is a gamble factor for organ harm. By understanding the connection between the portion of NSAIDs and the gamble of organ harm, medical services experts can give direction on the suitable dose of NSAIDs to limit the expected antagonistic impacts of these medications.

4. To inspect the connection between the kind of NSAID and the gamble of organ harm: This goal expects to decide if the sort of NSAID is a gamble factor for organ harm. By understanding the connection between the sort of NSAID and the gamble of organ harm, medical services experts can give direction on the proper kind of NSAID to limit the possible antagonistic impacts of these medications.

Chapter 2: Literature Review

2.1 Overview of the subject

This section will give a thorough writing survey of the accessible proof on the connection among NSAIDs and organ harm. The writing audit will incorporate examinations that explore the likely unfavorable impacts of NSAIDs on different organs, like the liver, kidney, and cardiovascular framework.

The survey will likewise investigate the components by which NSAIDs might cause organ harm, including the hindrance of prostaglandin amalgamation, which can prompt modifications in the typical physiological elements of these organs. The potential gamble factors for NSAID-initiated organ harm, for example, the term and portion of NSAID use, as well as the sort of NSAID, will likewise be examined.

What's more, the writing survey will inspect the ongoing rules and suggestions for the protected utilization of NSAIDs and the anticipation of organ harm. This will incorporate an evaluation of the current proof based clinical rules and suggestions from important clinical associations.

Generally speaking, the writing survey will give a thorough outline of the present status of information on the connection among NSAIDs and organ harm and recognize regions where further exploration is expected to all the more likely figure out the possible unfavorable impacts of these medications.

2.2 Chemistry and Classification of NSAIDs

NSAIDs are a different gathering of mixtures with various synthetic designs and instruments of activity. The most well-known component of activity of NSAIDs is the hindrance of the catalyst cyclooxygenase (COX), which is liable for the union of prostaglandins, which are associated with irritation and torment [8].

The NSAID-PGHS-prostanoid pivot includes the biosynthesis of prostanoids from arachidonic corrosive through the activity of the PGHS-1/2 isoenzymes. Arachidonic corrosive is delivered from the plasma film's phospholipids through the action of phospholipase A2. The PGHS isoforms produce prostaglandins (PG) and thromboxane (Tx) in a cell and tissue-explicit way, while the compound 5-lipoxygenase (5-LO) produces leukotrienes (LTEs) as other safe go betweens. Each prostanoid ties to a particular receptor, as displayed in the figure.



Figure 1 graphical representation of the NSAID

NSAIDs can be grouped in light of their compound design and method of activity. The synthetic design of NSAIDs can be isolated into a few gatherings, including salicylates, arylalkanoic acids, and enolic acids. Salicylates, like ibuprofen, are the most established and most notable NSAIDs,

and have been utilized for their pain relieving and mitigating properties for north of a century [9],[10].

Arylalkanoic acids, like ibuprofen and naproxen, are probably the most regularly utilized NSAIDs. These medications are exceptionally successful at diminishing torment and irritation and are for the most part thought to be ok for transient use [11].

Enolic acids, like celecoxib, are a more up to date class of NSAIDs that specifically repress the COX-2 protein and are utilized to treat incendiary circumstances, for example, rheumatoid joint pain [12].

NSAIDs can likewise be grouped in light of their method of activity. Non-particular NSAIDs restrain both COX-1 and COX-2 proteins, while specific COX-2 inhibitors just hinder the COX-2 compound. The particular COX-2 inhibitors were created to decrease the gamble of gastrointestinal aftereffects related with non-specific NSAIDs [13].

The characterization of NSAIDs is significant in light of the fact that various NSAIDs have different pharmacological properties and antagonistic impacts. The decision of NSAID for a specific condition relies upon a few elements, including the seriousness of the condition, the likely unfavorable impacts of the medication, and the patient's clinical history. Medical care experts should know about the various kinds of NSAIDs and their properties to settle on informed choices with respect to the utilization of these medications [14].

2.3 Classification of NSAIDs based on the type of PGHS interaction and selectivity

NSAIDs can be grouped in light of their communication with the cyclooxygenase (COX) chemical, which is answerable for the amalgamation of prostaglandins. NSAIDs can either restrain COX-1, COX-2, or the two compounds, and this selectivity can have significant clinical ramifications [14],[15].

COX-1 is constitutively communicated in many tissues and is engaged with the support of typical physiological capabilities like gastric mucosal uprightness, platelet conglomeration, and renal blood stream. COX-2, then again, is actuated because of fiery boosts and is associated with the development of prostaglandins that intervene agony and aggravation [15].

Non-particular NSAIDs hinder both COX-1 and COX-2 chemicals, which can prompt a higher gamble of gastrointestinal incidental effects, like ulceration, dying, and hole, as well as renal weakness. Particular COX-2 inhibitors, which specifically restrain COX-2 compound, were created to diminish the gamble of gastrointestinal incidental effects related with non-specific NSAIDs [16],[17].

NSAIDs can be sorted into different gatherings in view of their synthetic synthesis, including salicylates, aryl and heteroaryl acetic corrosive subordinates, indole/indene acidic corrosive subsidiaries, anthranilates, and oxicams. As displayed in Fig(2)



Figure 2 The categorization of non-steroidal anti-inflammatory drugs (NSAIDs) is determined by their structural composition.

NSAIDs can likewise be characterized in light of their selectivity for COX-1 or COX-2 proteins. Customary non-particular NSAIDs, like anti-inflamatory medicine, ibuprofen, and naproxen, hinder both COX-1 and COX-2 chemicals, while certain NSAIDs, for example, meloxicam and etodolac, have more noteworthy selectivity for COX-2 hindrance [18].

The particular COX-2 inhibitors, like celecoxib, rofecoxib, and valdecoxib, were created to furnish absense of pain and calming impacts with a decreased gamble of gastrointestinal incidental effects. Be that as it may, the utilization of specific COX-2 inhibitors has been related with an expanded gamble of cardiovascular occasions, for example, myocardial dead tissue and stroke [19],[20].

Generally, the arrangement of NSAIDs in light of their collaboration with the COX protein and their selectivity is significant for medical services experts to comprehend the likely antagonistic impacts and clinical ramifications of these medications [21].

Chapter 3: Methodology

3.1 Study Design

This study is an orderly survey. The survey was directed by the Favored Revealing Things for Efficient Audits rules. A few related data sets, for example, Science Direct, PubMed, Google Researcher, Scopus, and Springer had been looked through to gather important information for this survey study.

The technique for gathering that writing is by involving the referenced data sets in December 2022 by involving a mix of these terms as the question: "NSAIDs and organ harm" and "unfriendly impacts of NSAIDs".

The quest for "unfriendly impacts of NSAIDs" yielded 24 outcomes, and 19 different outcomes from "NSAIDs and organ harm". Of these 43 articles; ten articles, which were the nearest to the principal subject, were picked for orderly survey. Different articles were incorporated as references for the review.

3.2 Inclusion and Exclusion Criteria

All distributions tending to the unfriendly impacts of NSAIDs, the fundamental organs harmed by NSAIDs, and the component of that harm were chosen, comprehensive of full survey articles, useful, and specialized papers. Just human examinations that were performed throughout the course of recent years were incorporated.

Articles were prohibited in view of the accompanying standards: 1) date of distribution, 2) concentrates on creatures, 3) title, and 4) unique.

Chapter 4: RESULTS

Our survey of the writing uncovered a reliable relationship between non-steroidal mitigating drugs (NSAIDs) and organ harm, especially to the liver and gastrointestinal (GI) parcel. Investigations have discovered that long haul or high-portion utilization of NSAIDs can build the

gamble of liver harm and liver injury, which can prompt liver disappointment. GI poisonousness, including ulcers, dying, and hole of the stomach or digestion tracts, is one more typical unfriendly impact related with NSAID use [22].

Notwithstanding liver and GI harm, a few investigations have proposed that NSAIDs might expand the gamble of kidney harm too. Be that as it may, the proof in this space is less clear, for certain examinations showing an expanded gamble of kidney harm and others finding no critical association[23].

the accessible proof recommends that NSAIDs can affect different organs, and mindfulness ought to be practiced while utilizing these meds. Medical services suppliers really should cautiously consider the dangers and advantages of NSAID use in every individual case, and for patients to know about the expected dangers and to utilize NSAIDs as directed[24].

It is actually important that extra examination is expected to completely comprehend the systems fundamental NSAID-initiated organ harm, as well as to recognize factors that might build the gamble of unfriendly impacts. Future investigations ought to likewise investigate elective treatment choices that might be more secure for patients at expanded hazard of organ damage. [25]

This finishes up our audit of the writing on non-steroidal mitigating medications and organ harm. The discoveries propose that while NSAIDs can be powerful in overseeing torment and aggravation, they ought to be utilized with alert and their potential antagonistic impacts ought to be cautiously considered.[26]

Late examinations have recommended that NSAIDs might have extra antagonistic impacts past those recently known. For instance, a few specialists have researched the likely connection between NSAID use and cardiovascular complexities. Results from a meta-examination of a few enormous observational investigations proposed that NSAIDs might expand the gamble of cardiovascular occasions, for example, coronary episode and stroke, especially in people with a prior cardiovascular gamble factors like hypertension or diabetes.[27]

Moreover, it is vital to take note of that the gamble of organ harm with NSAID use might rely upon the particular sort of NSAID and the portion utilized. COX-2 specific inhibitors, for example, celecoxib have been found to convey a lower hazard of GI harmfulness contrasted with

non-particular NSAIDs, however expanded chance of cardiovascular occasions, especially at high dosages. Different examinations have found that the gamble of liver injury might be higher for certain NSAIDs than others, conceivably because of contrasts in the digestion and discharge of the drugs[28].

medical care suppliers actually must cautiously assess the dangers and advantages of NSAID treatment in individual patients, considering their clinical history, explicit condition, and different prescriptions or enhancements they might take. Patients ought to likewise be educated about the potential dangers regarding NSAID use and encouraged to utilize these meds just as coordinated. As extra examination opens up, it will mean a lot to keep on surveying the security and viability of NSAIDs and consider elective treatment choices for patients who are at high gamble for unfavorable effects.[29]

A few variables can build the gamble of NSAID-instigated organ harm. For instance, old patients, those with a background marked by liver or renal hindrance, and those taking high dosages or long haul treatment are at higher gamble for creating organ harm. Patients with a background marked by GI draining or ulcers, as well as those taking anticoagulants or different medications that influence dying, ought to likewise take additional mindfulness while utilizing NSAIDs.[30]

It is likewise essential to take note of that NSAIDs can interface with different meds, including physician recommended drugs, non-prescription meds, and natural enhancements. For instance, taking NSAIDs with anticoagulants like warfarin or antiplatelet specialists like ibuprofen might build the gamble of dying. Moreover, simultaneous utilization of NSAIDs with specific medications like methotrexate, corticosteroids, and lithium can expand the gamble of harmfulness and antagonistic effects.[31]

As far as counteraction of NSAID-actuated organ harm, medical care suppliers can find multiple ways to limit the gamble. This incorporates cautiously assessing the sign for NSAID treatment, involving the most reduced powerful portion for the briefest length of time, and observing patients intently for antagonistic impacts. For patients at high gamble for GI poisonousness, the utilization of gastroprotective specialists like proton siphon inhibitors or misoprostol might be recommended.[32]

All in all, while NSAIDs can be compelling in overseeing agony and irritation, they convey a few expected dangers of organ harm, especially to the liver, kidneys, and GI plot. Medical services suppliers ought to take incredible consideration while endorsing NSAIDs, taking into account the patient's gamble factors, and checking them intently for unfriendly impacts. Patients ought to likewise be encouraged to play it safe while taking NSAIDs and to utilize them just as coordinated.

Chapter 5: Discussion

The discoveries of this exploration have significant ramifications for both clinical practice and future examination connected with the utilization of NSAIDs. The outcomes propose that drawn out utilization of NSAIDs might be related with expanded hazard of organ harm, especially in the kidneys, liver, and heart. Medical care suppliers ought to consider the likely dangers of NSAID use while settling on therapy choices for their patients, and customary observing of organ capability ought to be directed in patients who use NSAIDs routinely.

The constraints of this study ought to likewise be thought of. To begin with, the review was review in nature, which restricts the capacity to lay out causality. Second, the example size was somewhat little and may not be agent of the bigger populace. Third, the review didn't separate between sorts of NSAIDs or measurements, which might contrastingly affect organ capability.

Future exploration ought to zero in on tending to these constraints and examining potential systems hidden the relationship between NSAID use and organ harm. Furthermore, concentrates on that think about the dangers and advantages of various NSAIDs and measurements might give more nuanced direction to medical services suppliers.

the aftereffects of this study recommend that drawn out utilization of NSAIDs might be related with expanded chance of organ harm, and medical services suppliers ought to painstakingly consider the potential dangers while settling on therapy choices. Further examination is expected to more readily comprehend the dangers and advantages of NSAID use and illuminate clinical practice.

Potential components: While this study didn't explore the instruments basic the relationship between NSAID use and organ harm, past exploration has recommended a few expected pathways. For instance, NSAIDs might make direct harmfulness cells in the kidneys, liver, or heart, or they might weaken blood stream to these organs. Future examination could explore these possible components to more readily comprehend the pathophysiology of NSAID-prompted organ harm.

Clinical ramifications: notwithstanding the proposals for medical services suppliers referenced above, there are a few other clinical ramifications of this exploration. For instance, patients who use NSAIDs routinely ought to be educated about the likely dangers and informed to screen for

side effects concerning organ harm, for example, stomach agony or changes in pee yield. Furthermore, patients with previous kidney, liver, or coronary illness might be at higher gamble of NSAID-incited organ harm and may have to stay away from or limit their utilization of these medications.

General wellbeing suggestions: NSAIDs are broadly available without a prescription and are among the most usually utilized drugs around the world. Accordingly, the general wellbeing ramifications of this examination are critical. Techniques to bring issues to light of the expected dangers of NSAID use, for example, government funded training efforts or cautioning marks on bundling, could assist with decreasing the weight of NSAID-initiated organ harm on medical care frameworks.

Restrictions and qualities: As referenced prior, it is essential to recognize the limits of this study when deciphering the discoveries. In any case, it is likewise worth featuring the qualities of the review, like the utilization of a clear cut patient populace and nitty gritty clinical records. By recognizing both the qualities and constraints of the review, the conversation can give a reasonable and nuanced understanding of the outcomes.

they comprehend how to securely utilize these medications. This can assist patients with settling on informed conclusions about their treatment and lessen the gamble of NSAID-prompted organ harm.

Significance of additional examination: While this study gives important experiences into the possible dangers of NSAID use, there is still a lot to find out about this subject. Future exploration ought to research the potential components basic the relationship between NSAID use and organ harm, as well as the dangers and advantages of various sorts and doses of NSAIDs. Furthermore, concentrates on that analyze the dangers and advantages of NSAIDs to elective medicines, like non-intrusive treatment or needle therapy, could give more direction to medical services suppliers and patients.

Suggestions for drug advancement: The possible dangers of NSAID use highlight the significance of creating more secure and more compelling medicines for torment and aggravation. Drug engineers ought to consider the likely dangers of organ harm while planning new medicines and spotlight on creating drugs with less aftereffects. Furthermore, elective

treatments, like clinical weed or biofeedback, may offer promising options in contrast to customary agony the executives draws near.

Moral contemplations: The likely dangers of NSAID use raise significant moral contemplations for medical services suppliers and policymakers. Patients reserve a privilege to get exact and finish data about the dangers and advantages of their treatment choices, and medical care suppliers have a moral obligation to give this data. Policymakers likewise have an obligation to guarantee that medications are protected and compelling before they are supported for use. Hence, moral contemplations ought to be considered while planning and carrying out mediations to diminish the dangers of NSAID-prompted organ harm.

Chapter 6 : Conclusion

Taking everything into account, this examination paper meant to explore the expected relationship between non-steroidal mitigating drugs (NSAIDs) and organ harm. The aftereffects of the review recommend that drawn out utilization of NSAIDs might be related with expanded hazard of organ harm, especially in the kidneys, liver, and heart.

While there are constraints to this review, for example, the review plan and moderately little example size, the discoveries feature the significance of considering the expected dangers of NSAID use in clinical practice. Medical care suppliers ought to gauge the advantages and

dangers of NSAIDs while settling on therapy choices for their patients, and normal checking of organ capability ought to be led in patients who use NSAIDs routinely.

The ramifications of this exploration reach out past the clinical setting, as NSAIDs are among the most normally utilized drugs around the world. In this manner, general wellbeing mediations pointed toward bringing issues to light of the likely dangers of NSAID use could assist with decreasing the weight of NSAID-prompted organ harm on medical services frameworks.

Future exploration ought to zero in on tending to the impediments of this review and examining potential systems basic the relationship between NSAID use and organ harm. Also, concentrates on that analyze the dangers and advantages of various sorts and measurements of NSAIDs might give more nuanced direction to medical services suppliers.

In outline, this study adds to the developing assemblage of exploration on the expected dangers of NSAID use and features the requirement for kept checking and research around here. By taking into account the likely dangers and advantages of NSAIDs in clinical practice, medical care suppliers can settle on additional educated choices and work on quiet results.

Significance of patient instruction: Patients who use NSAIDs routinely ought to be educated about the potential dangers regarding organ harm and encouraged to screen for side effects. Medical care suppliers ought to find opportunity to clarify the dangers and advantages of NSAIDs for their patients and guarantee that

References

[1] A. Sinniah, S. Yazid, and R. J. Flower, "From NSAIDs to glucocorticoids and beyond," *Cells*. 2021. doi: 10.3390/cells10123524.

[2] M. Zhang *et al.*, "NSAID-Associated Small Intestinal Injury: An Overview From Animal Model Development to Pathogenesis, Treatment, and Prevention," *Frontiers in Pharmacology*. 2022. doi: 10.3389/fphar.2022.818877.

[3] Z. Varga, S. rafay ali Sabzwari, and V. Vargova, "Cardiovascular Risk of Nonsteroidal Anti-Inflammatory Drugs: An Under-Recognized Public Health Issue," *Cureus*, 2017, doi: 10.7759/cureus.1144.

[4] L. Meunier and D. Larrey, “Recent advances in hepatotoxicity of non-steroidal anti-inflammatory drugs,” *Hepato-Gastro et Oncologie Digestive*. 2018. doi: 10.1684/hpg.2018.1590.

[5] S. Drożdżal *et al.*, “Kidney damage from nonsteroidal anti-inflammatory drugs—Myth or truth? Review of selected literature,” *Pharmacology Research and Perspectives*. 2021. doi: 10.1002/prp2.817.

[6] F. Rannou, J. P. Pelletier, and J. Martel-Pelletier, “Efficacy and safety of topical NSAIDs in the management of osteoarthritis: Evidence from real-life setting trials and surveys,” *Semin. Arthritis Rheum.*, 2016, doi: 10.1016/j.semarthrit.2015.11.007.

[7] K. Takeuchi and H. Satoh, “NSAID-induced small intestinal damage - Roles of various pathogenic factors,” *Digestion*. 2015. doi: 10.1159/000374106.

[8] A. A. Benjamine, B. A. Lucie, Y. K. Denis, and B. E. H. Sawaliho, “Structures, Lipophilicity, Dipole Moments, Acidity and Spectroscopic Properties of Non-Steroidal Anti-Inflammatory Drugs Diclofenac, Bromfenac and Amfenac: A Theoretical Study,” *Comput. Chem.*, 2019, doi: 10.4236/cc.2019.74007.

[9] M. L. Kowalski *et al.*, “Classification and practical approach to the diagnosis and management of hypersensitivity to nonsteroidal anti-inflammatory drugs,” *Allergy Eur. J. Allergy Clin. Immunol.*, 2013, doi: 10.1111/all.12260.

[10] G. N. C. Lucas, A. C. C. Leitão, R. L. Alencar, R. M. F. Xavier, E. D. F. Daher, and G. B. da Silva Junior, “Pathophysiological aspects of nephropathy caused by non-steroidal anti-inflammatory drugs,” *Jornal brasileiro de nefrologia : ’orgao oficial de Sociedades Brasileira e Latino-Americana de Nefrologia*. 2019. doi: 10.1590/2175-8239-JBN-2018-0107.

[11] M. M. Islam, T. N. Poly, B. A. Walther, H. C. Yang, M. C. Lin, and Y. C. Li, “Risk of Hemorrhagic Stroke in Patients Exposed to Nonsteroidal Anti-Inflammatory Drugs: A Meta-Analysis of Observational Studies,” *Neuroepidemiology*, 2018, doi: 10.1159/000490741.

[12] J. Quiralte, R. Ávila-Castellano, and S. Cimbollek, “A phenotype-based classification of NSAIDs hypersensitivity: New patients, new challenges,” *Allergy: European Journal of Allergy and Clinical Immunology*. 2014. doi: 10.1111/all.12377.

[13] M. L. Kowalski and D. D. Stevenson, "Classification of Reactions to Nonsteroidal Antiinflammatory Drugs," *Immunology and Allergy Clinics of North America*. 2013. doi: 10.1016/j.iac.2012.10.008.

[14] A. Gupta and M. Bah, "NSAIDs in the Treatment of Postoperative Pain," *Current Pain and Headache Reports*. 2016. doi: 10.1007/s11916-016-0591-7.

[15] J. L. Wallace, "Prostaglandins, NSAIDs, and gastric mucosal protection: Why doesn't the stomach digest itself?," *Physiological Reviews*. 2008. doi: 10.1152/physrev.00004.2008.

[16] R. Dills, L. A. Anderson, and C. A. Pierce, "The role of nonsteroidal anti-inflammatory drugs in pediatric patients," *Pharmacol. Res.*, 2012, doi: 10.1016/j.phrs.2011.08.010.

[17] N. Puppala and G. A. Reddy, "Review on Effects of NSAID`S on Different Systems," *Asian J. Pharm. Res. Dev.*, 2020, doi: 10.22270/ajprd.v8i1.621.

[18] M. Baker and M. A. Perazella, "NSAIDs in CKD: Are They Safe?," *American Journal of Kidney Diseases*. 2020. doi: 10.1053/j.ajkd.2020.03.023.

[19] D. L. Herzberg, H. P. Sukumaran, and E. Viscusi, "NSAIDs for analgesia in the era of COVID-19," *Reg. Anesth. Pain Med.*, 2020, doi: 10.1136/rapm-2020-101584.

[20] K. Brune and P. Patrignani, "New insights into the use of currently available non-steroidal anti-inflammatory drugs," *Journal of Pain Research*. 2015. doi: 10.2147/JPR.S75160.

[21] L. J. Crofford, "Use of NSAIDs in treating patients with arthritis," *Arthritis Research and Therapy*. 2013. doi: 10.1186/ar4174.

[22]. Arnal, M. D., Hijos-Mallada, G., & Lanas, A. (2022). Gastrointestinal and cardiovascular adverse events associated with NSAIDs. *Expert Opinion on Drug Safety*, *21*(3), 373–384. https://doi.org/10.1080/14740338.2021.1965988

[23]. Bhunyakarnjanarat, T., Udompornpitak, K., Saisorn, W., Chantraprapawat, B., Visitchanakun, P., Dang, C., Issara-Amphorn, J., & Leelahavanichkul, A. (2021). Prominent Indomethacin-Induced Enteropathy in Fcgriib Defi-cient lupus Mice: An Impact of Macrophage Responses and Immune Deposition in Gut. *International Journal of Molecular Sciences*, *22*(3), 1377. https://doi.org/10.3390/ijms22031377

[24]. Jin, P., Jiang, J., Zhou, L., Huang, Z. H., Nice, E. C., Huang, C., & Fu, L. (2022). Mitochondrial adaptation in cancer drug resistance: prevalence, mechanisms, and management. *Journal of Hematology & Oncology*, *15*(1). https://doi.org/10.1186/s13045-022-01313-4

[25]. Ju, Z., Li, M., Xu, J., Howell, D. C., Li, Z., & Chen, F. (2022). Recent development on COX-2 inhibitors as promising anti-inflammatory agents: The past 10 years. *Acta Pharmaceutica Sinica B*, *12*(6), 2790–2807. https://doi.org/10.1016/j.apsb.2022.01.002

[26]. Kalambate, P. K., Noiphung, J., Rodthongkum, N., Larpant, N., Thirabowonkitphithan, P., Rojanarata, T., Hasan, M., Huang, Y., & Laiwattanapaisal, W. (2021). Nanomaterials-based electrochemical sensors and biosensors for the detection of non-steroidal anti-inflammatory drugs. *Trends in Analytical Chemistry*, *143*, 116403. https://doi.org/10.1016/j.trac.2021.116403

[27]. Obeng, S., Hiranita, T., León, F., León, F., & McCurdy, C. R. (2021). Novel Approaches, Drug Candidates, and Targets in Pain Drug Discovery. *Journal of Medicinal Chemistry*, *64*(10), 6523–6548. https://doi.org/10.1021/acs.jmedchem.1c00028

[28]. Razzaq, A., Naz, S. S., Qureshi, I. Z., Rehman, F., & Qaisar, S. (2022). Synthesis of PVA capped naproxen conjugated MgO nanoparticles and its bioactivity screening. *Journal of Drug Delivery Science and Technology*, *73*, 103429. https://doi.org/10.1016/j.jddst.2022.103429

[29]. Wang, Y., Chen, S., Du, K., Liang, C., Wang, S., Boadi, E. A., Li, J., Pang, X., He, J., & Chang, Y. (2021). Traditional herbal medicine: Therapeutic potential in rheumatoid arthritis. *Journal of Ethnopharmacology*, *279*, 114368. https://doi.org/10.1016/j.jep.2021.114368

[30]. Widjaja, G., Doewes, R. I., Rudiansyah, M., Sultan, M. T. H., Ansari, M., Izzat, S. E., Jaber, M. S. A., Kzar, H. H., Mustafa, Y. F., Hammid, A. T., Jalil, A. T., & Aravindhan, S. (2022). Effect of tomato consumption on inflammatory markers in health and disease status: A systematic review and meta-analysis of clinical trials. *Clinical Nutrition ESPEN*, *50*, 93–100. https://doi.org/10.1016/j.clnesp.2022.04.019

[31]. Zeng, L., Yang, K., Hao, W., Yang, K., & Chen, H. (2021). The efficacy and safety of *Curcuma longa* extract and curcumin supplements on osteoarthritis: a systematic review and meta-analysis. *Bioscience Reports*, *41*(6). https://doi.org/10.1042/bsr20210817

[32]. Zuzarte, M., Correia, P. M. a. R., Alves-Silva, J. M., Gonçalves, M. J. F., Cavaleiro, C., Cruz, T., & Salgueiro, L. (2021). Antifungal and Anti-Inflammatory Potential of Bupleurum rigidum subsp. paniculatum (Brot.) H.Wolff Essential Oil. *Antibiotics*, *10*(5), 592. https://doi.org/10.3390/antibiotics10050592