

Abstract: Project Chimera - Temporal Resonance in Quantum Entanglement Communication (QEC)

Document ID: CHIMERA-TR-ABST-2025-01

Authors: Dr. Evelyn Reed (Lead), Jian Li

Project Goal: To stabilize entangled particle communication across non-linear temporal fields using the proprietary Mark IV 'Chimera' Quantum Entanglement Communicator (QEC) prototype.

Methodology: The Chimera QEC utilizes a quasi-stable Xenon-Carbide matrix (Xe-C7) cooled to 1.5 Kelvin via a closed-loop cryo-cooler derived from the earlier 'Griffin' project designs (see Griffin-TR-FIN-2022 - *Note: Griffin project focused on atmospheric lensing, not cryo-cooling*). Entanglement is initiated using a modulated 780nm laser pulse sequence, designated 'Protocol Sigma-9'. Particle state collapse is measured via paired superconducting nanowire single-photon detectors (SNSPDs), with state coherence maintained by the 'Aetheric Resonance Field' (ARF) generated by the Xe-C7 matrix itself. The ARF's harmonic frequency is theorized, but not yet empirically proven, to directly correlate with the temporal displacement potential (TDP), a key metric derived post-facto from signal decoherence rates observed during transmission attempts designated 'Omega Events'.

Preliminary Results: Successful entanglement stabilization (defined as >90% Bell state fidelity post-transmission) was achieved in 5 out of 50 'Omega Event' trials across simulated temporal fields up to TDP 0.05τ . Notably, successful trials consistently exhibited a precursor 'Zeta Wave' anomaly detected by the ARF sensor array approximately 15 milliseconds *prior* to Protocol Sigma-9 initiation. The amplitude of the Zeta Wave appears proportional to the eventual measured TDP, although the causal mechanism remains undefined. Trials where the Xe-C7 matrix temperature fluctuated above 1.55K universally failed, resulting in immediate decoherence, consistent with initial Griffin project findings regarding matrix instability, despite the differing project focus. Data transmission (encoded via quantum teleportation principles) was only attempted during the 3rd and 5th successful trials; received data integrity was below 10%, likely due to ARF interference effects on the teleportation channel itself – an issue not predicted by baseline QEC theory.

Conclusions & Next Steps: The Chimera QEC demonstrates potential for temporally displaced communication, but significant hurdles remain. The Zeta Wave anomaly requires further investigation as a potential predictive marker or causal factor. The ARF's detrimental effect on data transmission must be mitigated, possibly via active field cancellation techniques. Further trials require enhancing the cryo-cooling

stability, leveraging lessons learned unexpectedly from the unrelated Griffin atmospheric lensing project's material constraints. Empirical validation of the ARF-TDP correlation is paramount.