

# Journal of Advanced Pharmacy Research<sup>1</sup>



## A Phytochemical and Biological Review on Plants of the Family Aizoaceae<sup>2</sup>

Khayrya A. Youssif<sup>1</sup>, Ali Elshamy<sup>2</sup>, Mohamed A. Rabeh<sup>1,2</sup>, Nagwan M. Gabr<sup>3</sup>, Eman G. Haggag\*<sup>3</sup>

<sup>1</sup>Department of Pharmacognosy, Faculty of Pharmacy, Modern University for Technology and Information, Cairo, Egypt, <sup>2</sup>Department of Pharmacognosy, Faculty of Pharmacy, Cairo University, Cairo, Egypt, <sup>3</sup>Department of Pharmacognosy, Faculty of Pharmacy, Helwan University, Cairo, Egypt, 11795.

\*Corresponding author: Prof. Dr. Eman Gaber Haggag, Department of Pharmacognosy, Faculty of Pharmacy, Helwan University, Cairo, Egypt, 11795 Tel. +201000023022  
Email address: [wemisir@gmail.com](mailto:wemisir@gmail.com)

Submitted on: 26-04-2019; Revised on: 12-06-2019; Accepted on: 18-06-2019<sup>6</sup>

To cite this article: Youssif, K. A.; Elshamy, A.; Rabeh, M. A.; Gabr, N. M.; Haggag, E. G. A Phytochemical and Biological Review on Plants of the Family Aizoaceae. *J. Adv. Pharm. Res.* **2019**, 3 (4), 158-181. DOI: [10.21608/APRH.2019.12303.1083](https://doi.org/10.21608/APRH.2019.12303.1083)

### ABSTRACT<sup>8</sup>

**Objectives:** This study is aimed to be a comprehensive review of the phytochemical constituents and biological activities of Aizoaceae family plants (Mesembryanthemaceae). **Methods:** This study is covering articles between 1969 and 2018, reviewed from internationally accepted databases and scientific data from scientific Journals. **Results:** Phytochemically studied plants of family Aizoaceae have shown the presence of various classes of compounds including; alkaloids, triterpenes, sterols, lignans, phenolic compounds, betacyanins, and essential oils. Biological studies on plants of family Aizoaceae have indicated various bioactive potentials including antioxidant, antidiabetic, antimicrobial, antitumor, hepatoprotective, anti-inflammatory and other effects. The reported medicinal plants of family Aizoaceae were selected and summarized on the basis of their; phytochemical constituents and biological activities. **Conclusion:** The results of this study may inspire further ethno-botanical and ethno-pharmacological research and investigations toward drug discovery.

**Keywords:** Aizoaceae; Biological activities; Phytochemical constituents<sup>10</sup>

### INTRODUCTION<sup>11</sup>

The Aizoaceae (Mesembryanthemaceae)<sup>12</sup> family is mostly composed of succulent plants of which 99% are found in South or Southwest of Africa. The other one-percent are found in coastal areas of Australia, New Zealand, Mediterranean area, Canary Islands, and the western coasts of Chile and California<sup>1</sup>.

Family Aizoaceae constitutes a major part of the Southern African succulent flora. With approximately 1860 species in 127 genera, and in India the family is represented by three genera, namely, *Sesuvium*, *Trianthema* and *Zaleya* with seven species distributed in the coastal and lowland areas of

peninsular India and the Gangetic plains<sup>2</sup>. It is also South Africa's second largest plant family<sup>3</sup>. Based on the division given by Bittrich & Hartmann<sup>4</sup>, five subfamilies are recognized: Aizooideae, Mesembryanthmoideae, Rushioideae, Sesuvioideae and Tetragonioideae<sup>5</sup>. In Egypt, Boulos<sup>6</sup> recognized five genera, but Hosny<sup>7</sup> listed only four genera without considering *Sesuvium*. According to Boulos<sup>6</sup> and Hosny<sup>7</sup>, *Trianthema* is represented in Egypt by two species: *T. portulacastrum* L. and either *T. triquetra* Willd. Ex Spreng. (according to Boulos<sup>6</sup>) or *T. Crystalline* (Forssk.) Vahl (according to Hosny<sup>7</sup>). *Zaleya* is represented by two species: *Z. pentandra* (L.) C. Jeffrey and *Z. decandra* (L.) Burm. f.<sup>6</sup>, or by only

one species: *Z. pentandra* (L.) C. Jeffrey<sup>7</sup>. This family 1 is represented in Egypt by 6 genera and 10 species<sup>8</sup>. It typically inhabits dry subtropical deserts and wet tropical coasts<sup>5-9</sup>. Native species of Aizoaceae are distributed through Mediterranean coastal habitats and the Nile Valley as weeds of cultivated land, and extend to deserts.

Many members of the family are of economic 2 importance as ornamentals and in cultivation worldwide<sup>10</sup> as some of them are used to stabilize sand dunes in coastal regions<sup>11</sup> while others are important as medicinal plants<sup>12</sup> e.g. *Tetragonia tetragonoides* is used in treatment of enteritis and stomach ache as well as stomach cancer and ulcer<sup>12</sup>. Extracts from these plants are used as preservatives, as a remedy against throat infections and in soap making<sup>13</sup>. Although Aizoaceae is considered as one of the most diverse and abundant families, it was found that it is the least studied in terms of medicinal potential<sup>14</sup>. It would, therefore, be important to extensively investigate other plants of family Aizoaceae for future drug discovery. Thus, we aimed to make a comprehensive review of the chemical constituents and biological activities of family Aizoaceae covering the articles between 1969 and 2018.

## MATERIAL AND METHODS 3

The research strategy employed in the review 4 of family Aizoaceae includes internationally accepted databases like Science Direct, Scopus and Web of Science as well as scientific data collected from scientific Journals.

## RESULTS AND DISCUSSION 5

### Phytochemical constituents

A variety of chemicals have been isolated from 6 different species of family Aizoaceae including; alkaloids, triterpenes, sterols, lignans, flavonoids, phenolic compounds, essential oils, and some miscellaneous compounds<sup>5-15-16</sup>.

### Alkaloids 7

Mesembrine alkaloids are considered to be the 8 primary active constituents of the South African medicinal plant *Sceletium tortuosum* Aizoaceae), and it is used as the dried or fermented aerial material from the plant, which is known as Kanna (Aka, Channa, Kougoed). Traditional regional use ranged from relieving thirst, mild analgesia, and alteration of mood<sup>19</sup>. Only two genera of family Aizoaceae were reported to be rich with alkaloids which are genus *Sceletium* and genus *Trianthema* as shown in Table 1. *Sceletium spp.* is currently the only known genus having species with relatively high levels of mesembrine<sup>19</sup>, which belong to the crinine class of compounds based on the alkaloid skeleton<sup>17</sup>. Jeffs et al.

(1982) categorized the various *Sceletium* alkaloids into 9 four structural groups: (1) the 3a-aryl-*cis*-octahydroindole class (e.g. mesembrine and the new isolated alkaloid channaine)<sup>17-18</sup>, (2) C-seco mesembrine alkaloids (e.g. joubertiamine), (3) alkaloids containing a 2,3 disubstituted pyridine moiety and 2 nitrogen atoms (e.g. *Sceletium* alkaloid A4) and (4) a ring C-seco *Sceletium* alkaloid A4 group (e.g. tortuosamine)<sup>20</sup>. The mesembrine alkaloid is the most studied of the mesembrine alkaloids due to its relative abundance in *S. tortuosum* and biological activity, it was initially partially isolated, characterized, and named by Zwicky, and it was purified and crystallized as picrate<sup>19</sup>. Mesembranol and its acetylated derivative, 4-*O*-acetylmesebrenol was also isolated together with 4-*O*-demethylmesebrenol and 4-*O*-demethylmesebrenol from *Sceletium spp* in addition to mesebrenone. Joubertiamine and its derivatives represented a new structural class different from the known mesebrenes. Joubertiamine, dehydrojoubertiamine, and dihydro-joubertiamine were all referred to as the seco-mesebrenes and were isolated from *Sceletium joubertii*<sup>17</sup>. The preponderance of research on *Sceletium* alkaloids has revolved around isolation and structural elucidation. Very little is known about the distribution and chemotaxonomic patterns of these alkaloids within the genus<sup>17</sup>.

### Di-, Tri-, tetra-terpenes, and sterols: 10

Phytochemical screening and analysis of some 11 species of family Aizoaceae showed presence of di-, tri-, and tetraterpenes.  $\beta$ - amyrin, oleanolic acid, Uvaol were isolated from the leaves' methanolic extract of *C. edulis*<sup>16</sup>, while ecdysterone, were isolated from aqueous and organic extracts of *S. portulacastrum*—and *T. portulacastrum*<sup>15-24</sup> Pentandradione and pentandraone were isolated only from the methanolic extract of *Z. pentandra*<sup>50-51</sup>.

Sterols and their glycosides e.g. (Stigmasterol, 12  $\beta$ -Sitosterol, Stigmasterol 3-*O*- $\beta$ -D-glucoside and  $\beta$ -Sitosterol 3-*O*- $\beta$ -D-glucoside and the novel and recently isolated sterol 17-(5-ethyl-6-methylheptan-2-yl)-4,4,10,13-tetramethyl-hexadecahydro-1H-cyclopenta ( $\alpha$ ) phenanthren-3-ol ) have been reported to be isolated from *T. portulacastrum*<sup>15-16-21-22-24-25-26-27-28</sup> as shown in Table 2.

### Lignans and neolignans: 13

*Aptenia cordifolia* and *T. turgidifolium* leaves 14 are the only reported member of family Aizoaceae to show the presence of lignans and neolignans e.g. Pinorsinol, syringaresinol, di-erythro-syringylglycol- $\beta$ -*O*-4,4'-syringarsinol ether and apteniol A-G were isolated from the methanolic extract of the leaves of *A. cordifolia*<sup>22-29-30-31</sup>. Isoamericanin A was isolated from *T. turgidifolium*<sup>22</sup> as shown in Table 3.

Table 1. Alkaloids isolated from plants of family Aizoaceae

Source	Used part/extract	Chemical constituents	Compound no.
<i>Sceletium spp.</i> <i>Sceletium tortuosum</i> (L.) N. E. Br. <i>Sceletium joubertii</i> L. Bolus	Acid-base extraction of the entire plant	- $\Delta^7$ mesembrenone <sup>17</sup> - 4'-O-demethylmesembranol <sup>17</sup> - Mesembrenol <sup>17</sup> - Mesembranol <sup>17</sup> - 4'-O-demethylmesembrenone <sup>17</sup> - Sceletenone <sup>17</sup> - N-demethyl-N-formyl mesembrenone <sup>17</sup> - O-acetylmesebrenol <sup>17</sup> - Mesembrane <sup>17</sup> - N-demethylmesembrenol <sup>17</sup> - N-demethylmesembranol <sup>17</sup> - Hordenine <sup>17</sup>	1 $\longrightarrow$ 12
<i>Sceletium spp.</i> <i>Sceletium tortuosum</i> (L.) N. E. <i>Sceletium joubertii</i> L. Bolus	Acid-base extraction of the entire plant	- Dehydrojoubertiamine <sup>17</sup> - Dihydrojoubertiamine <sup>17</sup> - Joubertiamine <sup>17</sup> - Joubertinamine <sup>17</sup> - O-methyldehydro-joubertiamine <sup>17</sup> - O-methyljoubertiamine <sup>17</sup> - O-methyldihydro-joubertiamine <sup>17</sup> - 3'-methoxy-4'-O-methyljoubertiamine <sup>17</sup> - 4-(3,4-dimethoxyphenyl)-4-[2-acetylmethylamino) ethyl]cyclohexanone <sup>17</sup> - 3'-methoxy-4'-O-methyljoubertiaminol <sup>17</sup> - 4-(3-methoxy-4-hydroxyphenyl)-4-[2-(acetylmethylamino) ethyl]cyclohexadienone <sup>17</sup> - 3a-(3,4-dimethoxyphenyl)-1-methyl-1,2,3,3a,4,5,6,8,9,9b-decahydro-7H-pyrrolo[2,3-f] quinolin-7-one <sup>17</sup> - Sceletium alkaloid A4 <sup>17</sup> - Touruosamine <sup>17</sup> - N-formyltortuosamine <sup>17</sup> - N-acetyltortuosamine <sup>17</sup> - Mesembrenone <sup>17</sup> - Channaine <sup>18</sup> - Mesembrine <sup>17</sup> - 4'-O-demethylmesembrenol <sup>17</sup>	13 $\longrightarrow$ 32
<i>T. decandra</i> L.	Organic extracts of leaves, fruits and seeds	- Trianthemine <sup>22</sup>	-----
<i>T. portulacastrum</i> L.	Organic extracts of the entire plant	- Punarnavine <sup>23</sup> - Trianthemine <sup>23</sup>	33
<i>Sesuvium portulacastrum</i> L.	Aqueous, methanolic and organic extracts of the whole dried plant	- Capsaicin <sup>21-25</sup>	34

Table 2. Reported plants of family Aizoaceae showing sterols, di-, tri-, and tetra-terpenes: 1

Reported Source	Used part/extract	Chemical constituents	Compound no. 2
<i>Carpobrotus edulis</i> L. Bolus	Methanolic extraxct of the leaves	- $\beta$ - amyirin <sup>16</sup> - Oleanolic acid <sup>16</sup> - Uvaol <sup>16</sup>	35 $\longrightarrow$ 37
<i>Sesuvium portulacastrum</i> L.	Aqueous, ethanolic and dichloromethane extracts of leaves and stems	- Ecdysterone <sup>15</sup> - Ecdysone <sup>15</sup> - 22,23-Dihydrostigmasterol <sup>21-25</sup> - Ethyl iso-allocholate <sup>26</sup> - Squalene <sup>26</sup> - Phytol <sup>26</sup>	38 $\longrightarrow$ 43
- <i>Trianthema decandra</i> L.	Leaves, fruits and seeds organic extracts	- 3-Acetyl aleuritolic acid <sup>22</sup> - Stigmasterol 3- <i>O</i> - $\beta$ -D-glucoside <sup>22</sup> - $\beta$ -Sitosterol 3- <i>O</i> - $\beta$ -D-glucoside <sup>22</sup> - Triantheno <sup>22</sup>	44 $\longrightarrow$ 47
<i>Trianthema portulacastrum</i> L.	Organic extracts of the aerial parts	- Ecdysterone <sup>24</sup> - 3-Acetyl aleuritolic acid <sup>24</sup> - Stigmasterol <sup>24</sup> - $\beta$ -Sitosterol <sup>24</sup> - Stigmasterol 3- <i>O</i> - $\beta$ -D-glucoside <sup>24</sup> - $\beta$ -Sitosterol 3- <i>O</i> - $\beta$ -D-glucoside <sup>24</sup> - Triantheno <sup>22-24</sup> - 17-(5-ethyl-6-methylheptan -2-yl)-4, 4, 10, 13- vetramethyl-hexadecahydro-1H cyclopenta( $\alpha$ ) phenanthren-3-ol] <sup>27</sup>	38 44 $\longrightarrow$ 50
<i>Zaleya pentandra</i> L.	Methanolic extract of the aerial parts	- Pentandradione <sup>28</sup> - Pentandraone <sup>28</sup>	51, 52

### Fatty acids, and fatty alcohols: 3

*Sesuvium portulacastrum* is the only member 4 of family Aizoaceae with data reported on isolates of fatty acids and fatty alcohols as shown in Table 4. It showed the presence of; Linolenic acid, oleic acid, eicosyl ester, 9,12,15-Octadecatrienoic acid, 2,3-dihydroxypropyl ester, (Z,Z,Z)-, hexadecanoic acid ethyl ester, lauric acid, tridecanoic acid, myristic acid, pentadecanoic acid, palmitic acid, heptadecanoic acid, stearic acid, oleic acid, linoleic acid, nonadecanoic acid, arachidic acid, heneicosanoic acid, behenic acid, octadecanoic acid, 1-Docosanol and rhodopsin, which were isolated from dichloromethane extract of its leaves and stems<sup>11-26-32-33</sup>.

### Catechins and phenolic acids 5

Reported data showed that catechins and 6 phenolic acids were isolated from six members only of family Aizoaceae which are; *A. cordifolia*, *C. edulis*, *M. crystallinum*, *S. portulacastrum*, *T. decandra*, and *T. portulacastrum*<sup>14-21-22-25-26-29-33-34-35-36</sup>. It has been reported, that epicatechin was isolated from the organic extracts of the leaves of two different plant species *C. edulis* and *S. portulacastrum*<sup>14-21-22-25-26-29-33-34-35-36</sup>. On the other hand, we found, that 2,6-bis(1,1-dimethylethyl)-4-methylphenol is the only phenolic

compound isolated from *M. crystallinum*. Other isolated 7 catechins and phenolic compounds are shown in Table 5.

### Flavonoids 8

Several flavonoids were isolated from different 9 species of family Aizoaceae. Rutin was isolated from the leaves ethyl acetate extract of *C. edulis* and *Mesembryanthemum forsskaolii*. Moreover, the new compound C-methyl flavone was isolated from the organic extracts of the whole plant of the two species *T. portulacastrum* and *T. decandra*<sup>11-22-24-34-37-38-39-40</sup>. Moreover, 2-(3',4'-dihydroxyphenyl) 3,5,7- trihydroxy-chromen-4 one was also isolated from *T. decandra*<sup>40</sup>. Other isolated compounds are shown in Table 6.

### Betacyanin 10

The genus *Lampranthus* of family Aizoaceae 11 was observed according to the reported data in Table (7) to show the presence of betacyanins. This includes *Lampranthus peersu* and *Lampranthus sociorum*. Betanin was isolated from the two plant species *Drosanthemum floribundum* and *Lampranthus spp.* Also, dopaxanthin was the only betacyanin isolated from *Glottiphyllum longum*, while  $\beta$ -Cyanin was isolated from *T. portulacastrum*<sup>24-41-42-43-44</sup>.

Table 3. Lignans, neolignans isolated from plants of family Aizoaceae 1

Species	Used part/extract	Chemical constituents	Compound no.
<i>Aptenia cordifolia</i> L.F.	Methanolic extract of the leaves	- Pinorsinol <sup>29</sup> - Syringaresinol <sup>29</sup> - Di-erythro-syringylglycol- $\beta$ -o-4,4'-syringaresinol ether <sup>29</sup> - Apteniol A, B, C, D, E, F - Apteniol G <sup>30</sup>	53 $\longrightarrow$ 59
<i>Trianthema turgidifolium</i> F. Muell	Methanolic extract of the leaves	- Isoamericanin A <sup>22</sup>	

Table 4. Fatty acids and fatty alcohols isolated from plants of family Aizoaceae 3

Species	Used part/extract	Chemical constituents	Compound no.
<i>Sesuvium portulacastrum</i> L.	Dichloromethane extracts of leaves and stems	- Linolenic acid <sup>11-32</sup> - Oleic acid eicosyl ester <sup>26</sup> - 9,12,15-Octadecatrienoic acid, 2,3-dihydroxypropyl ester, (Z,Z,Z)- <sup>26</sup> - Hexadecanoic acid ethylester <sup>26</sup>	60 $\longrightarrow$ 63
<i>Sesuvium portulacastrum</i> L.	Dichloromethane extracts of leaves and stems	- Lauric acid <sup>32</sup> - Tridecanoic acid <sup>32</sup> - Myristic acid <sup>11,32</sup> - Pentadecanoic acid <sup>32</sup> - Palmitic acid <sup>11,32</sup> - Heptadecanoic acid <sup>32</sup> - Stearic acid <sup>32</sup> - Oleic acid <sup>11,26,32</sup> - Linoleic acid <sup>11,32</sup> - Nonadecanoic acid <sup>32</sup> - Arachidic acid <sup>32</sup> - Heneicosanoic acid <sup>32</sup> - Behenic acid <sup>11,32</sup> - Octadecanoic acid <sup>33</sup> - 1- Docosanol <sup>26</sup> - Rhodopin <sup>26</sup>	64 $\longrightarrow$ 79

#### Nitrogen containing compounds 5

The reported data showed that some nitrogen-containing compounds like 2-(dimethylamino)-1-phenylethanol, 3-(1H-inol-3-yl) propionic acid, 3-(1H-inol-3-yl)propionic acid methyl ester, (2S,E)-N-[2-Hydroxy-2-(4-hydroxyphenyl)ethyl] Ferulamide, (E)-N-[2-Hydroxy-2-(4-hydroxy-3-methoxyphenyl)-ethyl] ferulamide, and (E)-N-[2-(4-Hydroxyphenyl)-2-propoxyethyl] ferulamide has been isolated from the methanolic extract of *A. cordifolia*, in addition to many compounds e.g. Trans-4-hydroxyprolinebtaine, and Pyrrolo [1,2-A] Pyrazin-1,4-dione,hexahydro-3-(phenylmethyl) were isolated from aqueous and organic extracts of *S. portulacastrum*. Moreover, nicotinic acid is the only isolated nitrogen-containing compound isolated from *T. portulacastrum*<sup>15-21-23-25-29-31-33</sup>, as shown in Table 8.

#### Essential oils 7

The available data reported that only the leaves of *S. portulacastrum* yielded oil via hydro-distillation which upon analysis using GC-MS showed the following composition: o-cymene,  $\alpha$ -pinene, 2- $\beta$ -pinene, trans-caryophyllene, 1,8- cineole, limonene,  $\alpha$ -terpinene,  $\alpha$ -terpinolene, camphene, (-)-bornylacetate, tridecane and  $\alpha$ -humulene<sup>21-45</sup> as shown in Table 9.

#### Miscellaneous compounds: 9

It has been reported, that different miscellaneous compounds have been isolated from five different plants of family Aizoaceae including; *A. cordifolia*, *C. edulis*, *T. portulacastrum* and *S. portulacastrum*<sup>16-23-24-26-29-33-46-47</sup> as shown in Table 10.

Table 5. Catechins and phenolic acids and their esters isolated from plants of family Aizoaceae <sup>1</sup>

Species	Used part/extract	Chemical constituents	Compound no.
<i>Aptenia cordifolia</i> L.F.	Methanolic extract of the leaves	- 4-hydroxybenzoic acid <sup>29</sup> - Dihydrocinnamic acid <sup>29</sup> - 4-hydroxy-dihydrocinnamic acid <sup>29</sup> - Dihydrofrulic acid <sup>29</sup> - 3,4-dimethoxy-dihydrocinnamic acid <sup>29</sup> -3,4-dimethoxy-dihydro cinnamic acid methyl ester <sup>29</sup> - 3,4-dimethoxy-dihydrocinnamic acid ethyl ester <sup>29</sup> - Methylfrulate <sup>29</sup> - Sinapic acid <sup>29</sup> - Methyl 2,5-dihydroxybenzoate <sup>29</sup> - 3,4,5-Trimethoxyphenol <sup>29</sup> - 3-Hydroxy-7,8-dihydro- $\beta$ -ionone <sup>29</sup> - 3,4-dimethoxycinnamic acid <sup>29</sup>	80 $\longrightarrow$ 85
<i>Carpobrotus edulis</i> L. Bolus	Leaves methanolic and ethylacetate extract	- Ferulic acid <sup>34</sup> - Catechin <sup>14</sup> - Epicatechin <sup>14</sup> - Procyanidin (B5) <sup>14</sup>	82 86 $\longrightarrow$ 88
<i>Mesembryanthemum crystallinum</i> L.	Methanolic extract of the leaves	2,6-bis(1,1-dimethylethyl)-4-methylphenol <sup>35,36</sup>	89
<i>Sesuvium portulacastrum</i> L.	Aqueous, ethanolic and dichloromethane extracts of leaves and stems	- Epicatechin <sup>21,25</sup> - Phenol,2,4-bis(1,1-dimethylethyl) <sup>33</sup> - Gallic acid <sup>21,25</sup> - Benzoic acid <sup>21</sup> - Benzoic acid, 4-ethoxy-, ethyl ester <sup>26</sup>	87 90 $\longrightarrow$ 93
<i>Trianthema decandra</i> L.	Leaves, fruits and seeds organic extracts	- 5- Hydroxy-2-methoxy benzaldehyde <sup>22</sup> - <i>P</i> -Methoxybenzoic acid <sup>22</sup> - <i>P</i> -Propoxybenzoic acid <sup>22</sup> - Leptorumol <sup>22</sup>	94 $\longrightarrow$ 97
<i>Trianthema portulacastrum</i> L.	Organic extracts of the aerial parts	- 4-hydroxybenzoic acid <sup>24</sup> - 3,4-Dimethoxycinnamic acid <sup>24,35</sup> - Pyrogallol <sup>24</sup> - 5- Hydroxy-2-methoxy benzaldehyde <sup>35</sup> - Protocatechuic acid <sup>24</sup> - Vanillic acid <sup>24</sup> - <i>O</i> -Coumaric acid <sup>24</sup> - Caffeic acid <sup>24</sup> - <i>P</i> -Methoxybenzoic acid <sup>24,35</sup> - <i>P</i> -Propoxybenzoic acid <sup>24,35</sup> - Leptorumol <sup>24</sup>	80, 81 94 $\longrightarrow$ 102

### Biological activities: <sup>3</sup>

Members of family Aizoaceae are known to have diverse biological activities including anti-hyperlipidemic, antipyretic, diuretic, antioxidant, anticancer, larvicidal, analgesic, anti-rheumatic, anticholera, emetic, laxative, anti-inflammatory and antimicrobial. As well as being used in the treatment of; skin diseases, specific blood diseases, jaundice, cataract, night blindness, heart diseases, joint pain, dropsy, ascites, edema, and others as shown in Table 11.

### Antioxidant activity <sup>5</sup>

Oxidative stress can cause damage to tissues and cells. Free radicals, such as nitric oxide, superoxide anions, and hydroxyl radicals, can result in oxidative stress and may inflict damage in almost every organ. Furthermore, cancers also can arise from excess reactive oxygen species (ROS) that can damage cellular DNA. The antioxidant potential of the following plant organic extracts: *T. portulacastrum*, *Mesembryanthemum forsskaolii*, *Aizoon canariense*,



Table 6. Flavonoids isolated from plants of family Aizoaceae 1

Species	Used part/extract	Chemical constituents	Figure no.
<i>Carpobrotus edulis</i> L. Bolus	Ethyl acetate extract of leaves	- Rutin <sup>34</sup> - Hyperoside <sup>34</sup> - Neohesperidin <sup>34</sup>	103, 104
<i>Mesembryanthemum forsskaolii</i> Hochst. ex Bioss	Methanolic extract of the entire herb	- Rutin <sup>37</sup> - Apigenin <sup>37</sup> - Apigenin-7- <i>O</i> -glucoside <sup>37</sup> - Kaempferol-3- <i>O</i> -glucoside <sup>37</sup> - Isorhamnetin-3- <i>O</i> -β-glucopyranoside <sup>37</sup>	103 105 → 108
<i>Sesuvium portulacastrum</i> L.	Aqueous, ethanolic and dichloromethane extracts of the leaves and stems	- 3,5,4'-trihydroxy-6,7-dimethoxyflavone <sup>11</sup> - 3,5-dihydroxy-6,3',4'-trimethoxy-flavone-7- <i>O</i> -[α-L-rhamnopyranosyl-1 (1→6)- β-D-Glucopyranoside <sup>11</sup>	103, 109
<i>Trianthema portulacastrum</i> L.	Organic extracts of the dried plant	- Quercetin <sup>24</sup> - C-Methyl Flavone <sup>24,38</sup> - 5, 2'-dihydroxy-7-methoxy-6, 8-dimethylflavone <sup>39</sup>	103, 110
<i>Trianthema decandra</i> L.	Leaves, fruits and seeds organic extracts	- C-Methyl Flavone <sup>22</sup> - 2 - (3', 4' dihydroxyphenyl) 3, 5, 7 - trihydroxy-chromen-4 one <sup>40</sup>	103, 111

Table 7. Betacyanins isolated from plants of family Aizoaceae 3

Species	Used part/extract	Chemical constituents	Compound no.
<i>Drosanthemum floribundum</i> (Haw.) N.E.Br.	Methanolic extract of the flowers	- Betanidine <sup>41</sup> - Isobetanidine <sup>41</sup> - Betanin <sup>41</sup> - Isobtanin <sup>41</sup>	112 → 115
<i>Glottiphyllum longum</i> (Haw.) N.E.Br.	Methanolic extract of the flowers	Dopaxanthin <sup>42</sup>	116
<i>Lampranthus spp</i>	Aqueous extract of the flowers	- Betanin <sup>43</sup> - Isobtanin <sup>43</sup>	114, 115
<i>Lampranthus sociorum</i> N.E.Br.	Methanolic extract of the flowers	Betanidin 5- <i>O</i> -[2''- <i>O</i> -( <i>E</i> )-feruloyl- β -(1'',2')-glucuronosyl-β-glucoside] <sup>44</sup>	117
<i>Lampranthus peersii</i> N.E.Br.	Methanolic extract of the flowers	Betanidin- 5- <i>O</i> -[6'- <i>O</i> -( <i>E</i> )-feruloyl- β-glucoside] <sup>44</sup>	118
<i>Trianthema portulacastrum</i> L.	Organic extracts of the dried plant	β-Cyanin <sup>24</sup>	119

*Mesembryanthemum Nodiflorum*, *Mesembryanthemum crystallinum*, *Mesembryanthemum forsskaolii*, *S. portulacastrum* *Mesembryanthemum nodiflorum*, *Carpobrotus edulis*, *Mesembryanthemum edule* *Mesembryanthemum crystallinum* and *T. decandra* were investigated by 1,1-diphenyl-2-picryl hydrazyl (DPPH) and hydrogen peroxide assays. The results indicated that the organic extracts possessed a concentration-dependent free radical-scavenging activity against DPPH and hydrogen peroxide radicals, which was comparable with standard ascorbic acid<sup>15-22-24-36-37-48-49-50</sup>.

#### Anticancer activity: 6

The organic extracts of the following plants: *T. portulacastrum*, *S. portulacastrum*, *Carpobrotus edulis* and *T. decandra* showed anticancer activity against mouse lymphoma cells and hepatic carcinoma using MTT assay<sup>16-22-24-33</sup>.

The protective role of *T. portulacastrum* against diethylnitrosoamine-induced experimental hepatocarcinogenesis was evaluated. Morphometric evaluation of focal lesions showed a reduction of altered liver cell foci/cm<sup>2</sup> and a reduction of the average focal area. A decrease in the percentage of liver

Table 8. Nitrogen containing compounds isolated from plants of family Aizoaceae 1

Species	Used part/extract	Chemical constituents	Compound no. 2
<i>Aptenia cordifolia</i> L.F.	Leaves methanolic extract	<ul style="list-style-type: none"> <li>- 2-(dimethylamino)-1-phenylethanol<sup>29</sup></li> <li>- 3-(1H-inol-3-yl) propionic acid<sup>29</sup></li> <li>- 3-(1H-inol-3-yl) propionic acid methyl ester<sup>29</sup></li> <li>- (2S, E)-N-[2-Hydroxy-2-(4-hydroxyphenyl) ethyl] Ferulamide<sup>31</sup></li> <li>- (E)-N-[2-Hydroxy-2-(4-hydroxy-3-methoxyphenyl)-ethyl] ferulamide<sup>31</sup></li> <li>- (E)-N-[2-(4-Hydroxyphenyl)-2-propoxyethyl] ferulamide<sup>31</sup></li> <li>- (E, E)-N,N-Dityramin-4,40 -dihydroxy-3,50 -dimethoxy-b,30 -biccinnamamide<sup>31</sup></li> <li>- 7-Hydroxy-1-(4-hydroxy-3-methoxyphenyl)-N2, N3-bis(4-hydroxyphenethyl)-6-methoxy-1,2-dihydronaphthalene-2,3-dicarboxamide<sup>31</sup></li> </ul>	120 → 125
<i>Sesuvium portulacastrum</i> L.	Aqueous, methanolic and organic extracts of the whole dried plant	<ul style="list-style-type: none"> <li>- Trans-4-hydroxyprolinebaine<sup>15</sup></li> <li>- Pyrrolo[1,2-A] Pyrazin-1,4-dione, hexahydro-3-(phenylmethyl)<sup>33</sup></li> <li>- Pyrrolo[1,2-A] Pyrazin-1,4-dione, hexahydro-3-(2-methylpropyl)<sup>33</sup></li> <li>- Butanoic acid, pyrrolidide<sup>33</sup></li> <li>- L-proline, N-Valeryl-, hexadecyl ester<sup>33</sup></li> </ul>	126 → 130
<i>Trianthema portulacastrum</i> L.	Organic extracts of the whole dried plant	- Nicotinic acid <sup>23</sup>	131

Table 9. Essential oils isolated from plants of family Aizoaceae 3

Species	Used part/extract	Chemical constituents	Compound no. 4
<i>Sesuvium porulacastrum</i> L.	Leaves hydrodistillation	<ul style="list-style-type: none"> <li>- <i>O</i>-cymene<sup>21,45</sup></li> <li>- <math>\alpha</math>-Pinene<sup>21,45</sup></li> <li>- 2-<math>\beta</math>-Pinene<sup>45</sup></li> <li>- Trans-Caryophyllene<sup>21,45</sup></li> <li>- 1,8- Cineole<sup>21,45</sup></li> <li>- Limonene<sup>21,45</sup></li> <li>- <math>\alpha</math>-Terpinene<sup>21-45</sup></li> <li>- <math>\alpha</math>-Terpinolene<sup>21,45</sup></li> <li>- Camphene<sup>21,45</sup></li> <li>- (-)-Bornylacetate<sup>21,45</sup></li> <li>- Tridecane<sup>21,45</sup></li> <li>- <math>\alpha</math>- Humulene<sup>21,45</sup></li> </ul>	132 → 143



Table 10. Miscellaneous compounds isolated from plants of family Aizoaceae <sup>1</sup>

Species	Used part/extract	Chemical constituents	Compound no. <sup>2</sup>
<i>Aptenia cordifolia</i> L.F.	Methanolic extract of the leaves	- 4-(hydroxymethyl) phenol <sup>29</sup> - 4-(hydroxymethyl)-2,6 dimethoxyphenol <sup>29</sup> - Dehydrolololide <sup>29</sup> - (9R)-9-hydroxymegastigm-4-ene-3-one <sup>29</sup> - Mgastigm-4-ene-3,9-dione <sup>29</sup> - 4-oxo-7,8-dihydro- $\beta$ -ionone <sup>29</sup> - (3R,9R)-3,9-dihydroxymegastigm-5-en-4-one <sup>29</sup> - 3-O-methyl-chiro-inositol <sup>29</sup>	144 $\longrightarrow$ 149
<i>Carpobrotus edulis</i> L. Bolus	Methanolic extract of the leaves	Monogalactosyl diacylglycerol <sup>16</sup>	150
<i>Trianthema portulacastrum</i> L.	Organic extracts of the dried whole plant	- $\beta$ -Carotene <sup>24</sup> - Ascorbic acid <sup>23</sup>	151, 152
<i>Trianthema decandra</i> L.	Methanolic extract of the leaves	Bis (2-ethyl hexyl) phthalate <sup>46</sup>	153
<i>Sesuvium portulacastrum</i> L.	Aqueous, methanolic and organic extracts of the dried plant	- Hentriacontane <sup>33</sup> - L-(+)-ascorbic acid, 2,6-dihexaecoate <sup>33</sup> - Vitamin E <sup>26</sup> - 1-monolinoleoylglycerol-trimethylsilyl ether <sup>26</sup> - Dibutylphthalate <sup>47</sup> - Diisooctyl phthalate <sup>47</sup>	154 $\longrightarrow$ 159

parenchyma occupied by foci seems to suggest the anticarcinogenic potential of the plant extract in DENA-induced hepatocarcinogenesis<sup>22</sup>. <sup>3</sup>

#### Analgesic, antinociceptive, antihyperglycemic and hepatoprotective activity <sup>4</sup>

The methanol extract of the leaves of *T. portulacastrum* and *T. decandra* have remarkable analgesic, antinociceptive activity, antihyperglycemic and hepatoprotective activity<sup>22-24</sup>. <sup>5</sup>

The analgesic activity of the leaf extract of *T. decandra* was detected by hot plate and acetic acid-induced writhing response method. The results indicate that the administration of leaf extract of *T. decandra* exhibit central analgesic properties since it exerted a significant and dose-dependent effect on the chemical (acetic acid-induced) and thermic (heat) painful stimuli from the respective doses of 100 and 200 mg/kg<sup>22</sup>. <sup>6</sup>

Many natural plant extracts have been investigated with respect to the suppression of glucose production from carbohydrates in the gut or glucose absorption from the intestine. <sup>7</sup>

$\alpha$ - amylase catalyzes the hydrolysis of  $\alpha$  -1,4- glucosidic linkages of starch, glycogen and various <sup>8</sup>

oligosaccharides and  $\alpha$ -glucosidase further breaks down the disaccharides into simpler sugars, readily available for the intestinal absorption. The inhibition of their activity, in the digestive tract of humans, is considered to be effective to control diabetes by diminishing the absorption of glucose decomposed from starch by these enzymes. Therefore, effective and nontoxic inhibitors of  $\alpha$  - amylase and  $\alpha$ -glucosidase have long been sought, and for the first time, studies have revealed the anti-diabetic potential of *T. decandra* and these studies could be helpful to develop medicinal preparations for diabetes<sup>22</sup>. <sup>9</sup>

The probable mechanism by which *T. portulacastrum* exerts its hepatoprotective action against paracetamol or thioacetamide-induced hepatocellular metabolic alterations could be by the stimulation of hepatic regeneration through an improved synthesis of protein or accelerated detoxification and excretion<sup>22</sup>. <sup>10</sup>

#### Anti-inflammatory activity <sup>11</sup>

The anti-inflammatory activity of methanolic and organic extracts of the following plants: *T. portulacastrum*, *Aizoon canariense*, *Aizoon hispanicum*, <sup>12</sup>


**Table 11. Different plant species and their biological activities reported in family Aizoaceae**

1


Plant name	Biological activities	References
<i>Aptenia cordifolia</i> L.F.	- Algicide activity	29
<i>Aizoon canariense</i> L.	- Antifungal	48,51,52
	- Antioxidant	
	- Antibacterial	
	- Skin diseases	
	- Emetic	
	- Antiinflammatory	
	- Antifungal	
	- Wound healing	
<i>Aizoon hispanicum</i> L.	Antiinflammatory	51
<i>Carpobrotus acinaciformis</i> L.	Antidiarrhea and dysentery	50,53
<i>Carpobrotus edulis</i> L. Bolus	- Antioxidant	16,34,35,53,54
	- Anticancer	
	- Antibacterial	
	- Antidiarrhea and dysentery	
	- Anti HFFIV/AIDS	
<i>Carpobrotus muirii</i> L.	Antidiarrhea and dysentery	53
	- Antioxidant	
<i>Galenia Africana</i> L.	- Antituberculosis	55
	- Asthma and obstructive pulmonary diseases	
<i>Gisekia phernaceoides</i> L.	Antibacterial	56
<i>Lampranthus francisci</i> L. Bolus	Anticandidal	57
<i>Mesembryanthemum anatomicum</i> Haw.	- Mood altering	58
	- CNS stimulant	
<i>Mesembryanthemum crystallinum</i> L.	- Antioxidant	35,36,48,50
	- Antifungal	
	- Antibacterial	
<i>Mesembryanthemum forsskaolii</i> Hochst. ex Bioss	- Antioxidant	35,37
	- Antibacterial	
	- Antifungal	
<i>Mesembryanthemum nodiflorum</i> L.	- Antibacterial	35,49
	- Antioxidant	
	- Antifungal	
<i>Sceletium</i> spp.	- Cytotoxic	17,58
	- Antiinflammatory	
	- Asthma and obstructive pulmonary diseases	
	- Cytotoxic	
	- Psoriasis	
	- Cocaine like activity	
	- Antidepressant	
	- Psychiatric conditions	
<i>Sceletium tortuosum</i> (L.) N. E. Br.	- Antianxiety	17,58,59
	- Antidote in alcohol poisoning	
	- Antidepressant	
	- Psychiatric conditions	
	- Antianxiety	
<i>Sesuvium portulacastrum</i> L.	- Hypnotic and sedative	15,33
	- Antioxidant	
	- Anticancer	
	- Antifungal	
	- Antibacterial	
	- Antiulcerogenic	
	- Kidney disorders	

Table 11. Cont. 1


<i>Sesuvium verrucosum</i> Raf.	Cytotoxic	21	2
<i>Trianthema decandra</i> L.	<ul style="list-style-type: none"> <li>- Antioxidant</li> <li>- Anticancer</li> <li>- Analgesic and antinociceptive</li> <li>- Antihyperglycemic</li> <li>- Hepatoprotective</li> <li>- Antiinflammatory</li> <li>- Antibacterial</li> <li>- Antifungal</li> <li>- Antiulcerogenic</li> <li>- Wound healing</li> </ul>	22,40	
<i>Zaleya pentandra</i> L.	<ul style="list-style-type: none"> <li>- Antiacetylcholinesterase</li> <li>- Antibutyrylcholinesterase</li> <li>- Antifungal</li> <li>- Stomach diseases</li> <li>- Respiratory tract infection and cough</li> <li>- Asthma and obstructive pulmonary diseases</li> <li>- Laxative</li> <li>- Blood diseases</li> <li>- Jaundice</li> <li>- Larvicidal</li> <li>- Cataract and night blindness</li> </ul>	28,60,61	




(1)  $\Delta^7$  Mesembrenone 3




(2) 4'-O-demethylmesembranol 5




(3) Mesembrenol 7




(4) Mesembranol 9




(5) 4'-O-demethylmesembranone 11




(6) Sceletenone 12




(7) N-demethyl-N-formyl mesembrenone 14




(8) O-acetylmesebrenol 15




(9) Mesembrane 16




(10) N-demethylmesembrenol 2




(11) N-demethylmesembranol 4




(12) Hordinine 6




(13) Dehydrojoubertiamine 8




(14) Dihydrojoubertiamine 10




(15) Joubertiamine 12




(16) Joubertinamine 15




(17) O-methyldehydro joubertiamine 17




(18) O-methyl joubertiamine 18




(19) O-methyl dihydrojoubertiamin 20




(20) 3'-methoxy-4'-O-methyl joubertiamine 22




(21) 4-(3,4dimethoxyphenyl)-4-[2-(acetyl methyl) amino] ethyl cyclohexanone 24




(22) 3'-methoxy-4'-O-methyl joubertiaminol 26




(23) 4-(3-methoxy-4-hydroxy phenyl)-4-[2-(acetyl methyl) amino] ethyl cyclohexanone 29




(24) 3a-(3,4-dimethoxy phenyl)-1-methyl-1,2,3,3a,4,5,6,8,9,9b-decahydro-7H-pyrrolo[2,3-f]quinolin-7-one 30




(25) Sceletium alkaloid A4 35




(26) Touruosamine 34




(27) N-formyl tortuosamine 36




(28) *N*-acetyl tortuosamine




(29) Mesembrenone




(30) Mesembrine




(31) 4'-*O*-demethylmesemrenol




(32) Channaine




(33) Punarnavine




(34) Capsaicin




(35)  $\beta$ -Amyrin




(36) Oleanolic acid




(37) Uvaol




(38) Ecdysterone




(39) Ecdysone



(40) 22, 23-Dihydrostigmasterol




(41) Ethyl iso-allocholate




(42) Squalene


1




(43) Phytol




(44) 3-Acetyl aleuritic acid




(45) Stigmasterol 3- O-β-D-glucoside




(46) β-Sitosterol 3- O- β-D-glucoside




(47) Trianthanol




(48) Stigmasterol




(49) β-Sitosterol




(50) 17-(5-ethyl-6-methylheptan -2-yl)-4, 4, 10, 13- vetramethyl-hexadecahydro-1H cyclopenta(α) phenanthren-3-ol]



(51) Pentandradione




(52) Pentandraone



(53)

Pinorsinol R=R1=H  
Syringaresinol R=OCH<sub>3</sub>, R1=H


2



(54) Di-erythro-syringylglycerol-β-o-4,4'- syringaresinol ether


3






(55)


**Apteniol A** R1=R2=R3=R4  
**Apteniol B** R1=OMe; R2=R3=R4=H  
**Apteniol D** R1=R2=R3=OMe; R4=H  
**Apteniol E** R1=R2=R3=R4=OMe




(56) Apteniol C




(57) Apteniol F




(59) Isoamericanin A




(60) Linolenic acid



(58) Apteniol G




(61) Oleic acid eicosyl ester




(62) 9,12,15-Octadecatrienoic acid  
2,3-dihydroxy propyl ester




(63) Hexadecanoic acid ethyl ester




(64) Lauric acid



(65) Tridecanoic acid




(66) Myristic acid




(67) Pentadecanoic acid



(68) Palmitic acid




(69) Heptadecanoic acid




(70) Stearic acid



(71) Oleic acid



(72) Linoleic acid



(80) 4-hydroxy benzoic acid 3

Dihydrocinnamic acid  $R=R_1=R_2=H$  4  
4-hydroxy-dihydrocinnamic acid  $R=R_1=H, R_2=OH$




Dihydrofrulic acid  $R=H, R_1=OCH_3, R_2=OH$  5

3,4-dimethoxy-dihydrocinnamic acid 6  $R=H, R_1=R_2=OCH_3$  7


3,4-dimethoxy-dihydrocinnamic acid methyl ester 9  $R=CH_3, R_1=R_2=OCH_3$  8

3,4-dimethoxy-dihydrocinnamic acid ethyl ester 11  $R=CH_2CH_3, R_1=R_2=OCH_3$  10


3,4-dimethoxycinnamic acid  $R=H, R_1=R_2=OCH_3$  12




Ferulic acid  $R=R_1=R_2=H$  14  
Methyl ferulate  $R=CH_3, R_1=R_2=H$   
Sinapic acid  $R=R_2=H, R_1=OCH_3$   
3,4,5 tri-methoxy cinnamic acid  $R=H, R_1=OCH_3, R_2=CH_3$




(85) 3-Hydroxy-7,8-dihydro- $\beta$ -ionone 2




(86) Catechin 10




(87) Epicatechin 19




(88) Procyanidin 3




(89) 2,6-bis(1,1-dimethylethyl)-4-methylphenol 11




(90) Phenol 2,4-bis(1,1-dimethylethyl) 20




(91) Gallic acid 5




(92) Benzoic acid 14




(93) Benzoic acid, 4-ethoxy ethyl ester 22




(94) 5-Hydroxy-2-methoxy benzaldehyde 7




(95) *p*-methoxy benzoic acid 16




(96) *p*-propoxy benzoic acid 24



(97) Leptorumol 8



(98) Pyrogallol 17




(99) Protocatechuic acid

(100) Vanillic acid 23


(101) *O*- coumaric acid 18

(102) Caffeic acid 25




(103) 2


Compound	R1	R2	R3	R4	R5	R6	R7
Rutin	OH	OH	O-Rutinosyl	OH	H	OH	H
Quercetin	OH	OH	OH	OH	H	OH	H
3,5,4'-trihydroxy-6,7-dimethoxyflavone	OH	H	OH	OH	OCH3	OCH3	H
C-Methyl flavone	H	H	H	OCH3	CH3	OCH3	CH3
Hyperoside	OH	OH	O-Galactosyl	OH	H	OH	H




(104) Neohesperidin 5




(105) Apigenin 9




(106) Apigenin-7-O-glucoside 12




(107) Kaempferol 3-O-glucoside 8




(108) Isorhamnetin-3-O-beta-D-glucoside 14




(109) 3,5-dihydroxy-6,3',4'-trimethoxyflavone-7-O-[alpha-L-rhamnopyranosyl (1-6)-beta-D-glucopyranoside] 16




(110) 5,2'-dihydroxy-7-methoxy-6,8-dimethylflavone 18




(111) 2-(3',4'-dihydroxyphenyl)-3,5,7-trihydroxychromen-4-one 19




(112) Betanidine 21




(113) Isobetanine 23




(114) Betanine 25




(115) Isobetanine 27




(116) Dopaxanthin 2




(117) Betanidin 5-O-[2''-O-(E)-feruloyl-β-(1'',2'')-glucuronosyl-β-glucoside] 4




(118) Betanidin-5-O-[6'-O-(E)-feruloyl-β-glucoside] 6



(119) β-cyanine




(120) 2-(dimethylamino)-1-phenylethanol 9




(121) 11


3-(1H-inol-3-yl) propionic acid R=H 12  
3-(1H-inol-3-yl) propionic acid methyl ester R=CH<sub>3</sub>




(122) (2S, E)-N-[2-Hydroxy-2-(4-hydroxyphenyl) ethyl] ferulamide R=H 13




(123) (E)-N-[2-Hydroxy-2-(4-hydroxy-3-methoxyphenyl) ethyl] ferulamide 16




(124) (E, E)-N, N-Dityramin-4,40-dihydroxy-3,50-dimethoxy-b,30-bicinnamamide 18




(E)-N-[2-(4-Hydroxyphenyl)-2-propoxyethyl] ferulamide R= n-Propyl 14




(125) 7-Hydroxy-1-(4-hydroxy-3-methoxyphenyl)-N2, N3-bis(4-hydroxyphenethyl)-6-methoxy-1,2-dihydronaphthalene-2,3-dicarboxamide 20




(126) Trans-4-hydroxyproline betaine 22




(127) Pyrrolo[1,2-A] Pyrazin-1,4-dione, hexahydro-3-(phenylmethyl) 24




**(128) Pyrrolo[1,2-A]  
Pyrazin-1,4-dione, hexahydro  
-3-(2-methylpropyl)**




**(131) Nicotinic acid**




**(134) 2-β-Pinene**




**(137) Limonene**




**(140) Camphene**




**(143) α-Humulene**




**(129) Butanoic acid, pyrrolidide**




**(132) O-cymene**




**(135) Trans-Caryophyllene**



**(138) α-Terpinene**




**(141) Bornylacetate**




**(144)**


4-(hydroxymethyl) phenol R = H  
4-(hydroxymethyl)-2, 6 dimethoxyphenol R = OCH<sub>3</sub>




**(130) L-proline, N-Valeryl-,  
hexadecyl ester**



**(133) α-Pinene**




**(136) 1, 8- Cineole**



**(139) α-Terpinolene**




**(142) Tridecane**




**(145) Dehydrololilide**






(146)


(9R)-9-hydroxymegastigm-4-ene-3-one  
R=OR1=H, α OH  
Megastigm-4-ene-3,9-dione  
R=R1=O




(147) 4-oxo-7, 8-dihydro-β-ionone




(148) (3R, 9R)-3,9-dihydroxymegastigm-5-en-4-one




(149) 3-O-methyl-chiro-inositol




(150) Monogalactosyldiacylglycerol




(151) β-Carotene




(152) Ascorbic acid




(153) Bis (2-ethyl hexyl) phthalate




(154) Hentriacontane




(155) L-(+)-ascorbic acid, 2-6-dihexaecoate




(156) Vitamin E



(157) 1-monolinoleoylglyceroltrimethylsilyl ether



(158) Dibutylphthalate



(159) Diisooctyl phthalate

Structures of isolated metabolites from family Aizoaceae

*T. decandra* and *Sceletium* spp. was evaluated against formaldehyde-induced arthritis in rats, and a significant inhibition of chemically-induced arthritis, indicates anti-inflammatory potential<sup>17-22-24-51</sup>.

Anti-inflammatory activity was evaluated in acute and chronic models. Significant anti-inflammatory activity was observed for chloroform extract of *T. decandra* in both carrageenan, dextran, and mediators induced edema models. The chloroform extract showed maximum inhibition of 58.36% at the dose of 200 mg/kg after 3 hrs of drug treatment in carrageenan-induced paw edema. The chloroform extract of *T. decandra* also exhibited significant anti-inflammatory properties in dextran-induced paw edema model. Dextran-induced paw edema is known to be mediated both by histamine and serotonin. Dextran induces fluid accumulation, which contains little protein and few neutrophils, whereas carrageenan induces protein rich exudation containing large number of neutrophils. The extract effectively suppressed the inflammation produced by both carrageenan and dextran<sup>22</sup>.

#### Antimicrobial activity<sup>3</sup>

It has been reported that the methanolic extract of *M. nodiflorum*, *M. crystallinum*, *M. forsskaolii*, and *Aizoon canariense* showed a broad-spectrum antibacterial and antifungal activity against: *Bacillus subtilis*, *K. pneumonia*, *S. aureus*, *S. pyogenes*, *E. coli*, *A. fumigatus*, *A. nigar*, *C. albicans*, and *Mucor spp*<sup>48</sup>. For *C. edulis* five bioactive flavonoid compounds, rutin, neohesperidin, hyperoside, cactichin, and ferulic acid were isolated from the ethyl acetate fraction and individually or collectively were responsible for the antibacterial against 11 known human pathogenic bacteria, five Gram-positive: *Staphylococcus epidermidis*, *Staphylococcus aureus*, *Bacillus subtilis*, *Streptococcus pneumoniae*, *Streptococcus pyogenes*, and six Gram-negative: *Pseudomonas aeruginosa*, *Haemophilus influenzae*, *Escherichia coli*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Moraxella catarrhalis*<sup>34</sup>. Moreover, the isolated compounds from the ethanolic extract of *Galenia Africana* showed a remarkable antimycobacterial activity against *M. smegmatis* and *M. tuberculosis*<sup>55</sup>.

The extracts and the essential oil from the fresh leaves of *S. portulacastrum* showed antibacterial, antifungal. The ethanolic extract showed potential antibacterial activity against the causative agents and pathogens related to various gastrointestinal disorders leading to indigestion, dysentery, and diarrhea<sup>15</sup>. Moreover, the ethanolic extract of the *S. portulacastrum* showed potential against the causative agents of nosocomial infections, *Staphylococcus aureus* and *E. coli*, while the essential oil exhibited notable antibacterial activity against both Gram-positive and

Gram-negative bacteria as well as significant antifungal<sup>15</sup>. In addition, *S. portulacastrum*, showed positive activity against human immunodeficiency viruses<sup>15</sup>. Finally, it was reported that the new isolated flavonoid 2 - (3', 4' dihydroxy-phenyl) - 3, 5, 7-trihydroxy-chromen-4-one isolated from *T. decandra* showed antibacterial activity against *Pseudomonas aeruginosa* and by molecular docking it was found that FAS II  $\beta$ -hydroxyacyl-ACP (*FabZ*) of *P. aeruginosa* is a potential target of the isolated compound<sup>40</sup>. There are many other reported biological activities of members of family Aizoaceae as shown in Table 11.

#### CONCLUSIONS<sup>7</sup>

From this review, it can be deduced that the major compounds of family Aizoaceae are alkaloids, triterpenes, flavonoids, sterols, lignans, fatty acids, phenolic acids, and essential oils. The review also showed that different extracts of aerial parts of plants of family Aizoaceae posse diverse biological activities such as anti-acetylcholinesterase, anti-butyrylcholinesterase, anti-infective, anti-hyperlipidemic, antipyretic, antifertility, diuretic, nephroprotective and others. It would, therefore, be important to extensively investigate their phytochemicals and pharmacologically determine their activities for future drug discovery and development.

#### Conflict of interest<sup>9</sup>

The authors declare that there is no conflict of interest regarding the publication of this paper.

#### REFERENCES<sup>11</sup>

1. James, A.R. Mesembryanthemaceae, *Newsletter of the Tucson Cactus and Botanical Society*. **1982**, 12, 1.
2. Rasingam, L. Aizoaceae (Magnoliopsida: Caryophyllales) - a new family record to the flora of Andaman Islands, India, *JOTT*. **2012**, 4 (6), 2653-2655.
3. Chesselet, P.; Mössmer, M.; Smith, G. Research priorities in the succulent plant family Mesembryanthemaceae Fenzl, *S. Afr. J. Sci.* **1995**, 91, 197-209.
4. Bittrich, V.; Harbmann, E. The Aizoaceae - a new approach. *Bot. J. Linn. Soc.* **1988**, 97, 239-254.
5. Eggli, U.; Hartmann, K. Illustrated Handbook of succulent plants, Springer-Verlag Berlin Heidelberg GmbH. **2001**, pp. 1-2.
6. Boulos L. Flora of Egypt. AzollaceaeOxalidaceae, Cairo, Egypt: Al-Hadara Publishing, **1999**.
7. Hosny A. Aizoaceae. In: El Hadidi MN, editor. Flora Aegyptiaca. The Palm Press., Cairo I, **2000**, 96-100.

8. Boulos L. Flora of Egypt, checklist. Al-Hadara Publishing, **1999**.
9. Hartmann H. Aizoaceae. In: Kubitzki K, Rohwer JG, Bittrich V, editors. the families and genera of vascular plants. II. Flowering Plants. Dicotyledons: Magnoliid, Hamamelid and Caryophyllid families. Berlin: Springer, **1993**, 37-69.
10. Vivrette, J.; Bleck, E.; Ferren, R. Aizoaceae Flora of North America, ed. N. R. Morin. New York, Oxford: Oxford University Press. **2003**, 4, 75-91.
11. Heywood, H. Flowering plants of the world. Mayflower Books. Elsevier, New York, **2007**.
12. Walters, M.; Figueiredo, E.; Crouch, R.; Winter, D.; Smith, F.; Zimmermann, G.; Mashop, K. Aizoaceae, Council for Scientific and Industrial Research, South Africa. **2012**, 64-69.
13. Rood, B. Uit die veldapteeke. Tafelberg Publishers, Cape Town, **1994**.
14. Elmarie, W.; Johan, C. Pretorius. Purification and identification of active antibacterial components in *Carpobrotus edulis* L. *J. Ethnopharmacol.* **2001**, 76, 87-91.
15. Manbir, K.; Nitika. Review on Sea purslane, *J Pharmacogn Phytochem.* **2015**, 3(5), 22-24.
16. Martins, A.; Vasas, A.; Schelz, S.; Viveiros, M., Molnar, J.; Hohmann, J.; Amaral, L. Constituents of *Carpobrotus edulis* Inhibit P Glycoprotein of MDRI-transfected Mouse Lymphoma Cells, *Anticancer Res.* **2010**, 30(3), 829-836.
17. Gerickea, N.; Viljoenb, M. *Sceletium*— A review update, *J. Ethnopharmacol.* **2008**, 119(3), 653-663.
18. Vealea, C.; Chenc, W.; Chaudharyc, S.; Kituyid, S.; Isaacsb, M.; Hoppe, H.; Edkins, A.; Combrinck, S.; Meharic, B.; Viljoenc, A. NMR structural elucidation of channaine, an unusual alkaloid from *Sceletium tortuosum*, *Phytochem. Lett.* **2018**, 23, 189-193.
19. John, K. Mesembrine Alkaloids: Review of their Occurrence, Chemistry, and Pharmacology, *J. Ethnopharmacol.* **2017**, 195, 10-19.
20. Jeffs, W.; Capps, M.; Redfearn, R. *Sceletium* alkaloids. Structures of five new bases from *Sceletium namaquense*. *J. Org. Chem.* **1982**, 47, 3611-3617.
21. Cybulska, I.; Brudecki, G.; Alassali, A.; Thomsen, J. Phytochemical composition of some common coastal halophytes of the United Arab Emirates, *EJFA.* **2014**, 26(12), 1046-1056.
22. Geethalakshmi, R., Sarada, L.; Ramasamy, K. *Trianthema decandra* L: A review on its phytochemical and pharmacological profile, *Int. J. Eng. Sci. Technol.* **2010**, 2(5), 976-979.
23. Sukalingam, K.; Ganesan, K.; Ponnusamy, K.; Venugopal, V. Pharmacological Properties of *Trianthema portulacastrum* L and its Therapeutic Potential as Complementary Medicine, *JPCBS.* **2015**, 4 (2), 269-274.
24. Yamaki, J.; Kalyan C.; Venkata, N.; Mandal, A.; Bhattacharyya, P.; Bishayee, A. Health-promoting and disease-preventive potential of *Trianthema portulacastrum* Linn. (Gadabani) —An Indian medicinal and dietary plant, *J. Integr. Med.* **2016**, 14 (2), 84-99.
25. Al-Azzawi A.; Alguboori, A.; Hachim, Y., Najat M.; Al Shaimaa, A.; Sad, M. Preliminary phytochemical and antibacterial screening of *Sesuvium portulacastrum* in the United Arab Emirates, *Pharmacogn. Res.* **2012**, 4 (4), 219-224.
26. Sheela, D.; Uthayakumari, F. GC-MS analysis of bioactive constituents from coastal sand dune taxon *Sesuvium portulacastrum* (L.), *Bio. Disc.* **2013**, 4(1), 47-53.
27. Geethalakshmi, R.; Sarada, L. In vitro and in silico antimicrobial activity of sterol and flavonoid isolated from *Trianthema decandra* L. *Microb. pathog.* **2018**, 121, 77-86.
28. Afzal, S.; Chaudhary, B.; Uzair, M.; Afzal, K.; Bokhari, T. Isolation of pentandraone from methanolic extract of aerial parts of *Zaleya pentandra*, *Int. Res. J. Pharm.* **2013**, 4 (10), 21-23.
29. DellaGreca, M.; Fiorentinob, A.; Izzob, A.; Napolia, F.; Purcaroa, R.; Zarrellia, A. Phytotoxicity of Secondary Metabolites from *Aptenia cordifolia*, *Chem. Biodivers.* **2007**, 4, 118-128.
30. DellaGreca, M.; Marino, C.; Previtera, L.; Purcaro, R.; Zarrelli, A. Apteniols A-F, oxyneolignans from the leaves of *Aptenia cordifolia*. *Tetrahedron.* **2005**, 61, 11924-11929.
31. DellaGreca, M.; Previtera, L.; Purcaro, R.; Zarrelli, A. Cinnamic acid amides and lignanamides from *Aptenia cordifolia*. *Tetrahedron.* **2006**, 62, 2877-2882.
32. Chandrasekaran, M.; Senthilkumar, A.; Venkatesalu, V. Antibacterial and antifungal efficacy of fatty acid methyl esters from the leaves of *Sesuvium portulacastrum* L., *Eur. Rev. Med. Pharmacol. Sci.* **2011**, 15, 775-780.
33. Kumar, A.; Kumari, S.; Somasundaram, T. Gas Chromatography-Mass Spectrum (GC-MS) Analysis of Bioactive Components of the Methanol Extract of Halophyte, *Sesuvium portulacastrum* L., *IJAPB.* **2014**, 3 (3), 766-772.
34. Van der Watt, E.; Pretorius, C. Purification and identification of active antibacterial components in *Carpobrotus edulis* L., *J. Ethnopharmacol.* **2001**, 76, 87-91.
35. Ibtissem, B.; Abdelly, C.; Sfar, S. Antioxidant and Antibacterial Properties of *Mesembryanthemum crystallinum* and *Carpobrotus edulis* Extracts, *ACES.* **2012**, 2, 359-365.

36. Ibtissem, B.; Imen, M.; Souad, S. Dosage of 2, 6-Bis (1.1-Dimethylethyl)-4-Methylphenol (BHT) in the Plant Extract *Mesembryanthemum crystallinum*, *J Biomed Biotechnol.* **2010**, 2010, 1-5.
37. Moawad, A.; Amin, E.; Mohammed, R. Diffusion-ordered Spectroscopy of Flavonol Mixture from *Mesembryanthemum forsskaolii* (Aizoaceae), *EJMP.* **2016**, 16(1), 1-8.
38. Kavitha, D.; Parvatham, R.; Padma, R. Assessment of *trianthema portulacastrum* for its antimicrobial potential and investigation of their phytochemicals using HPTLC, GC-MS, and IR, *Int J Pharm Pharm Sci.* **2014**, 6 (1), 675-686.
39. Karim, M.; Kalam, A.; Alam, A.; Alam K.; Jahan, N.; Jafri, A. Biskhapra (*Trianthema portulacastrum* Linn) and its medicinal utility mentioned in Unani System of Medicine—A Review, *IJPSR.* **2015**, 6(4), 790-795.
40. Geethalakshmi, R.; Sundaramurthi, J.; Sarada, L. Antibacterial activity of flavonoid isolated from *Trianthema decandra* against *Pseudomonas aeruginosa* and molecular docking study of FabZ. *Microb. Pathog.* **2018**, 121, 87-92.
41. Impellizzeri, G.; Piattelli, M.; Sciuto, S. Acylated betacyanins from *Drosanthemum floribundum*, *Phytochemistry.* **1973**, 12, 2295-2296.
42. Impellizzeri, G.; Piattelli, M.; Sciuto, S. A New betaxanthin from *glottiphyllum longum*, *Phytochemistry.* **1973**, 12, 2293-2294.
43. Piattelli, M. Impellizzeri, G. Betacyanins from *lampranthus* sp. (aizoaceae), *Phytochemistry.* **1969**, 8, 1595-1596.
44. Strack, D.; Bokern, M.; Marxen, N.; Wray, V. Feruloylbetanin from petals of *Lampranthus* and feruloylamaranthin from cell suspension cultures of *Chenopodium rubrum*, *Phytochemistry.* **1988**, 27(11), 3529-3531.
45. Magwa, L.; Gundidza, M.; Gwerua, N.; Humphrey, G. Chemical composition and biological activities of essential oil from the leaves of *Sesuvium portulacastrum*, *J. Ethnopharmacol.* **2005**, 103, 85-89.
46. Veeresh; Kumar, P.; Lavanya, A.; Suresh, P.; Mounika, N. Pharmacological potentials of *Trianthema decandra*- A review, *J Pharmacogn Phytochem.* **2017**, 6(2), 290-294.
47. Gagare, B.; Jadhav, S. Comparative Phytochemical profiling of various extracts, from different parts of *Sesuvium portulacastrum* using GCMS, FTIR and ICP AES, *IJESM.* **2017**, 6(6), 267-276.
48. El-Amier, A.; Haroun, A.; El-Shehaby, O.; Al-hadithy, N. Antioxidant and Antimicrobial Properties of Some Wild Aizoaceae Species Growing in Egyptian Desert, *J. Environ. Sci.* **2016**, 5 (1), 1-10.
49. Doudach, L.; Meddah, B.; Benbacer, L.; Hammani, K.; El mzibri, M.; Elomri, A.; Cherrah, Y. *Ethnopharmacological studies of Mesembryanthemum nodiflorum*, *Pharmacology.* **2013**, 4(2), 246-258.
50. Mohammed, R.; El-Hawary, S.; Abo-youssef, M. Biological investigation of some wild *Aizoaceae* and *Chenopodiaceae* species growing in Egypt, *J. Nat. Prod.* **2012**, 5, 193-206.
51. Qasem, R. Prospects of wild medicinal and industrial plants of saline habitats in the Jordan valley, *Pak. J. Bot.* **2015**, 47 (2), 551-570.
52. Abdallah, A.; Ali Merito, A.; Hassan, S.; Aboubaker, D.; Djama, M.; Asfaw, Z.; Kelbessa, E. Medicinal plants and their uses by the people in the Region of Randa, Djibouti, *J. Ethnopharmacol.* **2013**, 148, 701-713.
53. Stark, D.; Mtui, J.; Balemba, B. Ethnopharmacological survey of plants used in the traditional treatment of gastrointestinal pain, inflammation and diarrhea in Africa: future perspectives for integration into modern medicine. *Animals* **2013**, 3, 158-227.
54. Omoruyi, E.; Bradley, G.; Afolayan, J. Antioxidant and phytochemical properties of *Carpobrotus edulis* (L.) bolus leaf used for the management of common infections in HIV/AIDS patients in Eastern Cape Province, *BMC Complem ALtern M.* **2012**, 12, 215.
55. Van Wyk, E.; Van Oudtshoorn, B.; Gericke, N. Medicinal Plants of South Africa. *Briza Publications*, Pretoria, ISBN 1-875093-09-5, **1997**.
56. De Ghosh, M.; Ramakrishna, S.; Ramakrishna, M. Antimicrobial activity and phytochemical analysis of medicinal plants, *WJPPS.* **2014**, 3(6), 1794-1799.
57. Moyo, B.; Mukanganyama, S. The Anticandidal and Toxicity Properties of *Lampranthus francisci*, *J. Mycol.* **2015**, 2015, 1-15.
58. Van Wyk, E.; Gericke, N. People's Plants: A Guide to Useful Plants of Southern Africa, *Briza Publications*, Pretoria, ISBN 978-1-875093-s19-9, **2000**.
59. VanWyk, E. A broad review of commercially important southern African medicinal plants, *J. Ethnopharmacol.* **2008**, 119, 342-355.
60. Afzal, S.; Chaudhary, B.; Ahmad, A.; Afzal, K. Preliminary phytochemical analysis and antifungal activities of crude extracts of *Zaleya pentandra* and *Corchorus depressus* linn., *Acta Pol Pharm.* **2015**, 72, 329-334.
61. Afzal, S.; Chaudhry, B.; Afzal, K.; Saeed, J.; Hakash, S.; Qadir, M. Acetyl cholinesterase and butyryl cholinesterase inhibitory activities of *Zaleya pentandra*. *Acta Pol Pharm.* **2017**, 74 (3), 891-894.