#### BASIC PROBABILITY: THEORY

Master of Logic, University of Amsterdam, 2018 — LICENSE CC BY-NC-SA 4.0 TEACHERS Alexandre Cremers and Jakub Dotlačil TA Bas Cornelissen

# **Board questions set 4**

# **Problem 1: Variances**

- (a) Prove that if  $X \sim \text{Bernoulli}(p)$ , then Var(X) = p(1-p).
- **(b)** Prove that if  $X \sim \text{Bin}(n, p)$ , then Var(X) = np(1 p).
- (c) Suppose  $X_1, X_2, ..., X_n$  are independent and all have the same standard deviation  $\sigma$ . Let  $\overline{X}$  be the average of  $X_1, X_2, ..., X_n$ . What is the standard deviation of  $\overline{X}$ ? What does this mean?

## Problem 2: Covariance

- (a) Flip a fair coin 3 times. Let X be the number of heads in the first 2 flips and let Y be the number of heads in the last 2 flips. Give a table describing the joint distribution of X and Y and directly compute Cov(X,Y).
- **(b)** Let  $X_1, X_2, X_3$  be the results of the three fair coin flips and let X and Y as before. Compute Cov(X, Y) without first using the joint distribution.

### Problem 3: More covariance

Toss a fair coin 2n+t times. Let X be number of heads in the first n+t flips and let Y be number of heads in the last n+t flips. Compute Cov(X,Y) and Cor(X,Y).

#### BASIC PROBABILITY: THEORY

Master of Logic, University of Amsterdam, 2018 — LICENSE CC BY-NC-SA 4.0 TEACHERS Alexandre Cremers and Takub Dotlačil TA Bas Cornelissen

# **Board questions set 4**

## **Problem 1: Variances**

- (a) Prove that if  $X \sim \text{Bernoulli}(p)$ , then Var(X) = p(1-p).
- **(b)** Prove that if  $X \sim \text{Bin}(n, p)$ , then Var(X) = np(1 p).
- (c) Suppose  $X_1, X_2, ..., X_n$  are independent and all have the same standard deviation  $\sigma$ . Let  $\overline{X}$  be the average of  $X_1, X_2, ..., X_n$ . What is the standard deviation of  $\overline{X}$ ? What does this mean?

## Problem 2: Covariance

- (a) Flip a fair coin 3 times. Let X be the number of heads in the first 2 flips and let Y be the number of heads in the last 2 flips. Give a table describing the joint distribution of X and Y and directly compute Cov(X,Y).
- **(b)** Let  $X_1, X_2, X_3$  be the results of the three fair coin flips and let X and Y as before. Compute Cov(X,Y) without first using the joint distribution.

### Problem 3: More covariance

Toss a fair coin 2n + t times. Let X be number of heads in the first n + t flips and let Y be number of heads in the last n + t flips. Compute Cov(X, Y) and Cor(X, Y).