

Homework III

Advanced Topics in Optimization

Basil R. Yap

2018 February 16

1 Question 1

system description: $1|prec|h_{\max}$

jobs	1	2	3	4	5	6	7
p_j	4	8	12	7	6	9	9
$h_j(C_j)$	$3C_1$	77	C_3^2	$1.5C_4$	$70 + \sqrt{C_5}$	$1.6C_6$	$1.4C_7$

$$1 \rightarrow 7 \rightarrow 6$$

$$5 \rightarrow 7$$

$$5 \rightarrow 4$$

Set $J = \emptyset$, $J^c = \{1, 2, \dots, 7\}$, $J' = \{2, 3, 4, 6\}$

$$h_{j*} \left(\sum_{k \in J^c} p_k \right) = \min_{j \in J'} \left(h_j \left(\sum_{k \in J^c} p_k \right) \right)$$

$$\sum_{k \in J^c} p_k = 55$$

$$h_2 \left(\sum_{k \in J^c} p_k \right) = 77$$

$$h_3 \left(\sum_{k \in J^c} p_k \right) = (55)^2$$

$$= 3025$$

$$h_4 \left(\sum_{k \in J^c} p_k \right) = 1.5 \times 55$$

$$= 82.5$$

$$h_6 \left(\sum_{k \in J^c} p_k \right) = 1.6 \times 55$$

$$= 88$$

$$\min_{j \in J'} (h_j (\sum_{k \in J^c} p_k)) = 77$$

$$J = \{2\}, \quad J^c = \{1, 3, 4, 5, 6, 7\}, \quad J' = \{3, 4, 6\}$$

$$\sum_{k \in J^c} p_k = 47$$

$$\begin{aligned} h_3 \left(\sum_{k \in J^c} p_k \right) &= (47)^2 \\ &= 2209 \end{aligned}$$

$$\begin{aligned} h_4 \left(\sum_{k \in J^c} p_k \right) &= 1.5 \times 47 \\ &= 70.5 \end{aligned}$$

$$\begin{aligned} h_6 \left(\sum_{k \in J^c} p_k \right) &= 1.6 \times 47 \\ &= 75.2 \end{aligned}$$

$$\min_{j \in J'} (h_j (\sum_{k \in J^c} p_k)) = 70.5$$

$$\begin{aligned} J &= \{2, 4\}, \quad J^c = \{1, 3, 5, 6, 7\}, \quad J' = \{3, 6\} \\ \sum_{k \in J^c} p_k &= 40 \end{aligned}$$

$$\begin{aligned} h_3 \left(\sum_{k \in J^c} p_k \right) &= (40)^2 \\ &= 1600 \end{aligned}$$

$$\begin{aligned} h_6 \left(\sum_{k \in J^c} p_k \right) &= 1.6 \times 40 \\ &= 64 \end{aligned}$$

$$\min_{j \in J'} (h_j (\sum_{k \in J^c} p_k)) = 64$$

$$\begin{aligned} J &= \{2, 4, 6\}, \quad J^c = \{1, 3, 5, 7\}, \quad J' = \{3, 7\} \\ \sum_{k \in J^c} p_k &= 31 \end{aligned}$$

$$\begin{aligned} h_3 \left(\sum_{k \in J^c} p_k \right) &= (31)^2 \\ &= 961 \end{aligned}$$

$$\begin{aligned} h_7 \left(\sum_{k \in J^c} p_k \right) &= 1.4 \times 31 \\ &= 43.4 \end{aligned}$$

$$\min_{j \in J'} (h_j (\sum_{k \in J^c} p_k)) = 43.4$$

$$\begin{aligned} J &= \{2, 4, 6, 7\}, \quad J^c = \{1, 3, 5\}, \quad J' = \{1, 3, 5\} \\ \sum_{k \in J^c} p_k &= 22 \end{aligned}$$

$$h_1 \left(\sum_{k \in J^c} p_k \right) = 3 \times 22$$

$$= 66$$

$$h_3 \left(\sum_{k \in J^c} p_k \right) = (22)^2$$

$$= 484$$

$$h_5 \left(\sum_{k \in J^c} p_k \right) = 70 + \sqrt{22}$$

$$= 74.69$$

$$\min_{j \in J'} (h_j (\sum_{k \in J^c} p_k)) = 66$$

$$J = \{2, 4, 6, 7, 1\}, \quad J^c = \{3, 5\}, \quad J' = \{3, 5\}$$

$$\sum_{k \in J^c} p_k = 18$$

$$h_3 \left(\sum_{k \in J^c} p_k \right) = (18)^2$$

$$= 324$$

$$h_5 \left(\sum_{k \in J^c} p_k \right) = 70 + \sqrt{18}$$

$$= 74.24$$

$$\min_{j \in J'} (h_j (\sum_{k \in J^c} p_k)) = 74.24$$

$$J = \{2, 4, 6, 7, 1, 5, 3\}, \quad J^c = \emptyset, \quad J' = \emptyset$$

2 Question 2

system description: $1 || \sum T_j$

jobs	1	2	3	4	5	6	7	8
p_j	6	18	12	10	10	11	5	7
d_j	8	42	44	24	26	26	70	75

3 Question 3

Consider the problem $1|prec|\max(h_1(S_1), \dots, h_n(S_n))$, where S_j denotes the starting time of job j . The cost function h_j , $j = 1, \dots, n$ is decreasing. Unforced idleness of the machine is not allowed. Describe a dynamic programming type algorithm for this problem similar to the one in Section 3.2. Why does one have to use here forward dynamic programming instead of backward dynamic programming?

placeholder

4 Question 4

5 Question 5

(a)

(b)

(c)

6 Question 6

(a)

(b)

(c)

(d)

(e)

(f)